

## *International Journal of Scientific Research and Reviews*

### **Molecular Phylogenetic study of larvae (*Chrysomya megacephala*) recovered from human corpse during crime scene investigation (Motinagar Thana suicidal case) at Sagar (India)**

**Jangir Shyam Sundar<sup>1\*</sup>, Patidar Pankaj<sup>2</sup> and Sharma J.D.<sup>3</sup>**

<sup>1,3</sup>Department of Criminology and Forensic Science, Dr. H.S. Gour University, Sagar, Madhya Pradesh, India 470003

Email: [shyamjangircfs@gmail.com](mailto:shyamjangircfs@gmail.com), [jdsharma29@gmail.com](mailto:jdsharma29@gmail.com), Mob. No: 9424438034

<sup>2</sup>Department of Police and Crime Scene unit, State Forensic science laboratory, Sagar, Madhya Pradesh, India 470003, Email: [pan\\_kaj469@yahoo.co.in](mailto:pan_kaj469@yahoo.co.in), Mob. No: 9981765547

#### **ABSTRACT**

DNA based identification of larvae sampled from Motinagar, Thana suicidal case of Sagar district, Madhya Pradesh was successfully demonstrated in the present study. It was found that the recovered larvae from the crime scene were identified morphologically as *Chrysomya rufifacies* but later molecularly confirmed to be *Chrysomya megacephala* by phylogenetic tree. Result from BLAST system revealed a 98-99% identity with *Chrysomya megacephala*. Phylogenetic analysis confirmed the presence of *Chrysomya megacephala* with 98-99% identity (match). The molecular study involved the sequencing of a total length of 1.3 kb encompassing the Cytochrome Oxidase I (COI) gene. This study will be very helpful to rule out the post mortem interval (PMI) due to correct species identification which can constrict the field of suspects and aid in the identification of the decedent in the near future also in India.

**KEYWORDS:** *Chrysomya megacephala*, mitochondrial DNA, COI, sequencing

#### **\*Corresponding author:**

**Shyam Sundar Jangir,**

Department of Criminology and Forensic Science,

Dr. H.S. Gour University, Sagar, M.P.

Email-[shyamjangircfs@gmail.com](mailto:shyamjangircfs@gmail.com)

## INTRODUCTION

Medico-legal entomological evidence plays diverse role as a forensic marker in criminal trials. In particular, determination of minimum postmortem interval ( $PMI_{min}$ ) of a decomposed dead body, locations and sometimes in the manner of death, therefore, accurate species identification is necessary for exploring the full potential of insect evidence. Taxonomic identification has some limitations in the procedure of identification. Because of this, to ensure correct species identification, DNA based methods are now increasingly employed in the field of medico- legal entomology<sup>1, 2</sup>. Calliphoridae flies viz. *Chrysomya rufifacies* and *Chrysomya megacephala* are one of the first visitors on human corpse<sup>3</sup>. Now a day, mitochondrial DNA (mtDNA) is becoming a very useful tool in correct insect identification<sup>4 5</sup>. Mitochondrial DNA (mtDNA) provides several advantages over nuclear DNA; therefore mtDNA becomes a better tool to determine closely related species for DNA based identification<sup>6</sup>. In India, there are many limitations to access human dead body at the crime scene, because forensic entomology is not very much popular in forensic practices. This study focus to examine the full potential of molecular identification of the first witness, that is fly and fly larvae recovered from human cadavers to compare the findings with the taxonomic identification of calliphoridae.

## MATERIAL AND METHODS

### (A) *Insect collection and rearing*

Insect larvae were collected with the help of soft nose forcep and kept into small plastic jar on 09.08.2016 at real crime scene (Temperature (24°C Avg.), relative humidity (97% Avg.) under Motinagar Police Station (0/16) Sagar. Maggots were reared (Temp 26.2°C) on an artificial diet using rearing method of Hung<sup>7</sup>. Emerge adult flies were used for taxonomic and molecular identification larvae were used for molecular identification (preserved in 90% ethanol).

### (B) *DNA extraction*

Genomic DNA from whole fly and whole larvae were extracted using the DNeasy tissue kit (Qiagen, Germany) following the manufacturer's instructions. For fly and tissue samples, 180ul buffer ATL was used separately, added 20ul proteinase K in each vial, mixed by vortexing and incubated at 56°C until completely lysed. Added 200ul of buffer AL and mixed thoroughly by vortexing then added 200ul of ethanol (96-100%) and again mixed. The mixture was transferred into a DNeasy mini spin column placed in a 2 ml collection tube and centrifuged at 8000 rpm for 1 min. Flow was discarded and spin was placed in a new 2 ml collection tube followed by 500 ul buffer AWZ and centrifuged at 14000 rpm for 3 min. Flow was again discarded and spin was transferred

into a new 1.5 ml MCT and DNA was eluted by adding 200 ul buffer AE and incubated for 1 min at room temperature (20°C) and finally centrifuged at 8000 rpm for 1 min. Extracted DNA was checked electrophoretically on 1% agarose gel (Fig 1). The fraction of extracted DNA was spectrophoretically quantitated and diluted to 50ng/μL prior to PCR amplification step.




Figure 1: Genomic DNA Loaded in 1% Agarose Gel. Lane 1: Larvae genomic DNA. 2: adult fly (*Chrysomya megacephala*) genomic DNA.

### (C) PCR amplification, sequencing and Phylogenetic analysis


Two opposing primers were selected for the PCR of the gene of Cytochrome Oxidase subunit 1 (COI) from published nucleotide sequences (Accession no. AY842621, AY842622, AY842623) and purchased from Chromous Biotech, Bangalore. The primers pairs composed of forward primer (Named C1-J-1751) of 5'GGAGCTCCTGACATAGCATTCCC3' and the reverse primer (Named UEA10) of 5'TCCAATGCACTAATCTGCCACATTA3' was designed to amplify a ~1.3 Kilo base pair of COI specific fragmented by PCR (Table 1). The PCR amplification mixture were prepared by containing the following components viz. 1X PCR reaction buffer, MgCl<sub>2</sub>, dNTPs, template DNA (100 ng), Taq Polymerase and Forward & Reverse primers each. Amplification reactions were performed in a Thermal Cycler (Applied Biosystem). PCR cycling conditions were as follows: initial denaturation at 96°C for 1 min, 25 cycles of denaturation at 96°C for 10 sec, hybridization at 50°C for 5 sec and followed by elongation at 60°C for 4 min. It was found that optimal annealing/hybridization temperature was 50°C for COI region. The PCR product were separated electrophoretically on 1% agarose gel and visualized after ethidium bromide staining (Fig 2).

Table 1: Detail of PCR primers

S.No.	Location	Sequence	Reference
1.	CJ-J-1751	5'GGAGCTCCTGACATAGCATTCCC3'	8
2.	UEA10	5'TCCAATGCACTAATCTGCCACATTA3'	9

DNA sequencing was carried out using an ABI 3500XL Genetic Analyser Big Dye Terminator Version 3.1' cycle sequencing kit, POP\_7 polymer & 50 cm Capillary array as polymer and capillary array, BDTv3-KB-Denovo\_v 5.2 as analysis protocol and Seq Scape\_ v 5.2 as data analysis software. The sequencing mixture containing the following components viz. Big Dye Terminator Ready Reaction Mix (4μL), Template (100ng/ul), Primer (10pmol/λ), Milli Q Water (3μL). The sequencing was determined for both forward and reversed strands. Sequence alignment

and phylogenetic tree by Maximum Likelihood Method were made using MEGA 7, bootstrap support derived from 5000 replicates. Aligned sequence of COI of *Chrysomya megacephala* after sequencing is shown in fig 3. The reference sequences of previously reported work on *Chrysomya megacephala* identification are shown in Table 2.



**Figure 2: PCR Product loaded on 1% Agarose Gel. Lane description: L: 500 bp ladder, 1: *Chrysomya megacephala* (~1.3kb)**

```

CCGTAGTCATATAACTTTATTATTAGTAAGTAGTATAGTAGAAAATGGGGCTGGAACAGGATGAACTGTTTACCCACCT
TTATCTTCTAATATTGCTCATGGAGGAGCATCAGTTGATTTAGCTATTTTCTCTTTACACTTAGCAGGAATTTCTTCAATT
TTAGGAGCTGTAAATTTTATTACAACGTAAATTAATATACGATCTACAGGAATTACATTTGATCGAATACCTTTATTTGT
ATGATCTGTAGTTATTACTGCTCTATTATTATTATTATCTTTACCAGTATTAGCTGGAGCTATTACTATATTATTAAGTGA
CCGAAATCTAAATACTTCATTCTTTGACCCAGCAGGAGGAGGAGATCCTATTTTATACCAACATTTATTTGATTCTTTG
GACATCCTGAAGTTTATATTTAATTTTACCTGGATTCGGAATAATTTCTCATATTATTAGTCAAGAATCAGGAAAGAAG
GAAACTTTCGGATCTTTAGGAATGATTTATGCTATACTAGCTATTGGTCTATTAGGATTTATTGTATGAGCTCACCACAT
GTTTACTGTTGGAATAGACGTAGACACACGAGCTTATTTCACTTCAGCTACATAATTATTGCTGTACCAACTGGAATTAA
GATTTTCAGTTGATTAGCAACTCTTTACGGAACACAATTAATTATTCTCCAGCTACTTTATGAGCTTTAGGATTTGTATT
TTTATTTACTGTAGGAGGATTAAGTGGAGTTGTTTATAGCTAATTCATCATTGACATTATTTACATGATACATATATGTA
GTAGCTCACTTCCATTATGTTCTATCATGGGAGCTGTATTTGCTATTATAGCAGGATTTGTTTCATTGATCCCTCTATTTA
CTGGATTAACCTTTAAATAGCAAGTTATTAAGAGTCAATTTGCTATTATATTTATCGGAGTAAATTTAACATTCTTCCCT
CAACATTTCTTAGGATTAGCAGGTATACCTCGACGATACTCAGACTATCCAGACGCTTACACAGCTTGAAATGTAATTT
CTACAATTGGTTCAACAATTTCAATTATTAGGAATTTATTCTTCTTTTTTCAATTATTGAGAAAGTTTAGTATCTCAACGAC
GAGTTTTATTCCCTGTTCCAAATAAATTCTATCTAATTGAATGATTAAGTAA
    
```

**Figure 3: Aligned sequence of *Chrysomya megacephala* (~1.3kb)**

**Table 2: Detail of NCBI GeneBank accessions (mtCOI) analysed**

Species	Accession number	Locality	Reference
<i>C. megacephala</i>	AF295551	USA	<sup>10</sup>
<i>C. megacephala</i>	AB112830	Australia	<sup>11</sup>
<i>C. megacephala</i>	AB112841	Australia	<sup>11</sup>
<i>C. megacephala</i>	AB112846	Australia	<sup>11</sup>
<i>C. megacephala</i>	AB112847	Australia	<sup>11</sup>
<i>C. megacephala</i>	AB112848	Australia	<sup>11</sup>
<i>C. megacephala</i>	AB112861	Australia	<sup>11</sup>
<i>C. megacephala</i>	AY092761	Taiwan	<sup>12</sup>
<i>C. megacephala</i>	AY842619	Australia	<sup>13</sup>
<i>C. megacephala</i>	AY909053	Malaysia	<sup>14</sup>
<i>C. megacephala</i>	FJ153258	Thailand	<sup>15</sup>
<i>C. megacephala</i>	FJ614818	China	<sup>16</sup>
<i>C. megacephala</i>	FJ614817	China	<sup>16</sup>
<i>C. megacephala</i>	FJ614816	China	<sup>16</sup>
<i>C. megacephala</i>	GQ912669	Pakistan	<sup>16</sup>
<i>C. megacephala</i>	KC249623	China	<sup>17</sup>
<i>C. megacephala</i>	KC249624	China	<sup>17</sup>
<i>C. megacephala</i>	KC249625	Egypt	<sup>17</sup>
<i>C. megacephala</i>	KC249626	Egypt	<sup>17</sup>
<i>C. megacephala</i>	KC249673	China	<sup>17</sup>
<i>C. megacephala</i>	KC249674	China	<sup>17</sup>
<i>C. megacephala</i>	KC249675	Egypt	<sup>17</sup>

**Table 3: Detail of specimens sequenced**

Species	Location	Police station	Date of collection	Methods and duration of preservation
<i>C. megacephala</i>	Sagar (India) (23°5'N, 78°5'E)	Motinagar (0/16)	09 Aug 2016	F, More than 10 month

Note: F, Frozen

## RESULT AND DISCUSSION

In this study, a suicide case of Motinagar, Thana of Sagar district, Madhya Pradesh was investigated (Fig 4). A case where the body was recovered from the closed room and declared as a suicide case after the investigation, during investigation of crime scene, entomological evidence as larvae were collected from the body as well as from the crime scene and the habitat was urban site, indoor with cloths on the body. The molecular identification of mitochondrial DNA (mtDNA) region was studied encompassing the Cytochrome oxidase subunit 1 gene (COI) gene. In this study, the complete 1.3 Kb nucleotide fragment was amplified (fig 3) and the phylogenetic analysis of collected larvae is shown in Fig 5. The detail of location, police station, date of collection and method & duration of preservation of *Chrysomya megacephala* are presented in table 3. The evolutionary history was inferred by using the Maximum Likelihood method based on the Kimura 2-parameter model <sup>18</sup>. The bootstrap consensus tree inferred from 5000 replicates <sup>19</sup> is taken to represent the evolutionary history of the taxa analyzed <sup>19</sup>. Branches corresponding to partitions reproduced in less than 50% bootstrap replicates are collapsed. The percentage of replicate trees in which the associated

taxa clustered together in the bootstrap test (5000 replicates) are shown next to the branches<sup>19</sup>. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. The analysis involved 100 nucleotide sequences. Codon positions included were 1st+2nd+3rd+Noncoding. All positions containing gaps and missing data were eliminated. There were a total of 536 positions in the final dataset. Evolutionary analyses were conducted in MEGA7<sup>20</sup>.

At motinagar thana case study, the recovered larvae were identified morphologically as *Chrysomya rufifacies* but later molecularly confirmed to be *Chrysomya megacephala* by phylogenetic tree. Result from BLAST system revealed a 98-99% identity with *Chrysomya megacephala*. Phylogenetic analysis confirmed the presence of *Chrysomya megacephala* with 98-99% identity (match). Out of 100 subject sequences, 95 sequences were of *Chrysomya megacephala* only and rest 5 other *Chrysomya* sp. sequences viz. *Chrysomya putoria* (98% identity), *Chrysomya cabrerai* (98% identity), *Chrysomya pacifica* (98% identity), *Chrysomya saffrana* (98% identity) and *Chrysomya chain* (98% identity) was also found based on phylogenetic tree analysis.

Hence it was found that the larvae recovered from human corpse during crime scene investigation were of *Chrysomya megacephala* larvae and it was also evident that the blow (Calliphoridae) flies viz. *Chrysomya rufifacies* and *Chrysomya megacephala* were first to visit on human corpse<sup>3</sup>. Forensic insect analysis during crime case provides valuable information by calculating Post mortem Interval (PMI) of larvae and insects found on human dead body and narrowing the investigation process.



Motinagar thana case -indoor with cloths

**Figure 4: Dead body recovered from crime scene investigation (Motinagar, Thana case) in Sagar, M.P**




Figure 5: Maximum Likelihood tree illustrating the presence of *Chrysomya megacephala* and phylogenetic relationships among blow flies (family: Calliphoridae) recovered from crime scene investigations based on COI gene of 1.3 Kb

## CONCLUSION

The current study successfully identified the collected larvae during crime scene investigation of Motinagar Thana case at Sagar, M.P. The entomological evidence collected was identified as

*Chrysomya megacephala* by DNA based analysis. This will further prove valuable in estimating the PMI in forensic investigation.

## ACKNOWLEDGEMENT

Authors thanks to University grants commission (UGC) for financial assistance, Dr. Devasish Bose (HOD, Department of Criminology and Forensic science, Dr. H.S Gour Central University, Sagar, M.P) for providing necessary facilities for the research work and Mr. Sachin Atulakar (Police Superident, Sagar) to provide facility to visit and collect entomological evidence at real Crime Scene.

## INTEREST OF CONFLICT

Author declares no conflict of interest.

## REFERENCES

1. Sperling, F. A.; Anderson, G. S.; Hickey, D. A., A DNA-based approach to the identification of insect species used for postmortem interval estimation. *Journal of Forensic Science* **1994**; 39: (2): 418-427.
2. Benecke, M., Random amplified polymorphic DNA (RAPD) typing of necrophagous insects (Diptera, Coleoptera) in criminal forensic studies: validation and use in practice. *Forensic Science International* **1998**; 98(3): 157-168.
3. Benecke, M., Arthropods and corpses. In *Forensic pathology reviews*, Springer: 2005; 207-240.
4. Harvey, M.; Gaudieri, S.; Villet, M.; Dadour, I., A global study of forensically significant calliphorids: implications for identification. *Forensic science international* **2008**; 177 (1): 66-76.
5. Wells, J. D.; Williams, D. W., Validation of a DNA-based method for identifying Chrysomyinae (Diptera: Calliphoridae) used in a death investigation. *International Journal of Legal Medicine* **2007**; 121 (1): 1.
6. Waugh, J., DNA barcoding in animal species: progress, potential and pitfalls. *BioEssays* **2007**; 29(2): 188-197.
7. Hung, T., The life table and mass rearing of *Chrysomya megacephala* (Fabricius). *Graduate Institute of Plant Pathology and Entomology, National Taiwan University, Taipei, Taiwan* **1995**.
8. Simon, C.; Frati, F.; Beckenbach, A.; Crespi, B.; Liu, H.; Flook, P., Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the entomological Society of America* **1994**; 87: (6): 651-701.
9. Lunt, D.; Zhang, D. X.; Szymura, J.; Hewlitt, O., The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies. *Insect molecular biology* **1996**; 5(3): 153-165.

10. Wells, J. D.; Sperling, F. A., DNA-based identification of forensically important Chrysomyinae (Diptera: Calliphoridae). *Forensic Science International* **2001**; 120(1): 110-115.
11. Harvey, M.; Mansell, M.; Villet, M.; Dadour, I., Molecular identification of some forensically important blowflies of southern Africa and Australia. *Medical and veterinary entomology* **2003**; 17(4): 363-369.
12. Chen, W.-Y.; Hung, T.-H.; Shiao, S.-F., Molecular identification of forensically important blow fly species (Diptera: Calliphoridae) in Taiwan. *Journal of Medical Entomology* **2004**; 41 (1):47-57.
13. Wallman, J. F.; Leys, R.; Hogendoorn, K., Molecular systematics of Australian carrion-breeding blowflies (Diptera: Calliphoridae) based on mitochondrial DNA. *Invertebrate Systematics* **2005**; 19(1): 1-15.
14. Tan, S. H.; Aris, E. M.; Surin, J.; Omar, B.; Kurahashi, H.; Mohamed, Z., Sequence variation in the cytochrome oxidase subunit I and II genes of two commonly found blow fly species, *Chrysomya megacephala* (Fabricius) and *Chrysomya rufifacies* (Macquart)(Diptera: Calliphoridae) in Malaysia. *Trop. Biomed* **2009**; 26: 173-181.
15. Preativatanyou, K.; Sirisup, N.; Payungporn, S.; Poovorawan, Y.; Thavara, U.; Tawatsin, A.; Sungpradit, S.; Siriyasatien, P., Mitochondrial DNA-based identification of some forensically important blowflies in Thailand. *Forensic science international* **2010**; 202 (1): 97-101.
16. Zaidi, F.; Wei, S.-J.; Shi, M.; Chen, X.-X., Utility of multi-gene loci for forensic species diagnosis of blowflies. *Journal of Insect Science* **2011**; 11;(1), 59.
17. Aly, S. M., Reliability of long vs short COI markers in identification of forensically important flies. *Croatian medical journal* **2014**; 55(1): 19-26.
18. Kimura, M., A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of molecular evolution* **1980**; 16(2):111-120.
19. Felsenstein, J., Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **1985**; 783-791.
20. Kumar, S.; Stecher, G.; Tamura, K., MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular biology and evolution* **2016**; 054.