

## *International Journal of Scientific Research and Reviews*

### **Stability Studies of Proteases Engendered From Enticing Microbial Primogenitors Screened From Estuarine Mangrove Detritus**

**Talreja Sarika Chhabria<sup>1\*</sup> and Chandra B. Maurya<sup>2</sup>**

<sup>1\*</sup> Guru Nanak Khalsa College, Matunga, Mumbai 400 019

<sup>2</sup> KLE College of Science and Commerce, Kalamboli, Navi Mumbai 410 218

E-mail: [sarikachhabriatalreja@gmail.com](mailto:sarikachhabriatalreja@gmail.com)

#### **ABSTRACT**

Marshy mangrove areas of Goa, India are opulent incumbents of halophilic bacteria resplendent in variety of Proteases and huge salinity-stress tolerant capacity. Bacteria are the most prevalent consortia of Protease primogenitors serving as quintessential enzyme source; with *Bacillus* being the premier genus. The colossal heterogeneity of Proteases, coupled with the ease with which they can be genetically manoeuvred with revamped properties desirable for diverse industrial applications, in contradistinction to their modus operandi and explicitness, has captivated worldwide recognition to exploit their commercial utility.

The present study is an attempt to harness bacterial isolates from Goa's mangrove areas for some neoteric proteolytic enzymes. Out of twenty isolates screened, B<sub>2</sub> and B<sub>3</sub> manifested persuasive proteolytic activity. Phylogenetic analysis further ensued that isolates were related to *Bacillus* Sp. Optimization studies by growth media variation revealed that optimal growth media temperature and pH was 40°C and 7.0 respectively. With respect to the characteristics influencing culture conditions and properties, it was considered sententious to characterize the enzyme through Stability studies. Studies showed that Protease was considerably stable at optimum conditions of 60°C in pH 7.0.

**KEY WORDS:** *Bacillus*, Characterization, Mangroves, Proteases, Stability

#### **\*Corresponding author**

**Sarika Chhabria Talreja**

Research Scholar,

Guru Nanak Khalsa College,

Matunga, Mumbai 400 019

E-mail: [sarikachhabriatalreja@gmail.com](mailto:sarikachhabriatalreja@gmail.com)

## **1. INTRODUCTION**

Proteases form a bulky cluster of enzymes which are ubiquitous in nature and the central category enzymes from industrial view point. Bacteria are the most prevalent consortia of Protease primogenitors serving as the quintessential enzyme source; with *Bacillus* being the premier genus<sup>1,2</sup>. Their rapid growth, limited space required for cultivation<sup>3</sup> and the ease with which they can be genetically manoeuvred with revamped properties desirable for diverse industrial applications makes microbial Proteases commercially conspicuous<sup>4</sup>.

The colossal heterogeneity of Proteases, in contradistinction to their *modus operandi* and explicitness, has captivated worldwide recognition to exploit their physiological and biotechnological relevance<sup>5,6</sup>. They are one of the most indispensable categories of industrial enzymes employed in brewing, meat and photography industries. They are rampantly exploited in food and pharmaceutical industry for peptide synthesis and leather industry for dehairing. Their application in detergent industry as additives for surfactant formulation<sup>1,7</sup> so as to replace currently used noxious chemicals, is a relatively recent advancement and has conferred added biotechnological significance<sup>6</sup>.

Proteases have traditionally detained a predominant share of industrial enzyme market accounting for about 60-65% of total global sale of enzymes<sup>6,8,9,10</sup>. About 35% of the aggregate microbial enzymes utilized in detergent industry are procured from bacterial sources and majority of them are produced by *Bacillus* Sp.<sup>11,12</sup>.

In this paper, we aimed to investigate the stability at varied pH and temperature conditions<sup>13</sup> of neutral Proteases extracted from *Bacillus safensis* for its possible and potential applications for industrial purposes.

## **2. EXPERIMENTAL SECTION**

Twenty Mangrove sediment samples (B<sub>1</sub> - B<sub>20</sub>) were collected from Panji, Goa, India and screened for their morphological and Protease production efficiencies. Out of 20, B<sub>2</sub> and B<sub>3</sub> were found to be competent cultivable source of Proteases (as described in our previous publication by Sarika et. al 2017). Protease activity was measured using Caseinolytic assay. Two media containing Casein and Gelatin as the protein substrate were studied for two selected positive cultures B<sub>2</sub> and B<sub>3</sub>. Growth curve studies revealed that the culture entered late log phase after 24h and entered stationary phase after 48h. Maximum Protease activity was obtained after 24h of incubation. The results significantly indicated that B<sub>3</sub> exhibited maximum Protease activity in lesser fermentation time which is a major prerequisite for commercial Protease production. The filtered enzyme supernatant was concentrated 10 fold through Tangential Flow Filtration method and used for further studies. Growth Optimization studies showed that the media containing specific protein substrate gave higher

activity than the basic protein mixture media. Growth media variation studies divulged that the optimal pH and temperature for growth medium was 7.0 and 40°C respectively. With respect to the characteristics influencing culture conditions, productivity and properties of Protease, it was considered sententious to purify and characterize the enzyme through its stability studies.

## **2.1 Stability Studies<sup>14</sup>**

Test buffer used was Sodium phosphate buffer for pH 7 and 10.

**2.1.1 Stability at optimum conditions:** To analyze the pH stability, the Protease that was obtained from TFF concentration method was diluted with the pH 7 buffer and incubated at 60°C for 1h. Enzyme aliquot was taken after every 15 minutes and a reaction was run with Casein.

**2.1.2 Stability at extreme conditions:** To analyze the pH and temperature stability, TFF concentrated Protease was diluted with the buffer of pH 10 and incubated at 80°C for 15 minutes. Enzyme aliquot was taken after every 5 minutes and a reaction was run with Casein.

**2.1.3 Stability at extreme pH and optimum temperature:** To analyze the pH stability, TFF concentrated Protease was diluted with the buffer of pH 10 and incubated at 60°C for 1 hour. Enzyme aliquot was taken after every 15 minutes and a reaction was run with Casein.

**2.1.4 Stability at extreme temperature and optimum pH:** To analyze the temperature stability, TFF concentrated Protease was diluted with the buffer of pH 7 and incubated at 80°C for 15 minutes. Enzyme aliquot was taken after every 5 minutes and a reaction was run with Casein.

## **3. RESULTS AND DISCUSSION**

### **3.1 Stability Studies**

100% activity shown in the data was obtained from enzyme activity measured at 60°C in pH 7.0 and other residual activities were measured accordingly. Activity at extreme conditions reduced to negligible i.e. 1U/ml within 15 minutes hence was not continued for 1h. Protease was considerably stable for 1h at optimum conditions though there was a decrease in the activity observed (**Table 1**) (**Figures 1, 2, 3, 4**).

Hideto Takami, 1998, extracted extremely thermostable Protease from an alkalophilic *Bacillus* sp. No. AH-101 and found that the enzyme production reached its maximum level of 1500 units/ml after about 24 hours in alkaline medium at pH 9.5<sup>[15]</sup>. Similarly, Enshasy H., 2008, studied extracellular alkaline Protease of *Bacillus licheniformis* and said that in optimized medium conditions, maximum Protease production was estimated to be 2400 UmL<sup>-1</sup> after 96 hours<sup>[16]</sup>. Joshi et al, 2011, also found that Protease was extracted from *Bacillus cereus* MTCC 6840 and noted that Protease production was found to be 120U/mL<sup>[17]</sup>. Ganesh Kumar et al 2004 also found the optimal pH and temperature for alkaline Protease extracted from *Bacillus clausii* were 11.5 and 80°C<sup>[18]</sup>.

Subba Rao et al, 2009 characterised the thermo and detergent stable serine alkaline protease isolated from *Bacillus circulans* that exhibited optimum activity at broad temperature range of maximum of 70°C under alkaline pH environment<sup>19,20</sup>.

**TABLES AND FIGURES**

**Table 1: Time v/s Enzyme Activity in U/ml at 60°C and pH 7**

Duration (minutes)	Blank Absorbance	Test Absorbance	Test - Blank	µmoles/ml	U/ml
0	0.434	0.227	0.207	69.00	207.00
15	0.312	0.231	0.081	27.00	81.00
30	0.31	0.237	0.073	24.33	73.00
45	0.304	0.232	0.072	24.00	72.00
60	0.323	0.252	0.071	23.67	71.00

**Table 2: Time v/s Enzyme Activity in U/ml at 60°C and pH 10**

Duration (minutes)	Blank Absorbance	Test Absorbance	Test - Blank	µmoles/ml	U/ml
0	0.45	0.202	0.248	82.67	99.20
15	0.311	0.237	0.074	24.67	74.00
30	0.295	0.232	0.063	21.00	63.00
45	0.284	0.242	0.042	14.00	42.00
60	0.257	0.252	0.005	1.67	5.00

**Table 3: Time v/s Enzyme Activity in U/ml at 80°C and pH 7**

Duration (minutes)	Blank Absorbance	Test Absorbance	Test - Blank	µmoles/ml	U/ml
0	0.4	0.202	0.198	66	79.20
5	0.35	0.21	0.14	46.67	56.00
10	0.276	0.24	0.036	12.00	24.00
15	0.232	0.23	0.002	0.67	1.33

**Table 4: Time v/s Enzyme Activity in U/ml at 80°C and pH 10**



Duration (minutes)	Blank Absorbance	Test Absorbance	Test - Blank	µmoles/ml	U/ml
0	0.40	0.202	0.198	66	79.20
5	0.344	0.29	0.054	18.00	21.60
10	0.294	0.251	0.043	14.33	17.20
15	0.254	0.252	0.002	0.67	0.80


Table 5: Statistical Analysis (SPSS v. 22) of Enzyme Stability Studies indicated the following results for each set of data:

Correlation: Spearman's Method using SPSS v. 22				
Table 1: Time v/s Enzyme Activity in U/ml				
1. 60°C and pH 7		3. 60°C and pH 10		
2. 80°C and pH 7		4. 80°C and pH 10		
		Time	U/ml	
Spearman's rho	Time	Correlation Coefficient	1.000	-1.000**
		Sig. (2-tailed)	.	.
		N	5	5
	U/ml	Correlation Coefficient	-1.000**	1.000
		Sig. (2-tailed)	.	.
		N	5	5
**. Correlation is significant at the 0.01 level (2-tailed).				

Statistical Interpretation:

- Time of the test and Enzyme activity have a statistically **significant linear relationship** ( $r = -1.00$ ) in each of the temperature and pH conditions viz. 60°C and pH 7, 60°C and pH 10, 80°C and pH 7 as well as 80°C and pH 10.
- The direction of relationship, for each condition of temperature and pH, is **negative**. The variables time and enzyme activity move in different directions. As time increases, enzyme activity decreases and vice-versa. It is concluded that Time and enzyme activity are **negatively correlated**.
- The magnitude or strength of the association in each of the temperature and pH condition is **highly strong**.





#### 4. CONCLUSION

The escalated cognizance of biocatalytic potentialities of enzymes has made feasible the progression of novel biological products in diverse industrial spheres. Exploration of new cost effective process for enzyme production by exploiting various novel sources is the basic need of today in the current competitive enzyme market. Additionally, the enzyme thermostability confers numerous advantages viz. enhanced reaction velocity, prolonged half life, diminished contamination menace and decrease in viscosity permitting higher concentration of less dissolvable materials.

In conclusion, the present work describes Protease production by moderately halophilic *Bacillus safensis*. The utilization of cheap substrate for enzyme production and higher purification fold with easy protocols could make this enzyme economically feasible. Collectively, the results justify the suitability of *Bacillus safensis* sp. for commercial production of Protease, using inexpensive materials. Some of the features of the secreted Proteases by this bacterial specie, such as optimum neutral pH (7.0), reasonable activity at high temperature up to 60°C and compatibility characteristics are credentials to future studies.

## ACKNOWLEDGEMENTS

Laboratory support from Principals, Smt. C.H.M College and Guru Nanak Khalsa College, Mumbai is gratefully acknowledged. Author is also thankful to University Grants Commission for sanctioning study leave under FDP Scheme.

## REFERENCES

1. Gupta R., Beg Q. and Lorenz P. Bacterial alkaline proteases; molecular approaches and industrial applications. *Applied Microbiology and Biotechnology* 2002; **59** (1):15 - 32.
2. Ramakrishna D.P.N., Gopi Reddy N., Rajgopal S.V. Purification and Properties of an extracellular alkaline protease produced by *Bacillus subtilis* (MTCC No. 10110), *International Journal of Biotechnology and Biochemistry*. 2010; **6**(4): 489 - 500.
3. Arulmani Manavalan, Aparanjini Kalaichelvan, Vasanthi Kalyanasundaram, Arumugam Perumal, Arivuchelvi Manavalan and Kalaichelvan Thangavelu P. Purification and partial characterization of serine protease from thermostable alkalophilic *Bacillus laterosporus*-AK1. *World Journal of Microbiology and Biotechnology*. 2007; **23** (4): 475-481.
4. Beg, Q.K. and Gupta Rani Purification and characterization of an oxidation-stable, thiol-dependent alkaline protease from *Bacillus mojavensis*, *Enzyme & Microbial Technology*. 2003; **32** (2): 294-304.
5. Binod, P., Piyush P., Raghavendra G., Madhavan K.N., Arvind D., Kakali D. and Ashok P. Industrial Enzymes: Present Status and Future Perspectives for India, *Journal of Scientific & Industrial Research*. 2013; **272**: 271-286.
6. Rao, M.B., Tanksale A.M., Ghatge M.S. and Deshpande V.V. Molecular and Biotechnological Aspects of Microbial Proteases. *Microbiology and Molecular Biology Reviews*. 1998; **62**(3): 597-635.
7. Mohsen Fathi Najafi, Deobagkar Dileep and Deobagkar Deepti Potential application of Protease isolated from *Pseudomonas aeruginosa* PD100. *Electronic Journal of Biotechnology*, 2005; **8**(2):198-207.
8. Banerjee, U.C., Sani R.K., Azmi W. and Soni R. Thermostable Alkaline Protease from *Bacillus brevis* and its Characterization as a laundry detergent additive. *Process Biochemistry*, 1999; **35**: 213-219.
9. Adinarayana, K., Bapi Raju K.V.V.S.N. and Ellaiah, P. Investigation on alkaline protease production with *B. subtilis* PE11 immobilized in calcium alginate gel beads. *Process Biochemistry* 2004; **39**(11): 1331-1339.

10. Adinarayana, K., Ellaiah P. and Prasad D.S. 2003 Purification and partial characterization of thermostable serine alkaline Protease from a newly isolated *Bacillus subtilis* PE-11. *AAPSP PharmSciTech* 4 (4), Article 56.
  11. Ferrero M.A., Castro G.R., Abate C.M., Baigori M.D. and Sineriz F. Thermostable alkaline Protease of *Bacillus licheniformis* MIR 29: isolation, production and characterization. *Applied Microbiology and Biotechnology*. 1996; **45**(3): 327-332.
  12. Ellaiah, P., Adinarayana, K., Rajyalaxmi P. and Srinivasalu B. Optimization of process parameters for alkaline protease production under solid state fermentation by alkalophilic *Bacillus* sp., *Asian Journal of Microbiology, Biotechnology and Environmental science* 20036; **5**: 49-54.?
  13. Dutta J.R. and Banerjee R. Isolation and Characterization of a Newly Isolated *Pseudomonas* Mutant for Protease Production, *Brazilian Archives of Biology and Technology*, 2006; **49**(1): 37-47.
  14. Ghorbel Basma, Alya Sellami-Kamoun, Moncef Nasri Stability studies of Protease from *Bacillus cereus* BG1, *Enzyme and Microbial Technology* 2003; **32**(5): 513-518.
  15. Hideto Takami, Teruhiko Akiba, Koki Horikoshi Production of extremely thermostable alkaline protease from *Bacillus* sp. No. AH-101, *Applied Microbiology and Biotechnology*. 1989; **30**(2): 120-124.
  16. El Enshasy, H., Abuoul-Enien A., Helmy, S. and El Ezaly, Y. Optimization of the Industrial Production of Alkaline Protease by *Bacillus licheniformis* in Different Production Scales, *Australian Journal of Basic and Applied Sciences*. 2008; **2**(3): 583-593.
  17. Gopal K. Joshi, Sarvesh Kumar, Vinay Sharma Production of Moderately Halotolerant, SDS Stable Alkaline Protease from *Bacillus cereus* MTCC 6840 Isolated from Lake Nanital, Uttaranchal State, India, *Brazilian Journal of Microbiology*. 2007; **38**: 773-779.
  18. Ganesh Kumar C., Han-Seung Joo, Yoon-Mu Koo, Seung R., Paik and Chung-Soon Chang Thermostable alkaline protease from a novel marine haloalkaliphilic *Bacillus clausii* Isolate. *World Journal of Microbiology and Biotechnology* 2003; **20**(4): 351-357.
  19. Rao C.S., Sathish T., Ravichandra P. and Prakasham R.S., Characterization of thermo and detergent stable serine protease isolated from *Bacillus circulans* and evaluation of ecofriendly applications. *Process Biochemistry*. 2009; **44**(3): 262-268.
  20. Kumar, C.G. Purification and characterization of a thermostable alkaline protease from alkalophilic *Bacillus pumilus*. *Letters in Applied Microbiology* 2002; **34**, 13-17.
-