

## *International Journal of Scientific Research and Reviews*

### **Enumerations on Ethnobotanical, Phytochemical and Pharmacological Aspects of Guduchi (*Tinospora Cordifolia* (Willd.)Miers Ex Hook. F. And Thoms)**

**DK Sharma**

Department of Science and Technology, Vardhaman Mahaveer Open University, Kota, Rajasthan, India

Email: [drdilipsharma12@gmail.com](mailto:drdilipsharma12@gmail.com), [dksharma@vmou.ac.in](mailto:dksharma@vmou.ac.in)

#### **ABSTRACT**

Guduchi (*Tinospora cordifolia* (Willd.)Miers ex Hook. F. and Thoms) is a semi-perennial, glabrous, succulent climber grown universally as wild or cultivated in warmer areas. It contains commercially important grey-brown, rough, thin stem bark which used in various drugs. The plant climbs on other plants with fleshy thread like aerial roots. The dry stem is odourless but freshly cut stem has very bitter taste. It produced bioactive compound or secondary metabolites *viz.* columbin (tinospurin), chasmanthin, palmarin, cordioside, tinoside and cordifoliside-A. Stem contains several phenylpropanoids (syringin, cordifolioside-A, cordifolioside-B, cordiol and sinapic acid). Pharmacologically it has bioactive isoquinoline alkaloids (berberine, jatrorrhizine, magnoflorine, tembetarine, N-formylanonaine, N-formylnornuciferine), lignans (a phenolic lignan), carbohydrates (an arabinogalactan polysaccharide) and aliphatic compounds.

#### **\*Corresponding author**

**DK Sharma**

Department of Science and Technology,

Vardhaman Mahaveer Open University, Kota, Rajasthan, India (Corresponding Author)

Email: [drdilipsharma12@gmail.com](mailto:drdilipsharma12@gmail.com), [dksharma@vmou.ac.in](mailto:dksharma@vmou.ac.in)

## INTRODUCTION

In recent trends of research various parts of medicinal plants are used universally due to their natural origin and lesser side effect. Guduchi or *Tinospora* [*Tinospora cordifolia* (Willd.) Miers ex Hook. F. and Thoms] (Syn. *Menispermum crispum* L., *T. gibbericaulis* Handel-Mazzetti, *T. mastersii* Diels, *T. rumphii* Boerlage and *T. thorelii* Gagnepain) is a glabrous, deciduous climber of family Menispermaceae. This family have about 70 genera and 450 species found in tropical lowland regions and most of them are generally climber or twiner but rarely shrubs<sup>1-3</sup>. The family is medicinally important due to rich source of alkaloid and terpenes<sup>4</sup>. It is listed amongst 29 highly prioritized medicinal plants of agro climatic zone 8 (Rajasthan, U.P. and M.P.) by National Medicinal Plant Board, India due to the presence of immense medicinal properties<sup>5</sup>.

The genus *Tinospora* having 32 species distributed in tropical Africa, Madagascar, Asia, Australia and the Pacific Islands<sup>6-8</sup>. In India, the genus is represented mainly by 4 species. Out of 4 two species as *T. cordifolia* (Thunb.) Miers and *T. sinensis* (Lour.) Merr occurs in South India and other two namely *T. crispa* (L.) Hook.f. & Thomson and *T. glabra* (Burm. F.) Merr reported in Northeast India and the Andaman Islands<sup>9</sup>. *Tinospora cordifolia*, *T. uliginosa*, *T. malabarica*, *T. tomentosa*, *T. crispa* are medicinally important species of genus.

*T. cordifolia* is deciduous plant of dry forest distributed throughout tropical Indian subcontinent and China height up to 300 m. It contains alkaloids, diterpenoid lactones, glycosides, steroids, sesquiterpenoid, phenolics, aliphatic compounds and polysaccharides<sup>10</sup>. Protoberberine alkaloids, terpenoids and polysaccharides are 3 major groups of compounds considered as putative active constituents<sup>11-12</sup>.

## BOTANICAL DESCRIPTION

*T. cordifolia* is glabrous, deciduous, perennial climber, bearing distinct male and female flowers (flowering in month of March to June) grow as wide range of hedges and trees with coiling branches<sup>1, 13, 14</sup>. Stem is succulent, long, filiform and fleshy with gray or creamy white bark which have large rosette shaped lenticels on its surface. The wood is white, soft and porous; but when it exposed to air become yellow tint. The fresh stem is covered by green succulent thin brown bark which normally separate from the wood when the stem dry or shrink. Aerial roots arise from the branches and characterized by tetra to penta-arch as primary structure. The inner soft parenchymatous cortex is covered by outer thick walled<sup>15-16</sup>.

Branches are coiled and pendulous; fleshy roots covered by striate, tubercled, pale-shining or glabrous bark. Leaves are simple, membranous, alternate, exstipulate, long petiolated (2.5-7.0 cm),

chordate in shape with multicoated reticulate venation (7-9 nerved). Small yellowish to greenish unisexual flowers emerge from leafless plant in summer.

Female flowers are solitary while males are found in cluster (terminal racemes). Six petals arranged equally in two whorls, broadly spatulate, claw cuneate, reflexed to apex, pistillode. Six sepals arranged in two whorls of 3 each, outer is very small, ovate-oblong, acute but the inner whorl has 3 sepals larger, membranous, broadly elliptical, concave and yellow. Solitary female flower has green sepals without reflex margins, staminode short, linear. Carpels 1-3, widely separated on the short fleshy gynophores, dorsally convexed and scarlet in colour<sup>14</sup>.

Red fruits are found in cluster with fleshy thick stalk and sub terminal style scars<sup>17</sup>. The fruits (drupes) are ovoid, pea-sized; glossy became red and mucilaginous when ripe in month November-January<sup>18-19</sup>. The embryo is curve automatically due to the curvature in seed (moonseed family). The endocarp is ornamented (taxonomic characters).

**Common Name(s):** Guduchi, Amrita (Sanskrit), Gadancha, Giloe, Gulancha (Bengali), Giloya, Giloe, Gulbel, Gurcha (Hindi), Gulo, Gado, Galo (Gujarati), Heartleaf Moonseed, Tinospora (English), Gilo (Arabic); Amarlata (Assamese); Amrytu, Sittamrytu (Malayalam); Ambarvel, Giroli, Gulvel (Marathi), Garjo (Nepali); Gulancha (Oriya); Gulbel (Persian); Gilo (Punjabi, Kashmiri), Gurjo (Sikkim); Shindila Kodi, Amridavalli, Niraidarudian (Tamil); Duyutige, Teppatige, Guduchi, Iruluchi (Telugu), Guruch (Urdu), Amrutha Balli (Kannada), Brotowali, Andawali, Putrawali (Indonesia), Makabuhay (Tagalog), Paliaban, Pañgiauban (Bisaya), Taganagtagua & Boraphet (Thailand), K'uan Chu Hsing (Chinese) and Culancha (French)<sup>1,13,20</sup>.

**DISTRIBUTION:** The plant is indigenous to India, Myanmar, Sri Lanka, China, Thailand, Philippines, Indonesia, Malaysia, Borneo, Vietnam, Bangladesh, North Africa, West Africa and South Africa<sup>21-23</sup>. In Indian subcontinent, it is found as climber on *Jatropha curcas*, *Azadirachta indica*, *Moringa oleifera* and other plants which provide support to it<sup>24,25</sup>. When it climbs on Neem (*Azadirachta indica*) as a climber it is called Neem giloy having combination of chemical constituents of both neem and giloy (tinospora)<sup>26</sup>.

**Growth Requirement-** The plant prefers wide range of soil; acid to alkaline with moderate soil moisture. It can grow in almost all climates but it prefers warm climate. Plantation in the months of July-August in well drained medium black or red soil with rich organic matter is good for plant growth. It is propagated by seeds, vegetative cuttings or clonal propagation<sup>27</sup>.

**Nutritive composition-** *T. cordifolia* contains high fibre (15.9%), protein (4.5%-11.2%), carbohydrate (61.66%) and low fat (3.1%) with 292.54 calories as nutritive value (per 100 g). It contains high amount of potassium (0.845%) and chromium (0.006%); sufficient amount of iron (0.28%) and calcium (0.131%)<sup>28</sup>.

## ETHNOBOTANICAL AND TRADITIONAL USES

Different tribal or ethnic groups have their own tradition, folk language, beliefs and knowledge to use natural resources as medicines<sup>29</sup>. All the plant parts of *T. cordifolia* are documented to be useful in ethno botanical surveys<sup>30-31</sup>. The plant is used in folkloric veterinary medicine and traditional ayurvedic medicine or reported to widely used for anti-inflammatory and anti-arthritic problems<sup>24, 21</sup>.

The leaves used in the treatment of gout and ulcer. Stem stimulates bile secretion, diuretic, enriches the blood, cures jaundice and useful in skin diseases. Extract of stem used to treat chronic diarrhea, chronic dysentery, and intestinal problems and improve digestion<sup>32</sup>. The juice of stem used to cure diabetes, vaginal and urethral discharges, low fevers and enlarged spleen<sup>13</sup>. Stem decoction used to washing sore eyes, syphilitic sores, antipyretic and anti-malarial.

In folk medicine the whole plant or powdered root and stem bark, root and stem decoction, juice of the root, paste of the leaves and stem used to treat various disease disorders viz. fever, jaundice, diarrhea, dysentery, general debility, cough, asthma, leucorhea, skin diseases, fractures, eye disorders and bites of poisonous insects. The combination of stem root with other drugs is used as antidote to snake bite and scorpion sting<sup>18, 33, 34</sup>. Dried fruits in combination of ghee or honey are used as tonic to cure jaundice and rheumatism.

## PHYTOCHEMISTRY

Stem contains several bioactive compounds like alkaloids viz. berberine, palmatine-D, choline-D, tinosporine, magnoflorine, tetrahydropalmatine, isocolumbin<sup>35-40</sup>, 18-norclerodane glycoside<sup>41</sup>. Furanoidditerpene glycoside<sup>42,43</sup> as tinocordiside<sup>44,45</sup>, syringin<sup>46,47</sup> syringin- apiosylglycoside<sup>41,42</sup> tinocordifolioside, cordioside, cordifolioside A, cordifolioside B, palmatoside-C31, palmatoside-F31, cordiofolioside-B2, cordifolioside-D2, cordifolioside<sup>43,44-48,49</sup> and sesquiterpenoid tinocordifolin<sup>50</sup>. The root contains palmatine as an important alkaloid<sup>38,51</sup>. The whole plant contains diterpenoid lactones, furanolactone, tinosporon, columbin<sup>52-53</sup>; aliphatic compound as octacosanol, heptacosanol<sup>54</sup> and other miscellaneous compound are nonacosan-15-one 3, (a-4-dihydroxy-3-methoxy-benzyl)-4-(4-hydroxy-3-methoxy-enzyl)-tetrahydrofuran, tinosponidine, 6-cordifol, 6-

cordifellone, 6-jatrorrhizine<sup>55,56</sup>. The aerial plant parts have steroids b-sitosterol, d-sitosterol, g-sitosterol b-hydroxyecdysone, ecdysterone, makisterone, giloinsterol jateorine and columbin<sup>51, 57</sup>.

## **USES IN INDIAN SYSTEM**

The stems, leaves and roots of guduchi are used in various medicines<sup>58, 59</sup>. In Hindu mythology giloy means heavenly elixir which saved celestial beings from old age and kept them eternally young<sup>10</sup>. In Ayurveda, various dosages in form of Swaras, Kalka, Churna, Kwatha, Fant, Arishta, Satwa, Ghana, Guduchi Ghrita and Guduchi Taila used for treatment of diseases<sup>60</sup>. It is used as Chhinnaruha/Chhinnodbhava (plant's capacity to grow from the cut side), Vatsadini (eaten by grazing animals), Pittaghi (bile destroying), Amruta (imparts immortality), Rasayana (capacity to improve quality of rasa, the primordial tissue which strengthens all other tissues), Chakrangi/Chakra-akshana (wheel-like appearance), Jwaranashi/ Jwarari (potent antipyretic), Vayastha (prevents ageing), Amrutsambhava (ambrosia), Bhishakpriya (favourite of physicians), Saumya (not harmful) and Tikta (bitter taste).

Guduchi used as several formulations to treat general weakness, fever, dyspepsia, skin diseases, anaemia, dysentery, asthma, anorexia, jaundice, gout, diabetes, gonorrhoea, secondary syphilis, urinary diseases, impotency, viral hepatitis, leprosy, chronic diarrhoea and dysentery<sup>61-70</sup>.

Charaka described guduchi rasayana as antiaging, immunostimulant, life span booster, promotes intelligence, memory enhancer, resistance, anti-spasmodic and anti-diarrhoeal<sup>71, 72</sup>, anti-stress; promote longevity and adaptogen<sup>17, 24, 73-75</sup>. In Sushruta Samhitaa guduchi described it as antiseptic, febrifuge, appetizer, abdominal dropsy, haemoptysis, urinary, vaginal disorders, skin diseases and oedema. It is component of various ointments used for healing of wounds and burns. It is used to treat hyperacidity, colic pain, sprue, dysentery, rheumatism and cellulitis. A crude aqueous dry stem starch preparation is used to cure gastrointestinal disorders or as a tonic in debilitating diseases<sup>76</sup>. The commercially available products to cure various diseases or as supplement to cure various diseases are Tinospora Pellets, Cirrholiv-DS syrup, Mussaffen, Madhu Mehari, Guduchi, Abhaibhubejhr, Safe herb, Brave Heart capsule, Cirrholiv capsules, Tonoplex, giloya vati and Rebuild<sup>27</sup>.

## **PHARMACOLOGICAL ACTIVITIES**

Recently guduchi extracts/fractions used in various forms (aqueous, alcoholic or methanolic, chloroform, ether, petroleum ether, acetone) or chemical constituents used in numerous pharmacological, pre-clinical and clinical investigations. The active adaptogenic constituents in plant

are diterpene compounds alkaloid, Giloinin(glucosidal bitter principle), polysaccharides, Picrotene and bergenin<sup>17, 74</sup>. It has anti-inflammatory, anti-oxidant, anti-pyretic and immunomodulatory activities<sup>17, 73-75</sup>. Bark of plant is anti-allergic, anti-spasmodic, anti-leprotic or used to treat urinary diseases, syphilis, skin diseases, bronchitis<sup>77</sup>, promote longevity and increase body's resistance<sup>77-81</sup> or induce the immunity<sup>82-84</sup>.

Guduchi has immunomodulatory and cytotoxic affects due to various compounds<sup>82, 85-97</sup>. Both humoral and cell-mediated immunity enhanced in a dose-dependent manner<sup>47</sup>. An immunologically active arabinogalactan identified from dried stem<sup>11</sup>. These natural compounds improve the phagocytic activity of macrophages, enhancement in nitric acid production by stimulation of splenocyte and production of reactive oxygen species (ROS) in human neutrophil cells<sup>27, 98-99</sup>. The aqueous extract improve phagocytic and intracellular bacterial capacities of neutrophils in mice reported against bacterial growth<sup>100</sup>.

The plant has various properties viz. antioxidant<sup>23, 101-107</sup>, anti-hyperglycaemic<sup>108-111</sup>, anti-cancerous or antitumor<sup>80, 89, 104, 112-115</sup>.

1. The stem used to cure diabetes due to the presence of alkaloids (Magnoflorine, Palmetine and Jatrorrhizine) tannins, glycosides, flavonoids, saponins, steroids by promoting insulin secretion or by inhibiting gluconeogenesis and glycogenolysis<sup>116-118</sup>. The stem extract in various organic solvents like ethylacetate, dichloromethane, chloroform and hexane inhibits the enzymes like salivary amylase and glucosidase resulting increase in post-prandial glucose level and shows potential activities to cure diabetes mellitus disease<sup>119</sup>. The root extract also has anti-diabetic properties, reduce the level of glycosylated haemoglobin, hydroperoxidase and vitamin E<sup>111, 120-122</sup>. Oral dose (aq.) of roots significantly reduce in thiobarbituric acid and increase in reduced glutathione, catalase and superoxide dismutase in alloxan diabetic rats<sup>89</sup>. Arabinogalactan polysaccharide protect against iron-mediated lipid peroxidation of rat brain homogenate (also vide infra)<sup>96</sup>.

The plant has antipyretic, analgesic, antiicteric<sup>80, 104</sup>, antigonorrhoeal<sup>120-122</sup>, antifertility<sup>123</sup>, anticancer<sup>112, 114, 124, 125</sup>, anti-stress<sup>127</sup>, hypolipidaemic<sup>104, 128-129</sup>, antimalarial<sup>120-122, 130</sup>, anti-allergic<sup>105, 131</sup>, antiamebic, antiatherogenic, antihelmentic<sup>80, 120-122</sup>, antigout, antiasthmatic, antiperiodic, androgenic<sup>78</sup>, anticoagulant, antimicrobial properties<sup>104</sup>. It help to dissolve urinary calculi<sup>132</sup>, infections<sup>133</sup>, phagocytes<sup>134</sup>, memory booster<sup>135</sup>, anti-ischemic<sup>136</sup>, anti-leprotic<sup>29</sup>, antiulcer<sup>137, 138</sup>, hepatoprotective<sup>89, 139</sup>, cure splenic injury<sup>95, 130</sup> and anti-inflammatory<sup>29, 80, 105, 140-145</sup>.

Stem extract (aqueous) exhibit adaptogenic properties<sup>146</sup>. In rat, it evidenced that liver damage by carbon tetrachloride (0.7 ml/kg/7 days) as evidenced by higher levels of enzymes (SGOT, SGPT,

AP) and serum bilirubin. When these rats treated with stem extract (100 mg/kg) for 15 days found significant reduction in the serum levels of these enzymes and bilirubin was observed<sup>95</sup>. In a clinical study of 30 patients (who received surgical treatment for extra hepatic obstructive jaundice) given aqueous extract (16 mg/kg/day/oral) of guduchi stem with usual medication showed a far superior postoperative survival rate<sup>83</sup>.

Aqueous extract of stem showed antiinflammatory, analgesic and antipyretic actions in rats. The extract @ 500 mg/kg (oral) significantly inhibited acute inflammatory response evoked by carrageenan. It is reported that syringin and cordiol found in stem extract have potent anticomplement properties<sup>147</sup>. In asthma, stem extract significantly reduced bronchospasms (allergic) induced by 5.0% histamine aerosol in guinea pigs and capillary permeability in mice<sup>79</sup>.

Plant has anti-neoplastic<sup>112, 148</sup>, antiasthmatic, antiperiodic, antiemetic, radioprotective<sup>149</sup>, anti-depression<sup>150</sup> and obstructive jaundice properties<sup>83, 151</sup>.

It has protected effect by lowering the concentration of thiobarbituric acid and enhancing the glutathione, vitamin-C and protein, antioxidant enzymes (superoxide dismutase, catalase, glutathione peroxidase, glutathione S-transferase), glutathione reductase in kidney<sup>152</sup>. In male albino, leaf and stem extract hepatoprotective effect against lead nitrate induced toxicity. Oral dose of plant extract showed prohibited the lead nitrate induced liver damage<sup>53</sup>.

Root extract shown radio protective activity due to extensively increase in body weight, tissue weight and tubular diameter. Dichloromethane extracts of guduchi shows cytotoxic effects owing to lipid peroxidation and release of LDH and decline in GSH<sup>125</sup>. In pre-irradiating mice, root extract has widely affected radiation, induced rise in lipid peroxidation and resulted in the decline of GSH in testes<sup>154</sup>. Most of the synthetic chemotherapeutic agents laid toxic side effects on the living organisms<sup>155</sup>.

In a study methylene dichloride extract of stem showed significant anticancer activity against cultured HeLa cells. Antineoplastic activity of extract was comparable to that of doxorubicin. Aqueous and alcoholic extracts of plant was also active, but less potent than the methylene dichloride extract<sup>112</sup>. A polysaccharide fraction from the plant was demonstrated inhibitory in experimental metastasis in mice<sup>105</sup>.

Methanolic extract of guduchi has antimicrobial activities and reported against *Staphylococcus aureus*, *Salmonella paratyphi*, *S. typhimurium*, *Klebsiella pneumonia*, *Escherichia coli*, *Enterobacter aerogene*, *Proteus vulgaris*, *S. typhi*, *Shigella flexneri*, *Pseudomonas aeruginosa* and *Enterobacter aeruginosa*<sup>156-158</sup>.

Methanolic extract of stem increase the erythrocytes, membrane lipid peroxide and catalase activity and decreases the activity of SOD, GPx in alloxan induced diabetic rats<sup>159-160</sup>. Leaf extract



have an alpha-glucosidase inhibitor, characterized as saponarin was found to be also significant antioxidant and hydroxyl radical scavenging activity<sup>155</sup>. Aqueous extract has a radio protective activity, enhancing the survival of mice against a sub-lethal dose of gamma radiation<sup>153, 161</sup>.

Purified polysaccharide fraction of stem showed photo-protective and effectively quenched free radicals generated by photons which protect the tissues against oxidative damage<sup>162</sup>. Stems extract (aq.) has radioprotective activity and inhibited radiation mediated 2-deoxyribose degradation in a dose-dependent manner<sup>161</sup>.

## ACKNOWLEDGEMENT

An author is grateful to all the scientists whom work is cited and could not acknowledge unknowingly and persons that directly or indirectly engaged in writing in this review.

## Conflict of interest

There is no conflict of interest in writing this review.

## REFERENCES

1. Anonymous. The Wealth of India: Raw Materials, Council of Scientific and Industrial Research, New Delhi. 1956; 10: 251–252.
2. Aima RK. Pictorial Guide to Plants. Natraj Publishers First Edition Dehradun 2003.p. 454-55.
3. Vaidya DB. Materia Medica of Tibetan Medicine. Delhi: Sri Satguru Publications 1994. p.163.
4. Sharma A, Gupta A, Singh S and Batra A. *Tinospora cordifolia* (Willd.) Hook. F. & Thomson- A plant with immense economic potential. J. Chem. Pharm. Research 2010; 2(5):327-333.
5. Anonymous. National Medicine Plant Board. (Last seen 21-1-2019 www.nmpb.nic.in).
6. Forman LL. A revision of *Tinospora* (Menispermaceae) in Asia to Australia and the Pacific. Kew Bull. 1981; 36(2): 375-421.
7. Kubitzki K, Rohwer JG and Bittrich V. The Families and Genera of Vascular Plants, vol. II. Berlin/Heidelberg, Germany: Springer-Verlag.1993
8. Mabberley DJ. The Plant-Book. A Portable Dictionary of the Vascular Plants Cambridge: Cambridge University Press. 2005.
9. Pramanik A and Gangopadhyay M. Menispermaceae. In: Sharma BD, Balakrishnan NP, Rao RR. And Hajra PK. (eds) Flora of India 1. Calcutta: Botanical Survey of India. 1993.



10. Bhandari C and Vanaushadhi Chandrodaya. Chaukhamba Sanskrit Sansthan. Varanasi, First Edition 2006; 3: 86.
11. Chintalwar G, Jain A, Sipahimalani A, Banerji A, Sumariwalla P, Ramakrishnan R, Sainis K. An immunologically active arabinogalactan from *Tinospora cordifolia*. *Phytochemistry* 1999; 52(6):1089-1093.
12. Bisset NG, Nwaiwu J. *Planta Medica* .1983; 48: 275.
13. Kirtikar KR and Basu BD. *Indian Medicinal Plants*, Lalit Mohan Basu, Allahabad, 1918; 1: 75-80.
14. Kirtikar KR, Basu BD. *Indian Medicinal Plants*. Edn 2, Vol. 1, M/S Bishen Singh, Mahendra Pal Singh; 1975. *Journal of Medicinal Plants Studies* 2014 p- 43.
15. Aiyer KN, Kolammal M. *Pharmacognosy of Ayurvedic Drugs*. Edn 1, The Central Research Institute 1963, series 1.
16. Khosa RL, Prasad S. Pharmacognostical studies on Guduchi (*Tinospora cordifolia* Miers). *J Res Ind Med*. 2010;6:261-269.
17. Anonymous. *British Pharmaceutical Codex (BPC)*. Published by Pharmaceutical Society of Great Britain (PSGB) 1907; current Twelve Edition, published in 1994.
18. Nadkarni KM and Nadkarni AK. *Indian Materia Medica*, Mumbai: Popular Prakashan Private Limited 3<sup>rd</sup> edtn Vol. 1. 1976. p. 356-365, p. 1220-1221.
19. Kirtikar KR and Basu BD. *Indian Medicinal Plants*. Second Edition, Edited and enlarged by Blatter E, Caius JF and Mhaskar KS. International Book Distributors, Dehra Dun, 2005; 1: 77-81.
20. Anonymous. 2001. *The ayurvedic pharmacopoeia of India*. Part I. 1<sup>st</sup> Edition Vol. 1. New Delhi: Department Of AYUSH, Ministry of Health and FW. 2001. p. 53-55.
21. Mia MMK, Kadir MF, Hossan MS and Rahmatullah M. Medicinal plants of the Garo tribe inhabiting the Madhupur forest region of Bangladesh. 2009.
22. Singh J, Sinha K, Sharma A, Mishra NP and Khanuja SP. Traditional uses of *Tinospora cordifolia* (Guduchi) *J Med Aromat plant Sci.*, 2003; 25: 748-51.
23. Jain S, Sherlekar B and Barik R. Evaluation of antioxidant potential of *Tinospora cordifolia* and *Tinospora sinensis*. *Int J Pharm Sci Res*, 2010; 1(11): 122-8.
24. Anonymous. *Ayurvedic Pharamacopoeia of India*. Guduchi, published by dept of AYUSH, ministry of health & family welfare, govt. of India. 1999; 1(1): 53-55.
25. Premila MS. *Ayurvedic Herbs: A clinical Guide to the Healing Plants of Traditional Indian Medicine*. New York: Haworth Press, 2006. p. 69-76, 175-176.

26. Chaudhari S, Shaikh N. Gaduchi-the best ayurvedic herb. The Pharma Innovation Journal 2013; 2(4):97-102.
27. Mittal J, Sharma MM and Batra A. *Tinospora cordifolia*: a multipurpose medicinal plant- A review. Journal of Medicinal Plants Studies, 2014; 2 (2): 32-47.
28. Nile SH and Khobragade CNN. Determination of nutritive value and mineral elements of some important medicinal plants from western part of India. J Med Plants, 2009; 8(5): 79-88.
29. Singh SS, Pandey SC, Srivastava S, Gupta VS and Patro B. Chemistry and medicinal properties of *Tinospora cordifolia* (Guduchi). Indian Journal of Pharmacology, 2003; 35:83-91.
30. Lalramnghinglova H. Ethno-medicinal plants of Mizoram Dehradun; Bishen Singh Mahendra Pal Singh.2003. p-293.
31. Sood SK, Parmar S and Lakhanpal TN. Ethnic plants of India used in cancer cure. Dehradun: Bishen Singh Mahendra Pal Singh 2005.p. 235.
32. Khory RN and Katrak NN: Materia, Medica of India and their therapeutics: Komal Prakash, Pg 31.
33. Kiritikar KR and Basu BD. 1933. Indian medicinal plants. Volume I. Lalit Mohan Publishers, India. 1933. p- 77.
34. Zhaotf, Wang X and Rimando AM. Chec. Folk tonic medicinal plants: *T. sgittata* var. *Cravaniana* and *Mahonia beabi*. Planta med 1991;57: 505
35. Qudrat -I- Khuda M, Khaleque M and Ray N. Sci Res (Dacca) 1964; 1, 177.
36. Pachaly P and Schneider C. Arch Pharma (Weinheim ger), 1981; 314, 251.
37. Bisset NG, Nwaiwu J Planta Medica .1983; 48, 275.
38. Sarma DNK, Padma P and Khosa RL. 1998. Fitoterapia, 69: 541.
39. Kumar S, Verma NS, Pande D and Srivastava PS. *In vitro* regeneration and screening of berberine in *T. cordifolia*. Journal of Medicinal and Aromatic Plant Sciences, 2000; 22: 61.
40. Padhya MA. Biosynthesis of Isoquinoline alkaloid berberine in tissue cultures of *T. cordifolia*. Indian drugs, 1986; 24:47-8.
41. Khan MA, Gray AL and Waterman PG. Tinosporaside, an 18- norclerodane glucoside from *T.cordifolia*. Phytochemistry 1986; 28: 273- 275.
42. Bhatt RK and Sabata BK. Furanoid diterpene glucoside from *T. cordifolia*. Phytochemistry, 1989; 28: 2419-2422.
43. Swaminathan K, Sinha UC, Bhatt RK, Sabuta RBK and Tavale SS. Structure of Tinosporide, a diterpenoid furanolactone from *T.cordifolia* miers. Acta crystallogm, 1989; 45: 134-6.

44. Maurya R, Wazir V, Tyagi A and Kapil RS. Clerodane diterpenoids from *T.cordifolia*." *Phytochemistry*1995; 38: 559-61.
45. Ghosal S and Vishwakarma RA. Tinocordiside, a new rearranged cadinane sesquiterpene glycoside from *T. cordifolia*. *Journal of Natural Product*, 1997; 60: 839-841.
46. Sipahimalani AT, Noerr H and Wagnor H. Phenyl propenoid glycosides and tetrahydro furanlignan glycosides from the adaptogenic plant drugs *T.cordifolia* and *Drypetes roxburghii*. *Planta Medica*, 1994; 60: 596-597.
47. Kapil A and Sharma S. Immuno potentiating compounds from *T.cordifolia*. *J. Ethnopharmacol*, 1997; 58: 89-95.
48. Wazir V, Maurya R and Kapil RS. *Phytochemistry* 1995; 38, 447.
49. Maurya R, Dhar KL and Handa SS. *Phytochemistry* 1997; 44, 749.
50. Maurya R and Hardass. Tinocordifolin, a sesquiterpene from *T.cordifolia*." *Phytochemistry*, 1998; 49: 1343-6.
51. Pathak AK, Agarwal PK, Jain DC, Sharma RP and Howarth OW. NMR studies of 20b-hydroxy ecdysone, a steroid, isolated from *T. cordifolia*." *Indian J. Chem Sec B*, 1995; 34: 674-6.
52. Ahmad M, Kazi AB, Karim R, Khaleque A and Miah MAW. Structure of tinosporide, a furanoid diterpene from *T. cordifolia*. *Journal of Bangladesh Academy of sciences*, 1978; 2: 25- 30.
53. Hanuman JB, Bhatt RK and Sabata BK. A diterpenoid furano lactone from *T. cordifolia*. *Phytochemistry*, 1986; 25: 1677-1680.
54. Qudrat-I-Khuda M, Khaleque A, Bashir A, Roufk MDA and Ray N. Studies on *T. cordifolia* Isolation of tinosporon, tinosporic acid and tinosporol from fresh creeper. *Scientific Research* 1966; 3: 9-12.
55. Dixit SN and Khosa RL. Chemical investigation of *T. cordifolia*. *Indian J. Appl Chem.*, 1971; 34: 46-7.
56. Khaleque A, Miah MAW, Huq MS and Abdul BK. *Sci Res (Dacca)* 1970; 7, 61.
57. Khaleque A, Miah MAW, Huq MS and Abdul BK. *Pak J Sci Ind Res*. 1971; 14, 481.
58. Gangan VD, Pradhan P and Sipahimalani AT. 1997. *Indian J Chem Sec B* 36: 787.
59. Pandey G. *Dravyaguna Vijnana* .Varuanasi: Krishnadas Academy 2002; Second Edition pp. 1: 697-710.
60. Krishna K, Jigar B and Jagruti P. Guduchi *Tinospora cordifolia*: Biological and Medicinal properties, a review. *The Internet Journal of Alternative Medicine*, 2009; 6 (2): 1-10.

61. Panchabhai TS, Kulkarni UP and Rege NN. Validation of Therapeutic Claims of *Tinospora cordifolia*: A Review Phytother. Res. 2007; 22, 425-441 (www.interscience.wiley.com).
62. Gupta SS, Verma SC, Garg VP and Rai M. Anti-diabetic effects of *Tinospora cordifolia*. Effect on fasting blood sugar level, glucose tolerance and adrenaline induced hyperglycaemia. Indian J Med Res, 1967; (55).p. 733-745.
63. Chopra RN, Chopra IC, Handa KL and Kapur LD. Chopra's Indigenous Drugs of India Second Edition Calcutta, India: B.K. Dhur of Academic Publishers, 1982; 426-428.
64. Chintalwar G, Jain A and Sipahimalani A. An immunologically active arabinogalactan from *Tinospora cordifolia*. Phytochemistry, 1999; 52: 1089-1093.
65. Tripathi B. Charak Samhita part I Varanasi: Chaukhambha Krishnadas Academy, 2003. P. 454.
66. Bhatta KR, Bhatta RK and Swami LR. Siddha Bhaisaiya Mani Mala Vaishwanara Hindi commentary. Third Edition Varanasi: Chaukhambha Krishnadas Academy 2003.p-31.
67. Sharma AR. Sushrut Samhita: Sushrutvimarshini Hindi Commentary along with special deliberation etc. Part I. Varanasi: Chaukhambha Surbharati Prakashan, 2000. p- 419.
68. Sharma AR. Sushrut Samhita: Sushrutvimarshini Hindi Commentary along with special Deliberation etc. Part II. Varanasi: Chaukhambha Surbharati Prakashan, 2001; 311- 317.
69. Chunekar KC and Pandey GS. Guduchyadi Varga Bhavprakash Nidhantu Varanasi: chaukhambha Bharati Academy, 2006. p- 269.
70. Tripathi I. 2006. Arkaprakash Tritiya Shatak Varanasi: Chaukhambha Krishnadas Academy 2006. p-45.
71. Sharma PV and Sharma GP. Kaiyadeva Nighantu” Varanasi: Chaukhambha Orientalia 2006; p- 5.
72. Warriar PK, Nambiar VPK, Ramankutty C and Nair RV. Indian medicinal plants: a compendium of 500 species 1996; 5: 283.
73. Badara VA, Thawani VR, Wakode PT, Shrivastava MP, Gharpure KJ and Hingorani LL. Efficacy of *Tinospora cordifolia* in allergic rhinitis. J Ethnopharmacol, 2005; 96: 445-9.
74. Patwardhan B and Gautam M. Drug Discov Today, 2005; 10:495–502
75. Patil M, Patki P, Kamath HV and Patwardhan B. Ind Drugs, 1997; 34:211-215.
76. Anonymous. Indian Pharmacopoeia. Guduchi, Published by Indian Pharmacopoeia Commission, Govt. Of India. 2007; (3): 2037-2034.
77. Bishayi B, Roychowdhury S, Ghosh S and Sengupta M. Hepatoprotective and immunomodulatory properties of *Tinospora cordifolia* in CCl<sub>4</sub> intoxicated mature albino rats. J Toxicol Sci, 2002; 27(3): 139-46.

78. Treadway S. Exploring the universe of ayurvedic botanicals to manage bacterial infections. *Clinical Nutrition Insights*, 1998; 6(17): 1-3.
79. Kapur P, Pereira BM, Wuttke W and Jarry H. 2008. Androgenic action of *Tinospora cordifolia* ethanolic extract in prostate cancer cell line LNCap, *Phytomedicine* 2009; 16(6-7):679-82.
80. Nayampalli, SS, Desai, NK, Ainapure, SS. Antiallergic properties of *Tinospora cordifolia* in animal models. *Indian J. Pharmacol*, 1986; 18 (4): 250-252.
81. Ikram M *et al.* *Journal of Ethnopharmacol.* 1987; 19(2): 185-192.
82. Asthana JG, Jain S, Mishra A and Vijaykanth MS. *Indian Drugs* 2001; 38, 82-6.
83. Rege N, Bapat RD, Koli R, Desai NK, Dahakar S. Immunotherapy with *Tinospora cordifolia*: a new lead in the management of obstructive jaundice. *Indian Journal of Gastroenterol*, 1993; 12 (1): 5-8.
84. Vasudevan DM and Sreekumari S. Text book of biochemistry for medical students. New Delhi: Jaypee Brothers Medical, 1995; 194-195.
85. Dahanukar SA, Thatte UM, Pai N, More PB and Karandikar SM. Immunotherapeutic modification by *Tinospora cordifolia* of abdominal sepsis induced by caecal ligation in rats. *Indian J Gastroenterol*, 1988; 7: 21-3.
86. Thatte UM, Kulkarni MR and Dahanukar SA. Immunotherapeutic modification of *Escherichia coli* peritonitis and bacteremia by *Tinospora cordifolia* *J Postgrad Med* 1992; 38:1; 13-5.
87. Thatte UM, Rao SG and Dahanukar SA. *Tinospora cordifolia* induces colony stimulating activity in serum. *J Postgrad Med* 1994; 40; 202-24.
88. Bapat RD, Rege NN, Koti RS, Desai NK and Dahanukar SA. Can we do way with PTBD? *HPB Surg* 1995; 9: (1); 5-11.
89. Prince PSM and Menon VP. Antioxidant activity of *Tinospora cordifolia* roots in experimental diabetes. *Journal of Ethnopharmacology* 1999; 65: 277-281.
90. Manjrekar PN, Jolly CI and Narayanan S. Comparative studies of the immunomodulatory activity of *Tinospora cordifolia* and *Tinospora sinensis*. *Fitoterapia* 2000; 71 (3): 254-257.
91. Dikshit V, Damre AS, Kulkarni KR, Gokhale A and Saraf MN. Preliminary screening of immunocin for immunomodulatory activity. *Indian J Phar Sci*, 2000; 62: 257.
92. Bishayi B, Roychowdhury S, Ghosh S, Sengupta M. Hepatoprotective properties of a *Tinospora cordifolia* in CCl<sub>4</sub> intoxicated mature albino rats. *J Toxic Sci*, 2002; 27:139-146.

93. Nair PK, Melnick SJ, Ramachandran R, Escalon E and Ramachandran C. Mechanism of macrophage activation by (1,4)-alpha-D-glucan isolated from *Tinospora cordifolia*. Int Immunopharmacol, 2006; 6:1815-24.
94. Tripathi YB, Sharma M, Manickam. Rubia 5 din, a new antioxidant from rubiacordifolia. Ind J Biochem Biophy, 1997; 34:302-306.
95. Bishayi B, Roychowdhury S, Ghosh S and Sengupta M. Hepatoprotective and immunomodulatory properties of *Tinospora cordifolia* in CCl<sub>4</sub> intoxicated mature albino rats. J Toxicol Sci 2002; 27:3; 139-46.
96. Subramanian M, Chintawar GJ, Chattopadhyay S. Antioxidant properties of *Tinospora cordifolia* polysaccharide against iron-mediated lipid damage and gamma-ray induced protein damage. Redox Rep 2002; 7:137-143.
97. Sharma P, Parmar J, Sharma P, Verma P, Goyal PK. Radiation- induced testicular injury and its amelioration by *T. cordifolia* (An Indian Medicinal plant) extract. Evid based comp altern Med. 2012; 643-647.
98. Handa and Kaul, Somatotype and Disease- A Review 2007; 74: 413, [Reviews.]
99. More P and Pai K. *In vitro* NADH-oxidase and myeloperoxidase activity of macrophages after *Tinospora cordifolia* (guduchi) treatment. Immuno Pharma Immuno toxic 2010; 34:368-372.
100. Sengupta S, Mukherjee A, Goswami R, Basu S. Hypoglycemic activity of the antioxidant saponarin, characterized as alpha-glucosidase inhibitor present in *Tinospora cordifolia*. J Enzyme Inhib Med Chem, 2009; 24:684-690.
101. Robak J and Gryglewski RJ. Flavonoids are scavengers of superoxide anions. Biochem Pharmacol, 1998; 37; 837-45.
102. Kumar PV, Shashidhara S, Kumar MM and Sridhara BY. Effect of *Luffa echinata* on lipid peroxidation and free radical scavenging activity. J Pharm Pharmacol., 2000; 52; 891-8.
103. Stanely P *et al.* Journal of Ethnopharmacol. 2000; 70: 9-15.
104. Mary N *et al.* Phytomedicine, 2003; 10: 474-482.
105. Leyon PV and Kuttan G. Effect of *Tinospora cordifolia* on the cytokine profile of angiogenesis-induced animals. International Immunopharmacology, 2004; 1569-1575.
106. Singh RP, Banerjee S, Kumar PV, Raveesha KA and Rao AR. *Tinospora cordifolia* induces enzymes of carcinogen/drug metabolism and antioxidant system, and inhibits lipid peroxidation in mice. Phytomedicine, 2006; 13 (1-2): 74-84.

107. Manjusha GV, Rajathi K, Alphonse JKM and Meera KS. Antioxidant potential and antimicrobial activity of *Andrographis paniculata* and *Tinospora cordifolia* against pathogenic organisms. J Pharm Res. 2011; 4-2: 452-5.
108. Raghunathan K and Sharma PV. The aqueous extract of Gulvel caused reduction of blood sugar in alloxan-induced hyperglycaemic rats and rabbits. J Res Indian Med, 1969; 3: 203-9.
109. Grover JK, Vats V and Rathi SS. Anti-hyperglycaemic effect of *Eugenia jambolana* and *Tinospora cordifolia* in experimental diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. J Ethnopharmacol. 2000; 73(3):461-70.
110. Nagaraja PK, Kammar KF and Devi S. Modulation of morphology and somegluconeogenic enzyme activity by *Tinospora cordifolia* (Willd.) in diabetic rat kidney. Biomed Res., 2007; 18; 179-83.
111. Umamaheshwari S and Mainzen Prince PS. Antihyperglycaemic effect of logen-Excel, an ayurvedic herbal formulation in streptozotocin - induced diabetes mellitus. Acta Pol Pharm, 2007; 10:1375-1386.
112. GC Jagetia, V. Nayak, MS. Vidyasagar, Cancer Lett., 1998; 127: 71.
113. Vedavathy S and Rao KN. Antipyretic activity of six indigenous medicinal plants of Tirumala hills, Andhra Pradesh, India. J Ethnopharmacol. 1991; 33: 193-6.
114. Mathew, S. Kuttan, G. Antioxidant activity of *Tinospora cordifolia* and its usefulness in the amelioration of cyclophosphamide induced toxicity J. Exp. Clin, Cancer Res., 1997; 16(4):407-11.
115. Perumalsamy R *et al.* Fitoterapia, 2005; 76: 697-699.
116. Patel MB, Mishra S. Hypoglycemic activity of alkaloidal fraction of *Tinospora cordifolia*. Phytomedicine, 2011; 18:1045- 1052.
117. Zinjarde SS, Bhargava SY, Kumar AR. Potent  $\alpha$ -amylase inhibitory activity of Indian Ayurvedic medicinal plants. BMC Complement Altern Med, 2011; 11:1.
118. Sangeetha MK, Balaji HR, Gayathri V, Vasanthi HR. *Tinospora cordifolia* attenuates oxidative stress and distorted carbohydrate metabolism in experimentally induced type 2 diabetes in rats. J Nat Med, 2011; 65:544-550.
119. Chougale AD, Ghadyale VA, Panaskar SN, Arvindekar AU. Alpha glycosidase inhibition by stem extract of *Tinospora cordifolia*. J Enzyme Inhib Med Chem, 2009; 24:998-1001.



120. Agarwal VS. 1997. Drug Plants of India. Kalyani Publishers, Ludhiana; First edition (Vol. II) 1997. p- 688.
121. Anonymous. Indian Herbal Pharmacopoeia. A joint Publication of Regional Research Laboratory, Jammu tawi and Indian Drug Manufacturer's Association, Mumbai,1998a; 1: 99-104; 156-64.
122. Anonymous. Bhavamishra. Bhavaprakasha Nigantu. Chaukhamba Bharti Academy, Varanasi 1998b. p- 270.
123. Gupta RS and Sharma A. The Antifertility effect of *Tinospora cordifolia* (Willd.) Stems extracts in male rats. Indian J Exp Biol. 2003; 41: 885-9.
124. Dhuley JN. Effect of some Indian herbs on macrophage functions in ochratoxin a treated mouse. J Ethnopharmacol.,1997; 58(1):15-20.
125. Jagetia GC and Rao SK. Evaluation of cytotoxic effects of dichloromethane extracts of guduchi *Tinospora cordifolia* Miers ex Hook f & Thoms on cultured HeLa Cells. Evidence based Complement Alternat Med., 2006; 3 (2):267-272.
126. Sarma DN, Khosa RL, Chansauria JP and Sahai M. Antiulcer activity of *Tinospora cordifolia* Miers and *Centella asiatica* Linn. Extracts. Phytother Res, 1995; 9: 589.
127. Mainzen S, Prince P, Menon VP and Gunasekaran G. Hypolipidaemic action of *Tinospora cordifolia* roots in alloxan diabetic rats. J Ethnopharmacol 1999; 64: 53-7.
128. Stanley M, Prince P, Menon VP and Gunasekaran G. Hypolipidaemic action of *T. cordifolia* roots in alloxan diabetic rats. Journal of Ethnopharmacol 1999; 64: 53-7.
129. Singh N, Singh SM and Shrivastava P. Immunomodulatory and antitumor actions of medicinal plant *Tinospora cordifolia* are mediated through activation of tumor-associated macrophages. Immunopharmacol Immunotoxicol, 2005; 26:145-62.
130. Chaudhary R, Jahan S and Goyal PK. Chemopreventive potential of a medicinal plant *Tinospora cordifolia* on skin carcinogenesis in mice. J Environ Pathol Toxicol Oncol., 2008; 27(3): 233-243.
131. Rai M and Gupta SS. Experimental evaluation of *Tinospora cordifolia* (guduchi) for dissolution of urinary calculi. J Res Ind Med, 1967; 2(1):115-16.
132. Lumba SP, Parmar TL, Bali H and Lumba R. Role of Septilin in Fungal Otitis Externa, Chronic sinusitis, Chronic Tonsillitis and Chronic Suppurative Otitis Media. Probe XXII: 1983; 3: 178-80.
133. Purandare H and Supe A. Immunomodulatory role of *Tinospora cordifolia* as an adjuvant in surgical treatment of diabetic foot ulcers: a prospective randomized controlled study. Indian J. Med Sci., 2007; 61(6): 347-35.

134. Agarwal A, Malini S, Bairy KL and Rao MS. Effect of *Tinospora cordifolia* on learning and memory in normal and memory deficit rats. Indian J Pharmacol.,2002; 34: 339-49.
135. SRao PR, Kumar VK, Viswanath RK and Subbaraju GV. Cardioprotective activity of alcoholic extract of *Tinospora cordifolia* in ischemia-reperfusion induced myocardial infarction in rats. Bio Pharm Bull, 2005; 28: 2319-22.
136. Sarma DNK, Khosa RL, Chansauria JPN and Ray AK. The effect of *Tinospora cordifolia* on brain neurotransmitters in stressed rats. Fitoterapia, 1995; 67(5): 421-2.
137. Bafna PA and Balaraman R. Anti-ulcer and anti-oxidant activity of pepticare: A herbo mineral formulation. Phytomed. 2005; 12; 264-70.
138. Mehrotra R, Katiyar CK and Gupta AP. Hepato protective compositions and composition for treatment of conditions related to hepatitis B and E infection. US patent 749296. 2000.
139. Chopra RN, Nayar SL and Chopra IC.1956. Glossary of Indian Medicinal plants. CSIR. New Delhi.
140. Patel SR, Goyal RK and Shah DS. Studies on the pharmacological effects of *Tinospora cordifolia*. J Res Ind Med., 1977; 13(2):46-57.
141. Gulati OD and Pandey DC. Anti-inflammatory activity of *Tinospora cordifolia* Rheumatism. 1982; 17: 76-83.
142. Gulati OD. Clinical trial of *Tinospora cordifolia* in Rheumatoid Arthritis. Rheumatism. 1980; 15: 143-8.
143. Jana U, Chattopadhyay RN and Shw BP. Preliminary studies on anti-inflammatory activity of *Zingerber officinale* Rosc. *Vitex negundo* L. and *Tinospora cordifolia* (Willd) Miers in albino rats. Indian J Pharm., 1999; 31: 232-33.
144. Faheem A, Ali M and Shokat RM. Anti-inflammatory activity and the standardization of *Tinospora cordifolia* stem bark. Traditional system of medicine. New Delhi: Narosa Publishing House, 2006; 414-7.
145. Rege, N.N. Thatte, U.M. and Dahanukar, S.A. Adaptogenic properties of six rasayana herbs used in Ayurvedic medicine. Phytotber Res, 1999; 13(4):275-91.
146. Kapil A and Sharma S. Immunopotentiating compounds from *Tinospora cordifolia*. J. Ethopharmacol, 1999; 58(2):89-95.
147. Pahadiya S and Sharma J. Short communication: Alteration of lethal effects of gamma rays in Swiss albino mice by *Tinospora cordifolia*. Phytotherapy Research, 2003; 17: 552-554.

148. Dhingra D and Goyal PK. Evidences for the involvement of monoaminergic and Gabaergic system in antidepressant-like activity of *Tinospora cordifolia* in mice. Indian J Pharm Sci, 2008; 70 (6): 761-7.
149. Rege NN, Nazareth HM, Bapat RD and Dahanukar SA. Modulation of immunosuppression in obstructive jaundice by *Tinospora cordifolia* Indian J Med Res, 1989; 90: 478-83.
150. Gupta R, Sharma V. Ameliorative effects of *Tinospora cordifolia* root extract on histopathological and biochemical changes induced by aflatoxin-b (1) in mice kidney. Toxicol Int., 2011; 18:94-98.
151. Sharma V, Pandey D. Beneficial effects of *Tinospora cordifolia* on blood profiles in male mice exposed to lead Toxic Int, 2010; 17:12-17.
152. Upadhyay PR, Sharma V, Anita KV. Assessment of the multifaceted immunomodulatory potential of the aqueous extract of *Tinospora cordifolia*. Res J Chem Sci, 2014; 1:71-79.
153. Diwanay S, Chitre D, Patwardhan B. Immunoprotection by botanical drug on experimental metastasis. J ethanopharmacol, 2004; 90:223-237.
154. Narayanan AS, Raja SS, Ponmurugan K, Kandekar SC, Maripandi A. Antibacterial activity of selected medicinal plant against multiple antibiotic resistant uropathogens: A study from Kolli Hills, Tamilnadu, India. Benef Microbes, 2011; 2:235-243.
155. Jayachandran R, Xavier TF, Anand SP. Antibacterial activity of stem extracts of *Tinospora cordifolia* (willd.) Hook. F. and Thoms. Anc Sci Life 2003; 22:40-43.
156. Ambekar DH, Khante BS, Chandak BR, Titare AS, Baralkar SS, Aghadte SN. Screening of antibacterial potential of some medicinal plants from Melghat forest in India. Afr J Trad Comp Altern Med, 2009; 6:228-232.
157. Stanely P, Menon VP. Antioxidant action of *Tinospora cordifolia* root extract in alloxan diabetic rats", Phytother Res 2001; 15:213- 218.
158. Stanely P, Menon VP. Hypoglycaemic and hypolipidemic action of alcohol extract of *Tinospora cordifolia* roots in chemical induced diabetes in rats. Phytother Res 2003; 17:410-413.
159. Kapur P, Wuttke W, Jarry H. Beneficial effects of beta-ecdysone on the joint epiphyseal cartilage tissue and trabecular bone in ovariectomized rats. Phytomedecine, 2010; 17:350-355.
160. Desia VA, Kamat JP, Sainis KB. Proc. Indian Acad. Sci. 2002; 114, 713.
161. Goel HC, Kamat IP, Rans SV. 202. Indian, J. Exp. Biol, 40, 727.