

International Journal of Scientific Research and Reviews

Preliminary Phytochemical Screening of Different Solvent Extracts of Selected Tropical Fruits

¹Vinitha S Babu * and ²Anice Kurian M

¹Department of Botany University of Kerala Kariavattom Thiruvananthapuram-695581 Kerala India

E-mail: vinithasbabu55@gmail.com Phone: +919061575820

²Head of the Department Department of Botany Devamatha College Kuravilangad

Kottayam-686633 Kerala India E-mail: anicebabu@gmail.com Phone: +919446272732

ABSTRACT

The present study evaluates the phytochemical constituents of selected tropical fruits viz, rambutan (*Nepheliumlappaceum*), mango (*Mangiferaindica*), avocado (*Perseaamericana*), and jackfruit (*Artocarpusheterophyllus*). Preliminary phytochemical compositions were carried out for the hexane, butanol, ethanol, chloroform and aqueous extracts. Solvents were taken based on increasing order of their polarity. Qualitative phytochemical analysis of these fruit extracts confirms the presence of various phytochemicals like alkaloids, carbohydrates, phenol, glycosides, terpenoids, flavonoids, saponins, proteins, steroids and tannins.

KEY WORDS: Fruits, Photochemical, Carbohydrates, Extracts

***Corresponding Author**

Vinitha S Babu *

MPhil Scholar Department of Botany

University of Kerala Kariavattom

Thiruvananthapuram-695581 Kerala India

E-mail: vinithasbabu55@gmail.com Phone: +919061575820

INTRODUCTION

Fruits constitutes an important part in our diet. The consumption of fruit is not just for a taste and it has become a concern of our health due to the nutrient contents. Fruits feature considerable amounts of micronutrients such as vitamins, minerals, dietary fibres and other secondary metabolites as bioactive compounds¹. This evidence shows the importance of several nutrients in fruits for human health. Consumption of fruits rich in several bioactive compounds has led to the reduction of several non-communicable diseases such as certain types of cancer, inflammation, cardiovascular diseases, cataracts, macular degeneration, and neurodegenerative disease². Recently, consumption of tropical fruit is increasing observed based on domestic and international market trends due to growing recognition of its nutritional and therapeutic value.

Photochemical constitute one of the most numerous and the widely distributed group of substances found in fruits, vegetables, grains and plants. Colourful fruits and vegetables contain hundreds of phytochemicals that work together with nutrients to promote health and prevent diseases³. The most important bioactive compounds are alkaloids, flavonoids, tannins and phenolic compounds⁴. They can have different mechanisms of actions in the body, including antioxidant effects, modulation of detoxification enzymes, stimulation of immune system, modulation of hormone metabolism, antibacterial and antiviral effect^{5,6,7}. During the last two decades, there has been a search for new plant derived drugs containing the medically useful phytochemicals. The phytochemical research based on ethno-pharmacological information is generally considered as an effective approach in the discovery of new therapeutic and novel agents. Scientists have identified thousands of phytochemicals, although only a small fraction has been studied closely. However, a key obstacle, which has hindered the acceptance of the alternative medicines in the world, is the lack of documentation and stringent quality control. There is a need for documentation of research work carried out on traditional medicines⁸. With this backdrop, it becomes extremely important to try towards standardization of the plant material to be used as medicine. The process of standardization can be achieved by stepwise pharmacognostic studies⁹. These studies help in identification and authentication of the phytochemicals. Correct identification and quality assurance of the starting materials are essential pre-requisites to ensure reproducible quality of natural medicine which will contribute to its safety and efficacy. Natural phytochemicals at the low levels present in fruits and vegetables offer health benefits, but antioxidant dietary supplement compounds may not be effective or safe when consumed at higher doses even in a pure dietary supplement form. Generally, taking higher doses increases the risk of toxicity¹⁰. Despite the numerous medicinal uses attributed to these fruits, there had been no documented evidence to our knowledge. Hence, the

objective of the present study is to evaluate the phytochemical constituent present in selected tropical fruits.

MATERIALS AND METHODOLOGY

Collection of fruit material

Selected fruits for the study were purchased from a local market in the month of May 2014. The fruits were thoroughly washed, shade dried, homogenized to fine powder and stored in air tight bottles.

Chemicals, Reagents and Solvents

All chemicals, reagents and solvents used during the experimentation were of analytical grade.

Preparation of Fruit Extracts

10 g of powdered samples were weighed and mixed with 100 ml of five different solvents (methanol, ethanol, acetone, chloroform and distilled water) in conical flasks and kept in rotatory shaker at 150 rpm for 24 hours. After 24 hours it was filtered with Whatman No.1 filter paper. The filtrates were evaporated in a hot air oven at 40⁰ C until dry. The extracts were stored in sample bottles at 40⁰ C prior to use.

Preliminary phytochemical screening:

The different solvent extracts of fruits rambutan (*Nephelium lappaceum*), mango (*Mangifera indica*), avocado (*Persea americana*), and jackfruit (*Artocarpus heterophyllus*) were used to screen the phytochemicals like reducing sugar, tannins, alkaloids, saponins, amino acids, flavonoids, phenolic compounds, sugar, sterols, terpenoids and glycosides by standard method¹¹.

Reducing sugar:

Fruit extracts were treated with 2 mL of Fehling's reagent; 3 mL of water was added and allowed to boil. The development of red orange colour indicated the presence of reducing sugar.

Tannins:

The test extracts were divided into two equal portions. Sodium chloride solution was added to one portion of the extract and 1% Gelatin solution to a second portion. The development of white precipitate indicated the presence of tannins. Positive tests were confirmed by the addition of ferric chloride solution to the extract and resulted in a characteristic blue, black or green colour.

Alkaloids:

The test extracts were taken with 2N hydrochloric acid. The aqueous layer was formed. It was decanted and to which were added one or two drops of Mayer's reagent. The test content changed into white turbidity or precipitated in positive reaction.

Saponins:

The test extracts were mixed with water and shaken well. The test solution changed into foamy leather indicated the presence of saponins.

Amino acids:

The test extracts were mixed with small quantity of ninhydrin. Formation of blue colour confirmed the presence of amino acid.

Flavonoids:

The test extract in alcohol mixed with a bit of magnesium and one or two drops of concentrated hydrochloric acid and heated. The test solution changed into red or orange red colour in the presence of flavonoids.

Phenolic compounds:

The test extract in alcohol was taken with a bit of magnesium and one drop of neutral ferric chloride. Change of intense colour in the test content, showed positive result for phenolic compounds.

Sugar:

The test extracts were mixed with minimum quantity of anthrone and few drops of concentrated sulphuric acid and heated. Change of colour from green to purple showed the presence of sugar.

Sterol:

The test extracts were mixed with minimum quantity (< 1 mL) of chloroform, 3 to 4 drops of acetic anhydride and one drop of concentrated sulphuric acid. The test content in purple colour changed into blue-green indicated the presence of sterol. The result was qualitatively determined and recorded.

Terpenoids:

The test extracts were mixed with chloroform, and concentrated H₂SO₄ was carefully added to form a layer. A reddish-brown colouration of the interface formed to show positive results for the presence of terpenoids.

Glycosides:

The test extracts were mixed with glacial acetic acid, few drops of 5% FeCl₃ and

concentrated H₂SO₄ were added reddish brown colour at the junction of the two liquid layers formed, and upper layer appeared bluish green which indicated the presence of glycosides.

RESULTS AND DISCUSSION

In the present study phytochemical screening of four tropical fruits like rambutan, mango, avocado and jackfruit were done. The phytochemical analysis conducted is presented in Tables 1 to 4. The results revealed that some of the phytochemicals analysed were present in the extracts of all the fruits. Of the eleven phytochemicals screened, sugars and amino acids were present commonly in all the studied fruits. From rambutan extracts, (Table-1) saponins, phenolic compounds and sugars were present in all the studied extracts. Aqueous extract had all the ten phytoconstituents except amino acids. From mango extracts, (Table-2) phenols and sugars were present in all the solvents. Amino acids were found in all the solvents except in the chloroform extract. Hexane extract showed the presence of all phytoconstituents except saponins and glycosides. From avocado extracts, (Table-3) amino acids and flavonoids were present in all the extracts. Ethanolic extract showed the absence of alkaloids and sugars, whereas the aqueous extract showed the absence of tannins and saponins. From jackfruit extracts, (Table-4) sugars were present in all the solvents whereas saponins were absent only in ethanolic extracts.

Table 1: Preliminary phytoconstituents of Rambutan extracts

Phytoconstituents	Solvents				
	Hexane	Butanol	Ethanol	Chloroform	Aqueous
Reducing Sugar	+	+	-	+	+
Tannins	-	-	+	-	+
Alkaloids	+	+	-	+	+
Saponins	+	+	+	+	+
Amino Acids	+	-	+	+	+
Flavonoids	-	+	-	-	+
Phenols	+	+	+	+	+
Sugars	+	+	+	+	+
Terpenes	+	+	-	+	+
Cardiac Glycosides	+	+	-	+	+
Steroids	+	-	+	+	+

Table 2: Preliminary phytoconstituents of Mango extracts

Phytoconstituents	Solvents				
	Hexane	Butanol	Ethanol	Chloroform	Aqueous
Reducing Sugar	+	+	+	-	-
Tannins	+	-	-	+	+
Alkaloids	+	-	+	-	-
Saponins	-	-	+	+	+
Amino Acids	+	+	+	-	+
Flavonoids	+	+	+	-	+
Phenols	+	+	+	+	+
Sugars	+	+	+	+	+
Terpenes	+	-	-	-	-
Cardiac Glycosides	-	-	+	-	+
Steroids	+	-	+	-	+

Table3: Preliminary phytoconstituents of Avocado extracts

Phytoconstituents	Solvents				
	Hexane	Butanol	Ethanol	Chloroform	Aqueous
Reducing Sugar	+	+	-	-	-
Tannins	+	+	+	-	-
Alkaloids	-	+	-	-	+
Saponins	+	-	+	+	-
Amino Acids	+	+	+	+	+
Flavonoids	-	+	+	+	+
Phenols	+	-	+	+	+
Sugars	+	+	-	+	+
Terpenes	+	-	+	+	+
Cardiac Glycosides	-	-	-	-	+
Steroids	+	-	+	+	+

Table4: Preliminary phytoconstituents of Jackfruit extracts

Phytoconstituents	Solvents				
	Hexane	Butanol	Ethanol	Chloroform	Aqueous
Reducing Sugar	+	-	+	-	-
Tannins	+	-	+	-	-
Alkaloids	-	-	+	-	+
Saponins	+	+	+	-	+
Amino Acids	-	+	-	+	+
Flavonoids	-	+	+	+	+
Phenols	-	-	+	-	-
Sugars	+	+	+	+	+
Terpenes	-	+	+	-	-
Cardiac Glycosides	-	-	+	+	+
Steroids	-	+	+	-	-

REFERENCES

1. Metafa, Economou A. Chemo metrical development and comprehensive validation of a solid phase microextraction/gas chromatography–mass spectrometry methodology for the determination of important free and bound primary aromatics in Greek wines. *Journal of Chromatography A*.2013; 1305: 244-258.
2. Clerici, MT and Carvalho-Silva. Nutritional bioactive compounds and technological aspects of minor fruits grown in Brazil. *Food Research International* 2011;7: 1658-1670.
3. Mahattanatawee, Kanjana, et al. Total antioxidant activity and fibre content of select Florida-grown tropical fruits. *Journal of agricultural and food chemistry*.2006; 54: 7355-7363.
4. Prior, Ronald L, Guohua Cao et al. Antioxidant phytochemicals in fruits and vegetables: diet and health implications. *Hort Science* 2000; 35: 588-592.
5. Wang, Youfa, and Qi Zhang. Are American children and adolescents of low socioeconomic status at increased risk of obesity? Changes in the association between overweight and family income between 1971 and 2002. *The American journal of clinical nutrition* 2006; 84: 707-716.

6. Luximon-Ramma, Amitabye, Theeshan Bahorun, Alan Crozier et al. Antioxidant actions and phenolic and vitamin C contents of common Mauritian exotic fruits. *Journal of the Science of Food and Agriculture* 2003;83: 496-502.
7. Babbar, Neha. Total phenolic content and antioxidant capacity of extracts obtained from six important fruit residues. *Food Research International* 2011; 44: 391-396.
8. Raghu, H. S., and P. Ravindra. Antimicrobial activity and photochemical study of *Phyllanthusemblica* Linn. *Int J Pharm Stu Res* 2010;11: 30-33.
9. Ozarkar, K. R. "Studies on anti-inflammatory effects of two herbs *Cissusquadrangularis* Linn. and *Valerianawallichii* DC using mouse model. University of Mumbai, Mumbai 2005;49:200-210.
10. Michel, Thomas, et al. Antimicrobial, antioxidant and phytochemical investigations of sea buckthorn (*Hippophaë rhamnoides* L.) leaf, stem, root and seed. *Food Chemistry* 2012;131: 754-760.
11. Evans, William Charles, Daphne Evans, and George Edward Trease. *Trease and Evans pharmacognosy*. Edinburgh; New York: Saunders/Elsevier, 2009.