

## *International Journal of Scientific Research and Reviews*

### **Identification of Major Volatile (Essential Oil) Constituents of “Carrom Seeds” and “Clove Buds”**

Sadanand S. Kadam<sup>1</sup> and Waghmare Jyotsna S.<sup>1\*</sup>

<sup>1</sup>Institute of Chemical Technology, Oils, Oleochemicals & Surfactants Technology, Mumbai. India

#### **ABSTRACT:**

The major volatile constituents of essential oil obtained from carrom seeds and clove buds have been analyzed by GC and FT-IR. Physio-chemical and phyto-chemical properties of carrom oil and clove bud oil were checked and confirmed with standard oil samples. Among carrom oil, thymol and p-cymene were the major constituents constituting 61% and 14%. In clove oil, eugenol was main compound constituting 78%. FT-IR results confirmed the presence of two major phenolic group i.e. thymol and eugenol in carrom oil and clove oil respectively.

**KEYWORDS:** Essential oil, GC, FT- IR, phyto-chemical, physio-chemical, eugenol, iso-eugenol, thymol, p-cymene, carrom seeds, clove buds, Phenolic.

#### **Corresponding Author:**

Dr. Jyotsna S. Waghmare<sup>1\*</sup>

Assistant professor, Institute of Chemical Technology,

Oils, Oleochemicals & Surfactants technology, Mumbai. India

Email id: [sksady@gmail.com](mailto:sksady@gmail.com) , [jt.waghmare@ictmumbai.edu.in](mailto:jt.waghmare@ictmumbai.edu.in).

Mob No: 9769912869

## **INTRODUCTION:**

Carron seeds and clove buds are traditional spices grown in India in majority of states. Though both the spices belong to different family still they possess almost similar medicinal properties. Carron seeds and clove buds categorize itself under family Apiaceae<sup>1</sup> and Myrtaceae<sup>2</sup> respectively. Carron seeds botanical name is *Trachyspermum ammi*<sup>1</sup> where *Trachyspermum* is genus and *ammi* is species. Botanical name of clove buds is *Syzygium aromaticum*<sup>2</sup> where *Syzygium* is genus and *aromaticum* is species. Both come under kingdom plantae. The Apiaceae family consists of almost 347 genera and 12816 species. Essential oil extracted from both the spices is a volatile oil having distinguishable organoleptic and physio-chemical characteristics. Both the oil possesses wide medicinal properties followed by wide range of compounds<sup>3</sup> useful in perfume, flavor and fragrance industry. The carron oil is majorly used in mouthwash<sup>4</sup> due to its good antimicrobial properties compared to other spices. Clove oil in turn is a natural antimicrobial<sup>5,6</sup>, antiseptic, and anti-fungal<sup>7</sup> agent. Clove oil is also one of the highest sources of nutrition as it contains manganese, vitamin C and vitamin K<sup>2</sup>. Clove oil are also a natural anesthetic the reason for which they were often used for dental treatment from centuries and are still used in some cultures to remedy toothache<sup>8</sup>. They are highly beneficial in reducing the pain of tooth<sup>9</sup>. Clove oil also possesses antibacterial properties<sup>10</sup>, thus they can be used a natural pesticides to kill bacteria. Apart from medicinal usage the seeds and buds of carron and clove are as such used as spices and herbs in preparing various edible dishes. Traditionally smoke of heated carron seeds can be sometimes used for treatment of cough. Essential oil derived from carron oil possesses high antimicrobial<sup>11</sup>, anti fungal<sup>12</sup> and insecticidal<sup>13,14</sup> activity. These pesticide properties are basically because of thymol the chief constituents of oil which largely contributes to success of oil<sup>15</sup>. Moreover this oil can also be used as natural antioxidant due to presence of phenolic compound.

## **EXPERIMENTAL SECTION:**

### **a) *Materials and Methods:***

The fresh seeds and buds of carron and clove were collected from local market in Maharashtra, India. The carron seeds were sunlight dried prior to extraction for 24 hour. After drying, seeds were stored away from moisture. In case of clove buds instead of drying and crushing they were directly used for isolation of oil.

b) ***Isolation of essential oil:***

The fresh crushed powder (100g) of carrom seeds and clove buds were hydro-distilled for 4 h at 100 °C using Clevenger apparatus. The time of distillation was decided by checking the aroma profile of oil. Based on density difference oil and water were separated in Clevenger apparatus. Collected oil was further dried using anhydrous sodium sulphate and stored in dark glass bottle until analysis.

c) ***Analysis of oil:***

The gas chromatography (GC) analysis of the essential oil was carried out on capillary column (HP-5 Agilent 19091J-413; 325 C 30 m X 320 µm X 0.25 µm). Temperature programming was 90 -190°C with 10 °C ramp/min and 190 – 290 °C with 5 °C ramp/min. Inlet temperature and detector temperature was 180 °C and 210 °C. Nitrogen was used as carrier gas at 1ml/min. FID section was used. The sample injection volume was 1.0 µl/min diluted in n-hexane, with split ratio of 100:1.

d) ***Physio-chemical analysis:***

Oil obtained from hydro-distillation was subjected to physio-chemical analysis such as refractive index, specific gravity, acid value, moisture. Further Organoleptic properties of oil were also determined such as color, physical appearance, odor, solubility. Result of Physio-chemical and organoleptic analysis of both oil are tabulated in table 1. All the analysis done was based on standard method of pharmacopeia.

e) ***Other analysis:***

Phyto-chemical analysis of both carrom and clove oil was carried out<sup>16,17</sup>. This analysis does not give any brief information about compounds of oil but it do show group of phyto-chemical compound present in oil. More so compound confirmatory test of oil is also done. Results of confirmatory and phyto-chemical analysis are presented in table 2 and 3.

f) ***Identification of the constituents:***

Identification of major volatile constituents of carrom oil and clove oil occurring in GC graph shown below was done by comparing both retention time and area of samples of, extracted oil and standard compound. Both the samples were diluted in same concentration followed by same method of programming for analyzing on GC. Few other compounds, whose standards were not available, were identified by comparing literature data from library. Calibration curve

was used for thymol, p-cymene and eugenol. Confirmation of these specific major volatile compounds present in carrom and clove oil was done by FT-IR analysis. In FT-IR analysis major (i.e. thymol and eugenol) compound were identified by confirming functional group of peaks adhering in FT-IR spectrum with standard frequency.

## RESULTS AND DISCUSSIONS:

Physio-chemical analysis followed by organoleptic properties of carrom oil and clove oil are very much similar to standard oil samples. Confirmatory test of both oil confirm presence of aldehyde and phenolic group. Phyto-chemicals analysis shows presence of terpenoids in oil. GC result presented in table, indicates the major volatile compound present in carrom oil and clove oil based on their retention time. From calibration curve it was concluded that carrom seed oil contains 61 % thymol followed by 14% p-cymene where as clove oil contains 78% of eugenol. Also, presence of thymol and eugenol were confirmed by FT-IR analysis. The entire bond present in structure of both thymol and eugenol were confirmed with standard frequency and functional group adhering in their spectrum.

**Table No. 1: Physio chemical and organoleptic analysis of Carrom seed and clove bud oil**

| Test                      | Carrom seed oil                     | Clove bud oil                                     |
|---------------------------|-------------------------------------|---|
| Moisture                  | 0.08 %                              | 0.02%   |
| Odour                     | Spicy                               | Woody spicy                                       |
| Solubility                | Soluble in solvent but not in water | Soluble in solvent and partially soluble in water |
| Physical state            | Liquid                              | Liquid  |
| Colour                    | Yellow brown                        | Light yellow                                      |
| Refractive index at 25 °C | 1.485                               | 1.579   |
| Specific gravity at 25 °C | 0.93                                | 1.04  |
| Acid value                | 1.32                                | 8.217   |

**Table No. 2: Confirmatory test of oil**

| Tests    | Carrom oil | Clove oil |
|----------|------------|-----------|
| Aldehyde | Present    | Present   |
| Ketone   | Absent     | Absent    |
| Phenol   | Present    | Present   |
| Ester    | Absent     | Absent    |
| Alcohol  | Absent     | Absent    |

**Table No. 3: Phyto-chemical analysis of oil**

| Tests      | Carrom oil | Clove oil |
|------------|------------|-----------|
| Terpenoids | Present    | Present   |
| Saponin    | Present    | Absent    |
| Tannin     | Absent     | Present   |

**Table No. 4: GC results of carrom oil**

| Sr. no | RT (min) | Constituents        |
|--------|----------|---------------------|
| 1      | 10.138   | $\beta$ -pinene     |
| 2      | 11.531   | p-cymene            |
| 3      | 12.516   | $\gamma$ -terpinene |
| 4      | 19.157   | Thymol              |

**Table No. 5: GC results of clove oil**

| Sr. no | RT (min) | Constituents    |
|--------|----------|-----------------|
| 1      | 20.978   | Eugenol         |
| 2      | 22.436   | caryophyllene   |
| 3      | 23.243   | Iso eugenol     |
| 4      | 24.875   | Eugenol acetate |

Table No. 6: FT-IR results of Carrom oil

For Thymol

| Sr.no | Bond            | Functional group            | Frequency, $\text{cm}^{-1}$ |
|-------|-----------------|-----------------------------|-----------------------------|
| 1     | O-H stretch     | Phenol                      | 3462.22                     |
| 2     | C-H stretch     | Aromatic ring               | 2960.73                     |
| 3     | C-C stretch     | Aromatic ring               | 1417.68                     |
| 4     | C-H substituted | Meta disubstituted aromatic | 808.17                      |

Table No. 7: FT-IR results of Clove oil

For Eugenol

| Sr.no | Bond            | Functional group            | Frequency, $\text{cm}^{-1}$ |
|-------|-----------------|-----------------------------|-----------------------------|
| 1     | O-H stretch     | Phenol                      | 3510.45                     |
| 2     | C-H stretch     | Aromatic ring               | 2937.59                     |
| 3     | C-C stretch     | Aromatic ring               | 1512.19                     |
| 4     | C-H substituted | Para disubstituted aromatic | 794.67                      |
| 5     | C-O stretch     | Ether                       | 1265.3                      |
| 6     | =C-H bend       | Alkene                      | 1010                        |

Figures:




Figure No. 1: GC graph of carrom oil




Figure No. 2: GC graph of clove bud oil




Figure No. 3: FT-IR spectrum of carrom oil




Figure No. 4: FT-IR spectrum of clove bud oil

## CONCLUSION:

From the above work it can be concluded that hydro-distillation is efficient technique to extract essential oil from spices considering yield, purity and aroma of oil. Also, carrom oil and clove oil are one of the natural sources to isolate thymol and eugenol.

## ACKNOWLEDGEMENT:

Authors are thankful to department of oils, oleo-chemicals and surfactant technology, Institute of chemical technology, (Mumbai) India and for providing appropriate facilities to carry to this entire work.

## REFERENCES:

1. Kamal J, Devi N, Thakur N, Tomar S et al., *Trachyspermum ammi* (ajwain): a comprehensive review. *International research journal of pharmacy*. 2012; 3(5): 133–138.



2. Bhowmik D, Kumar KP, Yadav A et al. Recent Trends in Indian Traditional Herbs *Syzygium aromaticum* and its Health Benefits. *Journal of Pharmacognosy and Phytochemistry*. 2012; 1(1): 13-22.
3. Nowak A, Kalembe D, Piotrowska M et al. Effects of thyme (*Thymus vulgaris* L.) and rosemary (*Rosmarinus officinalis* L.) essential oils on growth of *Brochothrix thermosphacta*. *African Journal of Microbiology Research*. 2013; 7(26): 3396-3404.
4. Hassan G, Sepideh A, Ashkan J & Aghi S. Chemical constituents, antimicrobial and antioxidative effects of *trachyspermum ammi* essential oil. *Journal of Food Processing and Preservation*. 2013: 1-5.
5. Chamdit S & Siripermpool P. Antimicrobial Effect of Clove and Lemongrass Oils against Planktonic Cells and Biofilms of *Staphylococcus aureus*. *Mahidol University Journal of Pharmaceutical Sciences*. 2012; 39(2): 28-36.
6. Fu Y, Zu Y, Chen L et al. Antimicrobial Activity of Clove and Rosemary Essential Oils Alone and in Combination. *Phytotherapy Research*. 2007; 21: 989-994.
7. Pinto E, Silva LV, Cavaleiro C et al. Antifungal activity of the clove essential oil from *Syzygium aromaticum* on *Candida*, *Aspergillus* and dermatophyte species. *Journal of Medical Microbiology*. 2009; 58: 1454–1462.
8. Bina R, Kachhawa GR, Yadav RK et al. Phytochemical effectiveness of clove oil: A review. *International journal of research in pharmacology and pharmacotherapeutics*. 2012; 1(2): 230-233.
9. Rusenova N & Parvanov P. Antimicrobial activities of twelve essential oils against microorganisms of veterinary importance. *Trakia Journal of Sciences*. 2009; 7(1): 37-43.
10. Ali HS, Kamal M & Mohamed SB. In vitro clove oil activity against periodontopathic bacteria. *Journal of Science and Technology*. 2009; 10(1): 1-7.
11. Hassanshahian M, Bayat Z, Saeidi S et al. Antimicrobial activity of *Trachyspermum ammi* essential oil against human bacterial. *International journal of Advanced Biological and Biomedical Research*. 2014; 2(1): 18-24.
12. Saad RE & Soad MA. Antifungal activity of some essential oil and their major chemical constituents against some phytopathogenic fungi. *Journal of pest control and environmental studies*. 2005; 13(1): 61 -72.

13. Chaubey MK. Insecticidal activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella sativa* (Ranunculaceae) essential oils against stored-product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *African Journal of Agricultural Research*. 2007; 2(11): 596-600.
14. Szczepanik M, Zawitowska B & Szumny A. Insecticidal activities of *Thymus vulgaris* essential oil and its components (thymol and carvacrol) against larvae of lesser mealworm, *Alphitobius diaperinus* Panzer (Coleoptera:Tenebrionidae). *Allelopathy Journal*. 2012; 30(1): 129-142.
15. Riccioni L & Orzali L. Activity of Tea Tree (*Melaleuca alternifolia*, Cheel) and thyme (*Thymus vulgaris*, Linnaeus.) Essential Oils against Some Pathogenic Seed Borne Fungi. *Journal of Essential Oil Research*. 2011; 23: 43-47.
16. Ayoola GA, Coker HA, Adesegun SA et al. Phytochemical Screening and Antioxidant Activities of Some Selected Medicinal Plants Used for Malaria Therapy in Southwestern Nigeria. *Tropical journal of Pharmaceutical Research*. 2008; 7(3): 1019-1024.
17. Ayoola G A, Lawore FM, Adelowotan T et al. Chemical analysis and antimicrobial activity of the essential oil of *Syzigium aromaticum* (clove). *African Journal of Microbiology Research*. 2008; 2(1): 162-166.