

International Journal of Scientific Research and Reviews

A Study To Compare The Effect Of Slow Breathing Exercise And Jacobson's Progressive Muscular Relaxation On Blood Pressure , Heart Rate And Respiratory Rate In Patients With Primary Hypertension : An Interventional Study

¹Kalyanee Jivrajani, ²Yagnik Dave and Vidhi Talati*³

¹Student(M.P.T Cardio-Respiratory) At Shri K.K. Sheth Physiotherapy College , Rajkot

²Assistant Professor (M.P.T. Cardio-Respiratory)At Harivandana Physiotherapy College Rajkot

^{3*}Assistant Professor (M.P.T. Cardio-Respiratory) At Shri K.K. Sheth Physiotherapy College Rajkot

ABSTRACT

Hypertension means persistent increase in both SBP and DBP. Breathing exercise controlled at 6 rate/min are known as slow breathing exercises .Jacobson's progressive muscle relaxation technique is a relaxation technique in which a person first tenses and releases major muscle of the body in a systematic order, usually beginning at the distal body parts and progressing proximally . To compare the effect of slow breathing exercise and Jacobson's progressive muscular relaxation on HR ,RR and BP in patients with primary hypertension.

30 subjects who fulfilled the selection criteria were selected for the study with age group 35-65 years. Subjects divided into 2 groups Group A - Slow breathing exercise , Group B Jacobson's Progressive Muscular Relaxation Technique for 1 week. Pre and post treatment evaluation of BP, HR and RR were taken and results were compared.

Result of present study shows a significant difference for Pre-Post treatment values of all outcome measures($p < 0.05$) for both groups. Inter group comparison shows no significant difference for HR and RR ,but significant difference ($p < 0.05$) for SBP and DBP .

This study concludes that patient receiving slow breathing exercise has more beneficial effect in SBP and DBP while there is no significant difference in HR and RR between the two groups.

KEY WORDS: Slow breathing exercises, Jacobson's progressive muscular relaxation , Essential Hypertension

***Corresponding author:**

Vidhi Talati

Assistant Professor

(M.P.T. Cardio-Respiratory)

At Shri K.K. Sheth Physiotherapy College , Rajkot

INTRODUCTION

Persistent increase in systemic arterial blood pressure is known as hypertension¹. Hypertension means increase in normal blood pressure both in systolic BP and diastolic BP². It is ranked as fourth top most disease on the basis of its prevalence. Hypertension is a major health problem and biggest of the challenges of the 21st century³.

It has been well documented as a major risk factor for cardiovascular disease, renal disease, Stroke, heart failure, peripheral artery disease. There are two main types of hypertension, "Primary" hypertension, also known as essential or idiopathic hypertension and "Secondary" hypertension which has some underlying cause. Essential hypertension is the most prevalent type, affecting 90-95% of hypertensive patients. With this type of hypertension, there is no single identifiable cause. The pathogenesis of hypertension is multifactorial and can trigger the risk as genetic factors which play an important role⁴.

Classification of hypertension for adults ages 18 and older has been provided by the Eighth Report of the Joint National Committee of High Blood Pressure (JNC 8 Classification, 2015) i.e. stage 1 hypertension (systolic BP 140-159 or diastolic BP 90-99 mmHg), and stage 2 hypertension (systolic BP \geq 160 or diastolic BP \geq 100 mmHg)⁵

Breathing exercises controlled at 6 rate per minute (rpm) are known as slow controlled breathing exercises and are characterized by increased tidal volume (respiratory amplitude) and decreased respiratory rate². It increases baroreflex sensitivity and reduces sympathetic activity and chemoreflex activation. Slow breathing improves vagal activity and therefore decreases baseline heart rate and blood pressure⁴.

Jacobson's progressive muscular relaxation is a relaxation technique in which a person first tenses and releases major muscle of the body in a systematic order, usually beginning at the distal body parts and progressing proximally³. The main rationale for this is when the body is aware of the presence of the tension, it will respond by triggering the muscles to relax, where the rest of the other components of the relaxation response will naturally follow which tends to cause parasympathetic dominance³.

NEED OF THE STUDY

Hypertension is a widespread health problem and has no warning signs or symptoms, and many people don't realize they have it. Despite the significance of the problem with respect to overall health; and its undesirable health consequences, high blood pressure is still not adequately controlled and far from being optimal. The side effects and cost of antihypertensive drugs have

stimulated the search for a non-pharmacological approach to control BP either as a first line or adjunctive treatment⁴.

Relaxation and stress-relieving techniques such as yoga, meditation and biofeedback have also been shown capable of lowering BP⁴. Also both the techniques are easy, non-invasive, painless, can be done at any place, less time consuming, cost effective

So the need of the study is to find the better non pharmacological treatment out of the two for the patient with primary hypertension.

AIMS AND OBJECTIVES

Aim of the study:

- To study and compare the effects of Slow breathing exercise and Jacobson's progressive muscular relaxation technique on HR, RR and B.P. in patients with primary hypertension.

Objective of the study:

- To study the effect of slow breathing exercise on blood pressure, HR and RR in patients with primary hypertension.
- To study the effect of Jacobson's progressive muscular relaxation technique on blood pressure, HR and RR in patients with primary hypertension.
- To compare the effect of Slow breathing exercise and Jacobson's progressive muscular relaxation technique on blood pressure, HR and RR in patients with primary hypertension.

HYPOTHESIS

- **Null Hypothesis** : There is no statistical significant difference in the effect of Slow breathing exercise and Jacobson's progressive muscular relaxation on B.P., HR and RR in patients with primary hypertension.
- **Experimental Hypothesis** : There is a statistical significant difference in the effect of Slow breathing exercise and Jacobson's progressive muscular relaxation on B.P., HR and RR in patients with primary hypertension.

METHODOLOGY

- STUDY DESIGN : An interventional study
- METHOD OF COLLECTION OF DATA:

Study population: Subjects with primary hypertension

Sampling method: Purposive sampling.

Sample size: 30 subjects.

Source of data: O.P.D in and around Rajkot city.

Study duration : 4 months (January to april 2019)

MATERIALS USED: (Figure 1)

- Mercury Sphygmomanometer
- Stethoscope
- Stop watch
- Pen
- Paper
- Assessment form
- Consent form
- Data collection sheet
- Couch
- Pillow
- Weighing machine
- Measure tape




Figure 1 : Materials used

CRITERIAS FOR SELECTION :

➤ Inclusion Criterias:

- 1) Age: 35-65 years²
- 2) Gender: male & female both will be included²
- 3) Patients with essential hypertension (SBP \geq 140 mmHg and DBP \geq 90 mmHg) diagnosed by a physician at least six months prior to study on stable anti-hypertensive treatment for a minimum of two months prior to the study and no change in medications during participation in the trial.⁶

➤ Exclusion Criterias :

- 1) Patients with secondary hypertension
- 2) Autoimmune diseases²
- 3) Cigarette smoking²
- 4) Alcohol consumption²
- 5) Use of oral contraceptives²
- 6) Use of neuroleptics/anti- arrhythmic ²
- 7) Use of neuroleptics/anti- arrhythmic ²
- 8) Impaired cognitive function⁸

➤ Method⁹

- On day 1- 30 Volunteers were briefed about the protocol and written consent , assessment and baseline readings of HR, RR and BP were taken. On day 2 Training was provided to Group A(n=15) for slow breathing exercise and Group B(n=15) for Jacobson's progressive relaxation technique . from 3rd day to 9th day Study duration of 20 mins twice daily was carried out and on day 9- post test readings were taken and the data obtained was analysed statistically.

PROCEDURE FOR SLOW BREATHING EXERCISE: (Figure 2)

- One has to sit comfortably in an easy and steady posture on a fairly soft seat
- Patients were asked to take deep breath in through nose and than were asked to exhale completely through mouth.
- I: E = 4:6
- This completes 1 cycle of slow breathing¹⁰.



Figure 2: Slow Breathing Exercise

PROCEDURE FOR JACOBSON'S PROGRESSIVE MUSCULAR RELAXATION: (Figure 3)

- In Jacobson's progressive muscle relaxation technique person first tenses and releases major muscle of the body in a systematic order, usually beginning at the distal body parts and progressing proximally³.
- The tensing time was kept for 5 seconds followed by relaxation time for 10 seconds.
- This constituted 1 session of 20 minutes.¹¹



Figure 3: Jacobson's Progressive Muscular Relaxation Technique

RESULTS

- Data of 30 subjects were analysed using SPSS V.20 and MS Excel 2010.
- Within the group : paired t- test
- Between the group : unpaired t-test
- Mean age of participants
- Group A = 57.8 ± 6.08 (8 Females , 7 males)
- Group B = 58.67 ± 7.32 (11 Females , 4 males)

Table 1 : Results Of Slow Breathing Exercise

RESULTS OF SLOW BREATHING EXERCISE				
	PRE	POST	MEAN DIFF	P-VALUE
Systolic B.P	144 ± 4.65	137.20 ± 4.32	6.8	0.00
Diastolic B.P	85.33 ± 5.43	80.40 ± 5.19	4.93	0.00
Heart rate	80.86 ± 7.78	75.66 ± 8.20	5.20	0.00
Respiratory rate	20.53 ± 3.04	16.06 ± 2.37	4.46	0.00


Table 2 : Results Of Jacobson’s Progressive Muscular Relaxation Technique

RESULTS OF JACOBSON’S PROGRESSIVE MUSCULAR RELAXATION TECHNIQUE				
	PRE	POST	MEAN DIFF	P-VALUE
SYSTOLIC B.P.	145.73 ± 5.28	142.0 ± 5.23	3.73	0.00
DIASTOLIC B.P.	85.46 ± 4.74	82.0 ± 4.14	3.46	0.00
HEART RATE	79.60 ± 8.78	75.13 ± 8.74	4.46	0.00
RESPIRATORY RATE	20 ± 4.58	16.33 ± 4.28	3.66	0.00

Table 3 : Between The Groups Comparison

BETWEEN GROUP ANALYSIS				
	GROUP A	GROUP B	MEAN DIFF	P VALUE
SYSTOLIC B.P.	6.8 ± 2.36	3.73 ± 1.27	3.06	0.00
DIASTOLIC B.P.	4.93 ± 1.27	3.46 ± 1.59	1.46	0.01
HEART RATE	5.20 ± 2.07	4.46 ± 1.18	0.73	0.245
RESPIRATORY RATE	4.46 ± 1.86	3.66 ± 1.14	0.93	0.110

Graph 1 : Between Group Analysis



Were 1= SBP
 2= DBP
 3=HR
 4= RR

DISCUSSION

The result of the present study support the alternative hypothesis for SBP and DBP which shows the statistical significant difference between slow breathing exercise and jacobson's progressive muscular relaxation. However the study also accepts the null hypothesis for HR and RR which shows no statistical significant difference between slow breathing exercise and jacobson's progressive muscular relaxation technique.

In hypertensive subjects practicing slow deep breathing exercise at 6 cycles/min, the tidal volume increases to maintain minute ventilation. The reflex mechanism involves the pulmonary stretch receptors that are stimulated and afferents via vagi reaches nucleus tractus solitarius, reduces the chemoreflex sensitivity and thus might enhance the baroreflex, with an effect on reducing blood pressure and sympathetic activity as reported by Narkiewicz et al.¹²

Dr. Labiba Abd El-kader Mohamed et al in 2013 studied the Effect of slow deep breathing exercise on blood pressure and heart rate among newly diagnosed patients with essential hypertension and concluded that practicing slow deep breathing exercise decreased the systolic and diastolic BP as well as heart rate of patients with essential hypertension which supports the result of the present study.⁴ When performing progressive muscle relaxation exercises with a calm, relaxed state and full concentration of tense and relaxed muscles trained for 15 minutes then the secretion of CRH (corticotropin releasing hormone) and ACTH (adrenocorticotrophic hormone) in the hypothalamus decreases¹³.

Decreased secretion of these two hormones cause decrease in sympathetic nerve activities, so that adrenaline and noradrenaline expenditure is reduced. The decrease in adrenaline and norepinephrine results in decreased heart rate, dilated blood vessels, reduced blood vessel resistance and decreased heart pumps so that cardiac arterial blood pressure decreases¹⁴.

CONCLUSION

- Result of present study shows a significant difference for Pre-Post treatment values of all outcome measures ($p < 0.05$) for both groups. Inter group comparison shows no significant difference for HR and RR, but significant difference ($p < 0.05$) for SBP and DBP.
- So this study concludes that patient receiving slow breathing exercise has more beneficial effect in SBP and DBP while there is no significant difference in HR and RR between the two groups.

LIMITATIONS OF THE STUDY:

- Sample size was small.

- Result of this study cannot be generalized on patients with secondary hypertension (due to liver, heart, renal failure, or on individuals with recent cardiovascular events, pulmonary diseases, diabetes mellitus, neuropathies, cardiac arrhythmias, with autoimmune diseases, cigarette smoking, alcohol consumption, using oral contraceptives, using neuroleptics, anti-arrhythmic and lithium)
- The result of present study could not consider the effects of medications on hemodynamic parameters.

FURTHER RECOMMENDATIONS:

- Large sample size can be recommended for the study.
- Study can be done in secondary hypertension.
- It can be done in elderly patients with hypertension.

REFERENCES

1. K Sembulingam, PremaSembulingam, Essential of Medical Physiology 6th edition, Jaypee brothers Arterial Blood Pressure, 2013; 103: 615.
2. BalasubramanianSundaram, Purvesh J Maniyar, Varghese John P, Vijay Pratap Singh. Slow breathing training on cardio-respiratory control and exercise capacity in persons with essential hypertension- A randomized controlled trial. *Indian Journal of Physiotherapy and Occupational Therapy*. 2012; 6(2): 22-7.
3. Nisha Shinde Shinde KJ , Khatri SM , Deepali Hande,Vichare Bhushan Immediate Effect of Jacobson's Progressive Muscular Relaxation in Hypertension *Sch. J. App. Med. Sci.*; 2013; 1(2):80-85
4. Labiba Abd El-kader Mohamed, Naglaa Fawzy Hanafy, Amel Gomaa Abd El- Naby. Effect of slow deep breathing exercise on blood pressure and heart rate among newly diagnosed patients with essential hypertension. *Journal of Education and Practice*; 2013;5(4):36-45
5. Kayce Bell, June Twiggs, Bernie R. Hypertension: The Silent Killer: Updated JNC-8 Guideline Recommendations 2015.
6. Kow Fei Ping, MMed, Adlina Bakar, MMed, Sivasangari Subramaniam, PhD, Punithavathi Narayanan, MBBS, Ng Kum Keong, MBBS, Ang Ah Heong, SRN, Ong Loke Meng The impact of music guided deep breathing exercise on blood pressure control - A participant blinded randomised controlled study *Clinical Research Centre, Hospital Pulau Pinang, Georgetown, Pulau Pinang, Malaysia*. 2018

7. Shirley Telles , Sachin Kumar Sharma ,Acharya balkrishna the effect of alternate nostril breathing on blood pressure and heart rate variability in normal healthy male volunteers *Med Sci Monit Basic Res*, (2014); 20: 184-193
8. Shirley Telles, Sadhana Verma , Sachin Kumar Sharma , Ram Kumar Gupta , Acharya Balkrishnathe effect of alternate nostril breathing exercise on blood pressure and stress vigilance test *Med Sci Monit Basic Res*,; 2017; 23: 392-398
9. A.V. Turankar, S. Jain, S.B. Patel, S.R. Sinha, A.D. Joshi, B.N. Vallish, P.R. Mane & S.A. Turankar Effects of slow breathing exercise on cardiovascular functions, pulmonary functions & galvanic skin resistance in healthy human volunteers - a pilot study *Indian J Med Res* 2013; 137: 916-921.
10. Tapas Pramanik, Hari Om Sharma, Suchita Mishra, , Anurag Mishra, Rajesh Prajapati, and Smriti Singh the Immediate Effect of Slow Pace Bhastrika Pranayama on Blood Pressure and Heart Rate *THE JOURNAL OF ALTERNATIVE AND COMPLEMENTARY MEDICINE* 2009; 15(3): 293–295
11. Bushra Mushtaq, and Ajaz Ahmad Khan Jacobson Muscle Relaxatation Technique (Jpmr) (20 Min) *JOJ nursing and health care* 2018 ; 8(1).
12. Dr.Vasuki G, Dr.Sweety L.M *IOSR Journal of Dental and Medical Sciences* 2017;16(2, IV): 59-62 .
13. Chellew, K., Evans, P., Fornes-Vives, J., Pérez, G. and Garcia Banda, G. The effect of progressive muscle relaxation on daily cortisol secretion *Stress*, 18 (5); 538-544.
14. Chris Cottier, Kenneth Shapiro, Stevo Julius Treatment of Mild Hypertension With Progressive Muscle Relaxation *Arch Intern med* oct 1984; 144