

## *International Journal of Scientific Research and Reviews*

### **Oxidation of Niacinamide by sodium N-chlorobenzenesulphonamide (CAB) in acid medium catalysed by Ru(III) Ion: Kinetic and Mechanistically**

**Chandrashekar**

<sup>1</sup>Department of Chemistry, PES College of Engineering, Mandya-571401, Karnataka

#### **ABSTRACT**

The kinetics of Ru (III) Catalysed oxidation of Niacinamide by sodium-N-chlorobenzene sulphonamide (CAB) has been studied in acidic medium at 303K. The reaction shows first order dependence of the rate on [CAB] [Ru(III)] and an inverse fractional order on [H<sup>+</sup>] and fractional order shows on [NA]. The addition of halide ions and the reduction product of CAB, benzene sulphonamide and dielectric constant of the medium have no influence on the rate. Thermodynamically parameters were computed by studying the reaction at different temperatures, a mechanism consistent with observed kinetics is presented.

**KEYWORDS:** Niacinamide, Oxidation, Chloramine-B, Ruthenium catalyst.

**\*Corresponding author**

**Dr. Chandra Shekar**

Department of Chemistry

PES College of Engineering,

Mandya-571401 Karnataka

Email: [chandrashekar-pes@rediffmail.com](mailto:chandrashekar-pes@rediffmail.com)

Mob No-9742167662

## INTRODUCTION

Niacinamide is an important component of vitamin-B complex group. It is water soluble vitamin and is part of the vitamin B group. It is also known as prevents pellagra. This disease is mainly found in those areas where maize is the main dietary component as the grain is deficient in niacin. Niacinamide also has anti-diabetogenic, antioxidant and anti-inflammatory properties<sup>1</sup>, anxiolytic (anti-anxiety) agent<sup>2</sup>. and also acts as a chemo and radio sensitizing agent by enhancing tumour blood, thereby reducing tumour hypoxia. Niacinamide is an activator of sirtuins but it inhibits at higher doses. Sharma ashok et al have reported Nicotinamide and Isonicotinamide were oxidized using permanganate ion in acidic medium<sup>3</sup>. Chandrashekar et al have reported oxidation of Nicotinamide by bromamine-T in acidic medium catalysed by Ru (III) ion<sup>4</sup>. Hence, the oxidation of niacinamide adds much to the knowledge of chemistry. These facts promoted us to undertake the study of kinetics of oxidation of Niacinamide by CAB in acidic media catalysed by ruthenium with a view to elucidate the reaction mechanism.

## EXPERIMENTAL

Chloramine-B (CAB) was prepared using standard method and its purity checked iodometrically and through IR and NMR spectral data<sup>5</sup> allowance was made for the amount of HCl present in the catalyst solution, while preparing reaction mixtures for kinetics runs. All other chemicals used were of accepted grade of purity. A constant ionic strength of the reaction mixtures was maintained by adding concentrated sodium perchlorate (NaClO<sub>4</sub>). A solution of RuCl<sub>3</sub>.3H<sub>2</sub>O (Arora matthey) in 0.5 M HCl was prepared and used as catalyst solution. Triply distilled water was utilized for preparing aqueous solutions.

## KINETIC MEASUREMENT

The kinetic runs were performed under pseudo-first order condition of [Niacinamide] >> [CAB]<sub>0</sub> at 303K. Mixtures contain requisite amount of substrate, oxidant, NaClO<sub>4</sub>, Ru(III) and HCl were taken in stoppered pyrex glass tube, whose outer surface was coated black to eliminate the photochemical effects. A required quantity of water was added to maintain constant total volume for all runs. The reaction vessel was thermostated in a water bath set at a temperature 303K, to this solution a measured amount of pre-equilibrated CAB solution was added to give a known concentration. The progress of the reaction was monitored iodometrically for two half-lives by withdrawing aliquots of the reaction mixture at regular time intervals. Under pseudo-first order conditions rate constant *k'* were reproducible within ±3%. The regression analysis of experimental data was carried out on an origin 5.0.

## STOICHIOMETRY

Reaction mixtures containing different composition of Niacinamide and CAB was equilibrated at 303K in acidic medium catalysed by Ru(III) ion for 24 hours. The analysis showed that one mole of CAB reacted with same mole of substrate is illustrated as in equation (1)



Niacinamide CAB

The reaction product benzene sulphonamide ( $\text{PhSO}_2\text{NH}_2$ ) was identified by TLC using light petroleum ether-chloroform-butanol (2:2:1 v/v) as the solvent system for a sending ( $R_f=0.88$ ), the 6-hydroxyl niacinamide present in the reaction mixture was identified with authenticated sample by TLC method. Further it was confirmed by conventional ferric chloride test<sup>6</sup>. The evolved  $\text{CO}_2$  was detected by the conventional lime water test. Attempts to quantitative measure of the  $\text{CO}_2$  evolved were unsuccessful. The GC-MS data for 6-hydroxy niacinamide obtained on a 17A shimadzu spectrometer showed a molecular ion peak at 133amu (figure.1), clearly confirming the formation of 6-hydroxy Niacinamide.




Fig 1: GC Mass Spectra of 6- Hydroxy Niacinamide.

## RESULTS

### *Effect of reactant concentration on the rate*

Under pseudo-first order conditions of  $[\text{Niacinamide}] \gg [\text{CAB}]_0$  plots of  $\log [\text{titre value}]$  versus time are linear ( $r=0.9980$ ) indicating a first order dependence of rate on  $[\text{CAB}]$ . The pseudo-first order rate constants  $k'$  is given in table 1. The values of  $k'$  remain unaffected with a change in  $[\text{CAB}]$ . The rate increased with increase in  $[\text{Niacinamide}]$  (table 1) and a plot of  $\log k'$  versus  $\log [\text{Niacinamide}]$  was linear with fractional slopes indicating the fractional order dependence of rate on  $[\text{Niacinamide}]$ .

Table-1 Effects of varying reactant concentrations on the reaction rate  $[\text{HCl}] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$ ;  $[\text{Ru(III)}] = 1.243 \times 10^{-6} \text{ mol dm}^{-3}$ ; Temp. = 303 K ;  $\mu = 0.2 \text{ mol dm}^{-3}$

$[\text{CAB}] \times 10^4$ $\text{mol dm}^{-3}$	$[\text{Niacinamide}] \times$ $10^3 \text{mol dm}^{-3}$	$k' \times 10^6$ $\text{sec}^{-1}$
1.15	2.0	3.00
1.61	2.0	2.90
2.00	2.0	2.80
2.53	2.0	2.85
2.99	2.0	3.15
2.00	1.0	1.25
2.00	1.5	2.00
2.00	2.0	2.80
2.00	3.0	3.70
2.00	4.0	4.60
2.00	5.0	5.40

### Effect of acid on the rate

The reaction was studied with varying  $[\text{HCl}]$  at constant  $[\text{CAB}]$ ,  $[\text{Niacinamide}]$ ,  $[\text{Ru (III)}]$ , ionic strength and temperature. The rate of reaction decreased with increase in  $[\text{HCl}]$  ( $r = -0.9990$ ), the plot of  $\log k'$  versus  $\log[\text{HCl}]$  was linear with negative slope equal to less than unity indicating inverse fractional order in HCl as shown in figure 2.




FIG.2. Plot of  $\log k'$  versus  $\log [\text{HCl}]$

### Effect of $[\text{H}^+]$ on the rate

At constant  $[\text{CAB}]$ ,  $[\text{Niacinamide}]$  and  $[\text{Cl}^-]$ , the rate of reaction decreased with increase in  $[\text{H}^+]$  ( $r = -0.9910$ ). The plot of  $\log k'$  versus  $\log [\text{H}^+]$  was linear with negative slope equal to less than unity ( $-0.62$ ) indicating an inverse fractional order in  $[\text{H}^+]$  table 2.

Table:2 Effect of varying  $[H^+]$  ion on the rate of reaction  $[NA] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$ ;  $[CAB]_0 = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$ ;  $[Ru(III)] = 1.243 \times 10^{-6} \text{ mol dm}^{-3}$  Temp. = 303K;  $\mu = 0.2 \text{ mol dm}^{-3}$

$[H^+] \times 10^4 \text{ (mol dm}^{-3}\text{)}$	$k' \times 10^6 \text{ (sec}^{-1}\text{)}$
2.4	3.04
4.8	2.05
7.2	1.75
9.6	1.40
12.0	1.20
13.4	1.15

### *Effect of Ru(III) catalysed reaction on the rate*

The rate increased with increase in  $[Ru(III)]$  and plots of  $\log k'$  versus  $\log [Ru(III)]$  was linear with unit slope indicating a first order dependence of rate on  $[Ru(III)]$  table.3.

Table : 3. Effects of varying  $[RuCl_2]$  on the reaction rate  $[NA] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$ ;  $[CAB]_0 = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$ ;  $[HCl] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$ ; Temp. = 303K;  $\mu = 0.2 \text{ mol dm}^{-3}$

$[RuCl_2] \times 10^6 \text{ mol dm}^{-3}$	$k' \times 10^6 \text{ sec}^{-1}$
3.729	2.75
7.458	7.40
11.187	16.24
14.187	20.12
18.640	27.08

### *Effect of halide ion on the rate on the rate*

Addition of  $Cl^-$  or  $Br^-$  ions in the form of NaCl or NaBr at constant  $[H^+]$  did not affect the rate, suggesting that chloride or bromide ions were not involved in the reaction rate.

### *Effect of benzenesulphonamide and ionic strength on the rate*


The addition of benzenesulphonamide had no effect on the rate indicating that it is not involved in a pre-equilibrium to the rate determining step. The variation of ionic strength of the medium using  $NaClO_4$  had no effect on the rate.

### *Effect of dielectric constant and temperature on the rate*



The variation of the solvent composition using methanol (5.0-20% v/v) did not affect the rate. The reaction was studied at varying temperatures 303-318K. The activation parameters namely energy of activation ( $E_a$ ), enthalpy of activation ( $\Delta H^\ddagger$ ), entropy of activation and free energy of activation ( $\Delta G^\ddagger$ ), were obtained from the Arrhenius plots of  $\log k'$  versus  $1/T$ .

## MECHANISM


Chloramine-B ( $\text{PhSO}_2\text{NCINa}$ ) like chloramine-T behaves as a strong electrolyte in aqueous solutions forming different species as shown in Equation 2-6<sup>7,8,9</sup>.



In acid solutions, the probable oxidizing species are the free acid  $\text{PhSO}_2\text{NHCl}$ ,  $\text{PhSO}_2\text{NHCl}_2$ ,  $\text{HOCl}$  and  $\text{H}_2\text{O}^+\text{Cl}$ . The involvement of  $\text{PhSO}_2\text{NCl}_2$  in mechanism leads to a second-order rate law according to equation (5), which is contrary to the experimental observations. The monohaloamines can be further protonated at  $\text{p}^{\text{H}} < 2$  as in equation (7) and (8) for chloramine-T and chloramine-B respectively.<sup>10,11</sup>




Therefore in acidic conditions, for chloramine-B,  $\text{PhSO}_2\text{NHCl}$  is expected to protonate as follows.




Electronic spectral studies have shown that coordination species such as  $[\text{RuCl}_5(\text{H}_2\text{O})]^{-2}$ ,  $[\text{RuCl}_4(\text{H}_2\text{O})_2]^-$ ,  $[\text{RuCl}_3(\text{H}_2\text{O})_3]$ ,  $[\text{RuCl}_2(\text{H}_2\text{O})_4]^+$  and  $[\text{RuCl}_5(\text{H}_2\text{O})_5]^{2+}$  do not exist in the aqueous solution of  $\text{RuCl}_3$ . Ruthenium (III) however exists in the following ligand substitution equilibrium in acid medium



In the present study the oxidation of Niacinamide in the presence of  $\text{Ru(III)}$  as catalyst, the inverse fractional order in  $[\text{H}^+]$  suggest that, the deprotonation of  $\text{PhSO}_2\text{N}^+\text{H}_2\text{Cl}$  in step (i) results in the formation of regeneration of  $\text{PhSO}_2\text{NHCl}$ . A retardation by the added benzenesulphonamides ( $\text{PhSO}_2\text{NH}_2$ ) i.e. an inverse fractional order on  $[\text{PhSO}_2\text{NH}_2]$  indicates hydrolysis of monobromamine  $[\text{PhSO}_2\text{NHCl}]$  to form  $\text{HOCl}$  in step (ii) which act as the active species in fast pre-equilibrium step. The reaction rate shows fractional order in  $\text{Ru(III)}$  concentration and first order on  $[\text{Niacinamide}]$ . Based on the preceding discussion a mechanism scheme 1 is proposed to account for the experimental observation.



**Scheme – 1**

$$\text{Rate} = - \frac{d[\text{CAB}]_t}{dt} = k_3 [\text{X}] [\text{Ru(III)}] \quad \dots (11)$$

From step (ii)

$$[\text{PhSO}_2\text{NHCl}] = \frac{[\text{X}]}{K_2[\text{S}]} \quad \dots (12)$$

From step (i)

$$\begin{aligned}
 [\text{PhSO}_2\text{N}^+\text{H}_2\text{Cl}] &= \frac{[\text{PhSO}_2\text{NHCl}] [\text{H}^+]}{K_1} \\
 \therefore [\text{PhSO}_2\text{N}^+\text{H}_2\text{Cl}] &= \frac{[\text{X}][\text{H}^+]}{K_1 K_2 [\text{S}]} \quad \dots(13)
 \end{aligned}$$

Total effective concentration of  $[\text{CAB}]_t$

$$[\text{CAB}]_t = [\text{PhSO}_2\text{N}^+\text{H}_2\text{Cl}] + [\text{PhSO}_2\text{NHCl}] + [\text{X}] \quad \dots (14)$$

$$[\text{CAB}]_t = \frac{[\text{X}] [\text{H}^+]}{K_1 K_2 [\text{S}]} + \frac{[\text{X}]}{K_2 [\text{S}]} + [\text{X}] \quad \dots(15)$$






$$\begin{aligned}
 [\text{CAB}]_t &= [\text{X}] \left\{ \frac{[\text{H}^+]}{K_1 K_2 [\text{S}]} + \frac{1}{K_2 [\text{S}]} + 1 \right\} \\
 &= [\text{X}] \left\{ \frac{[\text{H}^+] + K_1 + K_1 K_2 [\text{S}]}{K_1 K_2 [\text{S}]} \right\}
 \end{aligned}$$

$$\therefore [\text{X}] = \frac{K_1 K_2 [\text{S}] [\text{CAB}]_t}{[\text{H}^+] K_1 + K_1 K_2 [\text{S}]} \quad \dots(16)$$


$$\therefore \text{Rate} = \frac{K_1 K_2 k_3 [\text{S}] [\text{CAB}] [\text{Ru(III)}]}{[\text{H}^+] + K_1 + K_1 K_2 [\text{S}]} \quad \dots(17)$$

This rate law equation (17) is in good agreement with the experimental observations, including first order in  $[\text{CAB}]$ ,  $[\text{Ru(III)}]$  and inverse fractional order in  $[\text{H}^+]$  and fractional order on  $[\text{Niacinamide}]$ .

This is in good agreement with the experimental results. A detailed mechanism of  $\text{Ru(III)}$  catalyzed oxidation of niacinamide by CAB in HCl medium is given in Scheme 2.







Since rate =  $k'$  under pseudo first order condition of [Niacinamide]  $\gg$  [CAB], the rate equation (17) can be transformed into equation

$$k' = \frac{K_1 K_2 k_3 [S][\text{Ru(III)}]}{K_1 + [\text{H}^+] + K_1 K_2 [S]} \quad \dots(18)$$

$$\frac{1}{k'} = \frac{1}{K_2 k_3 [S][\text{Ru(III)}]} + \frac{[\text{H}^+]}{K_1 K_2 k_3 [S][\text{Ru(III)}]} + \frac{1}{k_3 [\text{Ru(III)}]} \quad \dots(19)$$

$$\frac{1}{k'} = \frac{1}{[S]K_2 k_3 [\text{Ru(III)}]} \left\{ \frac{K_1 + [\text{H}^+]}{K_1} \right\} + \frac{1}{k_3 [\text{Ru(III)}]} \quad \dots(20)$$

$$\frac{1}{k'} = \frac{[\text{H}^+]}{K_1 K_2 k_3 [S][\text{Ru(III)}]} + \left\{ \frac{1}{K_2 k_3 [S][\text{Ru(III)}]} + \frac{1}{k_3 [\text{Ru(III)}]} \right\} \dots(21)$$

Plots  $1/k'$  versus  $\frac{1}{[S]}$  at constant  $[\text{H}^+]$  and  $[\text{Ru(III)}]$  from equation (20) and  $1/k'$  versus  $[\text{H}^+]$  at constant  $[S]$  and  $[\text{Ru(III)}]$  from equation (21) were found to be linear (Figure 3). From the intercepts and slopes of the plots, the values of  $K_1$ ,  $K_2$  and  $k_3$  were evaluated. The protonation constant ( $K_P = 1/K_1$ ) value obtained, 18.20 for the species  $\text{PhSO}_2\text{NHBr}$  is in good agreement with the previously published work<sup>12,13</sup>. This gives indirect evidence for the proposed mechanism of the scheme 1. The thermodynamic parameters  $E_a$ ,  $\Delta H^\ddagger$ ,  $\Delta S^\ddagger$ , and  $\Delta G^\ddagger$  were calculated as shown in Table 3. The moderate value of enthalpy of activation is supportive for the proposed mechanism in scheme 1. The high negative value of entropy of activation ( $\Delta S^\ddagger$ ) indicates the formation of a rigid transition state by associative process




Fig.3. Plot of  $1/k'$  versus  $1/[S]$

## ACKNOWLEDGEMENT

We thank our Management and Principal of PES College of Engineering Mandya Karnataka, for encouragement and motivation.

## REFERENCES

1. N.M .Niven, Pharmacologic doses of Nicotinamide in the treatment of Inflammatory Skin conditions a review. *Cutis* 77 (1 suppl): PMID 168774. 2006; 11-6.
2. J.F. Tallman, S.M. Paul, P. Skolnick and D.W Gallager, 198. Receptors for the age of Anxiety Pharmacology of the Benzodiazepines Science, 207 (4428): 274-81.
3. Sharma Ashok, K. Mudgal Punit, and K.S. Gupta, 'Nicotinamide and Isonicotinamide was oxidized using permanganate ion in acidic medium' *Journal of the Indian Chemical Society*, 2008; 85; 920\_25.
4. Chandrashekar, Radhika, B.M. Venkatesha, and S. Ananda. Oxidation of Nicotinamide by bromamine-T in acidic medium catalysed by Ru(III) ion. *International Journal of Current Research*, 2014; 6(01): 4567\_71.
5. B.S Furniss, A.J. Hannatord, P.W.G Smith and A.R. Tatchell,. *Vogels Text book of organic chemistry*, Pearson Publication, 2009; 1213.
6. M.S. Ahmed, D.S. Mahadevappa, *Talanta*, 1980; 27: 669.
7. B. G. Pryde, and F.G. Soper *J. Chem. Soc*; 1926; 1582 and 1931; 1514.
8. E.Bishop, and Jennings, *Talanta* 1958; 1:197, 1961; 8: 22, 34, 697, 1962; 9: 581.
9. J.C. Morris, J.A. Salazar. and M.A. Wineman, *J. Am. Chem. Soc*, 1948; 70:2036
10. S.S. Narayana, and V.R.S Rao, *Radio chem acta*, 1983; 32: 211.
11. M. Subhashini, M. Subramanian, and V. R. S. Rao. *Talanta*, 1985; 32: 1082.
12. S. Ananda, M.B. Jagadeesha, and M.G. Ramananda, Kinetics of oxidation of cysteine by bromamine-T in acid Medium. *Asian journal of Chemistry*, 1998; 10 (4): 937\_46
13. B.M.Venkatesha, S. Ananda, and D.S. Mahadevappa. Kinetics of oxidation of chloroacetic acid by BAT in HCl medium and catalysed by Ru(III) ion. *Indian journal of Chemistry*, 1993; 32A :128\_35