

International Journal of Scientific Research and Reviews

Human Papillomavirus associated burden of Oral Squamous Cell Carcinoma in an Indian inhabitants

Turakhia Bhavika, Patel Amisha, and Shah Sejal *

Department of Microbiology, School of Science, RK University, Gujarat, India

ABSTRACT

Oral Squamous Cell Carcinoma (OSCC) is the eighth most common global malignancy and one of the leading cancer in Indian inhabitants. Gujarat is considered nowadays to be the high-risk region for oral cancer. The occurrence of OSCC mainly due to tobacco addiction; however 15-20% of overall oral cancer incidence in the non-smoker and non-alcoholic patients. Previous studies indicated that oral cancer might be caused by infection of Human Papillomavirus (HPV) along with consumption of tobacco. We have combined “PubMed” and “Medline” articles published from 1983 to 2017 and retrospective review has been done to identify the association of HPV and oral cancer. There are > 100 types of HPV have been found in oral lesions, including type 1, 2, 3, 4, 6, 7, 10, 11, 13, 16, 18, 30, 31, 32, 33, 35, 45, 52, 55, 57, 59, 69, 72, and 73. The prevalence of HPV in OSCC has been reported the overall prevalence ranges between 20-50%. The risk difference of OSCC with HPV infection varies ethnicity to ethnicity. The meta-analysis has shown HPV 16 and 18 strain found to be associated with the higher risk of developing OSCC in the Indian subcontinent. In view of the relationship between HPV and OSCC, further studies are required to reveal the role of HPV onset of oral neoplasm especially the oral cancer patients without the history of tobacco in any form.

KEYWORDS: HPV, OSCC, Oral cancer, India, Tobacco, PCR.

***Corresponding author**

Sejal Shah

Department of Microbiology,
School of Science,
RK University, Gujarat, India

INTRODUCTION:

Oral cancer is a perilous and emergent malignant neoplasm in many parts of the globe ¹, it represents the carcinoma of the tongue, lip, buccal mucosa, soft palate, hard palate, tonsil, alveolus and gingival^{1,2}. It has been reported that 90% of cases are Squamous Cell Carcinoma ^{3,4}. Oral cancer is the eight most common fatal disorder worldwide ⁵. In the Indian subcontinent, it is the leading malignancy among men and third most common type of cancer among women. Annually ~128,000 deaths occur from >260,000 new cases of Oral Squamous Cell Carcinoma (OSCC) ⁶.

Tobacco and alcohol consumption are the well-established risk factors for oral cancer ⁷. In the Asia-Pacific region, the concomitant and current form of tobacco is composed of areca nut, slaked lime, catechu, betel leaf and betel quid ⁸. Chewing betel quid is a common dietary habit widespread in many parts of the world^{1,3}. The frequent use of betel quid leads to a 50-fold increase in reactive oxygen species generated ^{2,9}. Another risk factor for developing OSCC is smoking cigars, pipes, and cigarettes ³. The western part of India, Gujarat is the tobacco growing belt, and it is obvious to be a high-risk region for OSCC due to the habit of chewing tobacco in various forms like pan, bidi and Gutkha. In our previous study we found 80% patients found to be associated with the history of tobacco in form of chewing and smoking, out of 71% patients had a history of tobacco chewing only, it clearly indicates the higher amount of tobacco consumption in form of chewing leads to OSCC⁸. However, approximately 20% of oral cancers developed in patients lacking these traditional risk factors ^{10,11}.




Figure 1: Various risk factors for developing OSCC.

There is increasing evidence that Human Papillomavirus (HPV) is an etiologic agent for developing oral cancer along with tobacco and alcohol addiction ¹¹. The co-existence of HPV in oropharyngeal and oral carcinoma was first studied ¹², and then supported the notion by many studies

^{13,14}. HPV is a small, non-enveloped, icosahedral, double-stranded DNA virus with the genome of 8 kb and it belongs to papillomaviridae family ¹⁵⁻¹⁸. HPV genome comprised of three distinct regions: The early region, the late region and URR (upstream regulatory region). The early part contains seven early proteins (E1, E2, E4, E5A, E5B, E6, and E7), the late includes two late proteins (L1, L2). The URR comprised promoter and enhancer DNA sequences that regulate viral replication ¹⁹. There are more than 150 different subtypes of HPV has been reported, and approximately 120 types are entirely sequenced ²⁰.




Figure 2: HPV structure and function of viral proteins¹¹¹

More than 10% of human cancers are developed by infection of HPV ²¹. On the bases of potential oncogenic HPVs can be categorized into two subsets: low-risk type and high-risk type ³. It causes both benign and malignant tumors of squamous epithelia ²². There are 24 types of HPV have been associated with benign lesions of OSCC including 1, 2, 3, 4, 6, 7, 10, 11, 13, 16, 18, 30, 31, 32, 33, 35, 45, 52, 55, 57, 59, 69, 72, and 73, and 12 types with malignant OSCC lesions including 2, 3, 6, 11, 13, 16, 18, 31, 33, 35, 52, and 57 ²³. The most predominant type detected in oral cancer is the HPV type 16^{24, 25}.

ROLE OF HPV IN OSCC PATHOGENESIS:

HPVs are mostly transmitted by close contact, especially oro-genital sex and high-risk sexual behavior through oral sexual activities such as the number of oral sex partners, and self-inoculation ^{26, 27}.

The buccal mucosa is the most exposed site to chemical carcinogens, infections, and trauma end up with carcinogenesis. The continuous exposure cause abrasion leads the mucosal surface more susceptible to HPV by penetration into the basal cells^{28, 29}.

The life cycle of HPV initiated through micro lesions in the epithelium and characterized by a special trophism keratinocytes, a specialized differentiation program of the infected squamous epithelial cell. Capsid proteins are linked to viral DNA synthesis and expression in keratinocytes. Usually, infection initiates in the basal and para-basal cells of the squamous epithelium. For a productive cell replication, a suitable micro-environment provided by the changes in the keratinocytes from the basal layer to the surface of the epithelium. Late and early promoter activation and synthesis of capsid occur when the basal layer is infected by virions released from the stratum corneum and granulosum³⁰.

The early and late proteins have a different function. E1 initiate viral DNA replication and transcription. E2 regulates DNA replication and viral transcription by controlling ORFs E6 to E7. E4 interacts with cytoskeleton proteins due to altering extracellular matrix cells. E5 interact with cellular proteins and down regulates MHC class I molecules; subtypes E5A and E5B, both have a role in membrane signaling. E6 degrades *p53*, which is tumor suppressor gene, and E7 binds to *pRb* oncoprotein. L1(major viral capsid proteins) and L2(the minor viral capsid proteins) in the late region, require for viral capsid formation in the final stage of replication³¹.




Figure-3: The proposed mechanism of high-risk HPV oncogenes in tumorigenesis

HPV integrates into the host genome at the E1/E2 sequence, hence up regulates E6 and E7 (encoded in an early open reading frame) and increase its tumor igenicity. E2 protein negatively regulates E6 and E7 protein expression encoded by the ORF of the virus. *p53* and *pRb* are the tumour suppressor genes that disrupted by E6 and E7 proteins by altering its host genome leads to cell proliferation and tumour formation.¹¹²

MATERIALS AND METHODS:

We collected the information using “PubMed” and “Medline” database from 1983 to 2017 and the retrospective review has been done to check the association between HPV and oral cancers and its occurrence at multiple sub-sites in the oral cavity like tongue, buccal mucosa, soft palate, hard palate, lips and gingiva. The keywords searched for the data collection are “HPV”, “OSCC”, “oral cancer”, “HNSCC”, “Indian population”, “Human Papillomavirus”, “cervical cancer” and “prognosis”, etc. Meta-analysis of HPV screening was done by type-specific PCR-based assays or in situ hybridization (ISH) ^{15, 32-35}.

AGE AND GENDER:

Oral cancer is a lethal disorder that occurs mainly in the elder age. However, most of the cases of oral cancer prevalence between 50 and 70 years of age ³⁶, There are certain cases where, occurrence of OSCC identified as early as ten years of age without any type of history³⁷. In most of the countries, the mean age of OSCC prevalence is usually between 51-55 years of age (Table-1). Around 17% of the patients are younger below 40 years of age ³⁸. The younger patients are affected three times more with the high-risk HPV types than elders, due to viral transmission by direct physical contact ³⁹. Considering all the age groups, men are more affected than women for oral cancer ⁴⁰.

SITE:

There has been an extensive variation of HPV positive rates in cancers at different sites of the oral region. Approximately 40% of gingivobuccal cancers have tested HPV positive ⁴¹. Previously 43% of tonsillar cancers were found positive for oncogenic subtypes of HPV ⁴², In contrast, the other study has reported 66% of tonsillar cancer, followed by the base of tongue 34% ⁴³. The earlier research has shown the prevalence of HPV ranged from 70% in the case of buccal mucosa, 67% in the case of the floor of the mouth and 82% in the case of tongue ⁴⁴. The previous study has reported 61% head and neck cancer samples were positive for HPV ⁴⁵. The number of studies has indicated HPV 72.5-91% prevalence of HPV in tongue cancers ⁴⁶. According to these results, oncogenicity of HPV should be considered as a high risk for tongue cancer. The tongue is the foremost among all sites contributing up to 42% of all oral cavity sites in Japan, Taiwan, Thailand, Yemen, India and Iran ⁴⁰. The Indian scenario is showing high occurrence of buccal mucosa SCC, which is due to tobacco chewing at specific buccal site⁸.

PREVALENCE OF HPV IN ORAL CANCERS IN INDIA:

Overall 23.5% of prevalence of HPV was reported from the geographic region⁴⁷. In an Indian scenario, petite data is available for HPV induced oral carcinomas. In Western countries, the floor of mouth and tongue are the generic sites for developing squamous cell carcinomas (SCC). The most exposed site observed was buccal mucosa due to tobacco chewing site for OSCC in Indian subset.⁴⁸ The earlier research demonstrated 33.6% prevalence of HPV in OSCC patients in Eastern India⁴⁹, in contrast, the studying the same region reported 69% HPV prevalence⁵⁰. As etiology differs, the study in Western India showed 15% HPV prevalence¹⁴. The incidence of HPV in South India was reported higher as compared to Eastern and Western India. The earlier study identified 16.8% of HPV prevalence in South Indian region⁵¹ whereas, the same region showed 31% and 74% prevalence of HPV^{52, 45}. The study performed in western India has shown 0% HPV prevalence in OSCC while 59.6% HPV prevalence found in cervical cancer¹⁰. A petite study has been performed in the Western region of the India regarding OSCC, as it is the leading neoplasm, the further study need to be validated for the HPV detection.

Table 1: Role of HPV and its association with OSCC- Global incidence rate

References	Year	Country	OSCC Cases	Mean age	Tobacco	Smoking	Alcohol	HPV (+ve)	HPV genotype	
									HPV 16	HPV 18
15	2016	India	47	-	13	10	5	19	-	-
44	2015	Netherlands	176	63	-	33	-	47	42	1
66	2014	China	312	-	265	272	227	52	13	6
67	2014	Germany	88	-	-	54	64	5	3	-
68	2014	Germany	93	57	-	73	72	25	-	-
69	2014	China	200	81	100	-	71	55	39	15
70	2013	China	400	52	225	222	195	87	20	18
10	2013	India	97	46.7	86	-	-	0	0	0
71	2013	Colombia	67	65.9	-	-	-	33	27	6
72	2013	Mexico	80	63	38	-	33	4	2	2
73	2013	USA	409	-	-	-	-	24	15	2
74	2012	China	103	49	79	90	72	31	16	7
75	2012	Belgium	162	57	-	52	44	65	-	-
76	2012	China	173	51	-	-	-	28	16	12
77	2012	China	65	53	-	-	-	24	-	-
78	2012	Brazil	114	-	36	-	28	22	15	-
3	2012	Malaysia	30	30	6	6	-	1	-	1
79	2012	Hradec-Kralove	24	63	-	9	-	5	4	-
80	2011	Carolina	25	30	-	-	-	11	2	-
81	2011	India	60	55	-	-	-	30	29	-
32	2011	India	34	-	-	-	-	24	10	13
82	2011	France	209	59	-	-	-	22		
83	2009	China	52	-	-	31	21	21	33	16
84	2008	Brazil	29	23	26	-	18	5	-	-
85	2007	Japan	66	-	-	23	24	66	24	-
86	2007	Japan	20	-	-	-	-	20	8	13
12	2006	Italy	63	68.89	-	11	6	24	8	10

87	2006	South- Africa	59		-	-	-	7	-	7
88	2006	USA	94	56.6	-	-	-	12	10	-
89	2005	Belgrade	50	32	-	30	27	32	10	6
90	2004	Japan	73	-	42	-	14	54	43	18
91	2004	Sweden	110	-	-	-	-	12	9	-
92	2004	Venezuela	16	-	-	-	-	8	-	-
52	2003	India	42	-	-	-	-	13	9	-
93	2003	USA	141	-	114	-	106	29	24	1
94	2003	Japan	86	-	-	-	-	30	28	2
94	2003	Brazil	46	-	-	-	-	18	-	-
96	2003	Japan	103	51.1	-	-	-	51	-	-
49	2002	India	110	-	-	-	-	37	25	16
97	2002	Taiwan	29	-	19	18	-	24	24	20
34	2001	USA	254	54.2	-	-	-	62	62	-
33	2001	Venezuela	50	-	-	26	19	30	-	-
98	2000	Japan	83	-	-	-	-	47	-	-
17	2000	Greece	53	52.7	-	-	-	48	24	11
51	1999	India	61	-	-	-	-	17	-	-
99	1999	Greece	81	-	-	-	-	40	5	22
29	1998	USA	284	-	-	-	-	64	41	2
100	1998	Venezuela	50	55	-	-	-	35	13	-
101	1998	Korea	42	-	-	-	-	22	15	6
102	1998	Sudan	88	-	-	-	-	0	0	0
103	1996	USA	64	-	-	-	-	20	11	2
104	1996	Netherland	35	-	-	-	-	19	15	-
105	1995	South- Africa	66	-	-	-	-	1	0	1
106	1995	Japan	77	-	-	-	-	24	23	1
107	1993	USA	39	-	-	-	-	7	7	1
108	1991	Spain	15	-	-	-	-	0	0	0
109	1990	USA	100	-	10	-	-	6	-	-
110	1990	Finland	40	-	-	-	-	16	10	1
35	1988	Finland	51	-	-	-	-	6	5	-

[OSCC= Oral Squamous Cell Carcinoma, HPV= Human Papillomavirus,- = Unknown]

OTHER CANCERS:

HPV infections have been found in many other cancers. Globally, 99.7% prevalence of HPV has been reported in cervical cancer⁵³. The rate of cervical cancer is maximum in South-Eastern Asia with an especially large burden in India, Latin America and sub-Saharan Africa⁵⁴. The previous study showed a prevalence of HPV in 71% of invasive anal cancers⁵⁵. Serological studies have confirmed HPV-16 and HPV-18 are the etiologic agent, and HPV-DNA is detectable in about 40–50% of all penile cancers. Overall, 17.5% HPV prevalence reported in patients with ovarian cancer⁵⁶. Approximately 13,000 cases of penile cancer, 35,000 of anal cancer, 8,500 of vulvar cancer and 12,000 of vaginal cancer are attributable to HPV worldwide. Globally, ~ 38,000 cases of head and neck cancer are attributable to HPV, and HPV 16/18 are responsible for 85% of the head and neck carcinoma⁵⁷. The prevalence of HR-HPV in the esophageal squamous cell carcinoma was reported 24.8% in Chinese population⁵⁸. The overall prevalence of HPV was reported 28.0% in laryngeal cancer⁵⁹. A study screened for HPV DNA and found 65-85 %of HPV infection in breast cancers positive⁶⁰. An association between HPV and SCC of the urinary bladder ranging from 0% to 17% in present outcomes⁶¹. The prevalence of HPV has been reported 37.57% among lung cancer patients in China⁶².

FUTURE DIRECTION:

Global survey of HPV infection and oral cancers has been reported. According to an Indian data, 20-50% of the HPV prevalence detected in OSCC in the different region of the India¹¹². The variation in the data showed, there is a strong recommendation of the new hypothesis to screen the HPV, especially in the western part of the India, as it has found higher number of OSCC patients compared to other region of the India. The number of studies has reported that overall survival significantly higher among HPV-positive patients as compared to patients with HPV-negative tumour⁶³⁻⁶⁵. If the percentage of the HPV affected patients have known, the prognosis of the patients could be better.

The further study needs more polishing touch, which includes 1) The investigation of the epidemiological profile of oral HPV infection and occurrence of OSCCs, 2) HPV transmission into oral cavity and how it causes oral cancers, 3) To investigate whether the HPV vaccines are biologically effective for preventing infection of oral HPV and related disease, 4) Complete examination is needed on the effects of HPV vaccines for preventing HPV positive oral carcinoma, and 5) analyze the status of HPV whether it should be used in clinical examinations for anti-cancer

treatment through forthcoming studies. Understanding the etiological importance of HPV in oral carcinoma might be required in prognosis, treatment and prevention of disease.

CONCLUSION:

In the Indian subcontinent, oral cancer is a major personal tragedy and public health problem. The meta-analysis has shown HPV 16 and 18 strain found to be associated with the higher risk of developing OSCC. In view of the relationship between HPV and OSCC, it is must to design additional studies to validate its role in development of oral carcinomas especially the oral cancer patients without the history of tobacco in any form. The future studies should be carried out to screen HPV infection in oral cancer patients in Gujarat region especially as it is the blazing dilemma in the current area.

ACKNOWLEDGEMENT

This work has been presented into NCRIS 2018. I sincerely acknowledge RK University to provide me such a platform for the poster presentation.

REFERENCES

1. Warnakulasuriya, S., Causes of oral cancer—an appraisal of controversies. *British dental journal*, 2009; 207(10): 471-475.
2. Anantharaman, D., Chaubal, P.M., Kannan, S., Bhisey, R.A. and Mahimkar, M.B., Susceptibility to oral cancer by genetic polymorphisms at CYP1A1, GSTM1 and GSTT1 loci among Indians: tobacco exposure as a risk modulator. *Carcinogenesis*, 2007; 28(7): 1455-1462.
3. Goot-Heah, K., Kwai-Lin, T., Froemming, G.R.A., Abraham, M.T., Rosdy, N.M.M.N.M. and Zain, R.B., Human papilloma virus 18 detection in oral squamous cell carcinoma and potentially malignant lesions using saliva samples. *Asian Pacific Journal of Cancer Prevention*, 2012; 13(12): 6109-6113.
4. Cooper, J.S., Porter, K., Mallin, K., Hoffman, H.T., Weber, R.S., Ang, K.K., Gay, E.G. and Langer, C.J., National Cancer Database report on cancer of the head and neck: 10-Year update. *Head & neck*, 2009; 31(6): 748-758.
5. Maitra, A., Biswas, N.K., Amin, K., Kowtal, P., Kumar, S., Das, S., Sarin, R., Majumder, P.P., Bagchi, I., Bairagya, B.B. and Basu, A., Mutational landscape of gingivo-buccal oral

- squamous cell carcinoma reveals new recurrently-mutated genes and molecular subgroups. *Nature communications*, 2013; 4: 2873.
6. Shah, S., Shah, S., Padh, H. and Kalia, K., Genetic alterations of the PIK3CA oncogene in human oral squamous cell carcinoma in an Indian population. *Oral surgery, oral medicine, oral pathology and oral radiology*, 2015; 120(5): 628-635.
 7. Mork, J., Lie, A.K., Glatte, E., Clark, S., Hallmans, G., Jellum, E., Koskela, P., Møller, B., Pukkala, E., Schiller, J.T. and Wang, Z., Human papillomavirus infection as a risk factor for squamous-cell carcinoma of the head and neck. *New England Journal of Medicine*, 2001; 344(15): 1125-1131.
 8. Shah, S., Jajal, D., Mishra, G. and Kalia, K., Genetic profile of PTEN gene in Indian oral squamous cell carcinoma primary tumors. *Journal of Oral Pathology & Medicine*, 2017; 46(2): 106-111.
 9. Nair, U.J., Nair, J., Mathew, B. and Bartsch, H., Glutathione S-transferase M1 and T1 null genotypes as risk factors for oral leukoplakia in ethnic Indian betel quid/tobacco chewers. *Carcinogenesis*, 1999; 20(5): 743-748.
 10. Patel, K.R., Vajaria, B.N., Begum, R., Desai, A., Patel, J.B., Shah, F.D., Shukla, S.N. and Patel, P.S., Prevalence of high-risk human papillomavirus type 16 and 18 in oral and cervical cancers in population from Gujarat, West India. *Journal of oral pathology & medicine*, 2014; 43(4): 293-297.
 11. Campisi, G., Giovannelli, L., Calvino, F., Matranga, D., Colella, G., Di Liberto, C., Capra, G., Leao, J.C., Muzio, L.L., Capogreco, M. and D'Angelo, M., HPV infection in relation to OSCC histological grading and TNM stage. Evaluation by traditional statistics and fuzzy logic model. *Oral oncology*, 2006; 42(6): 638-645.
 12. Syrjänen, K., Syrjänen, S., Lamberg, M., Pyrhönen, S. and Nuutinen, J., Morphological and immunohistochemical evidence suggesting human papillomavirus (HPV) involvement in oral squamous cell carcinogenesis. *International journal of oral surgery*, 1983; 12(6): 418-424.
 13. De Villiers, E.M., Weidauer, H., Otto, H. and ZurHausen, H., Papillomavirus DNA in human tongue carcinomas. *International journal of cancer*, 1985; 36(5): 575-578.
 14. D'Costa, J., Saranath, D., Dedhia, P., Sanghvi, V. and Mehta, A.R., Detection of HPV-16 genome in human oral cancers and potentially malignant lesions from India. *Oral oncology*, 1998; 34(5): 413-420.
 15. Bijina, B.R., Ahmed, J., Shenoy, N., Ongole, R., Shenoy, S. and Baliga, S., Detection of human papilloma virus in potentially malignant and malignant lesions of the oral cavity and a study of associated risk factors. *South Asian journal of cancer*, 2016; 5(4): 179.

16. Hagensee, M.E., Yaegashi, N. and Galloway, D.A., Self-assembly of human papillomavirus type 1 capsids by expression of the L1 protein alone or by coexpression of the L1 and L2 capsid proteins. *Journal of virology*, 1993;67(1): .315-322.
17. Bouda, M., Gorgoulis, V.G., Kastrinakis, N.G., Giannoudis, A., Tsoli, E., Danassi-Afentaki, D., Foukas, P., Kyroudi, A., Laskaris, G., Herrington, C.S. and Kittas, C., “High risk” HPV types are frequently detected in potentially malignant and malignant oral lesions, but not in normal oral mucosa. *Modern Pathology*, 2000; 13(6): 644-653.
18. Datta, S., Agarwal, M., Chatterjee, S., Gogoi, H.K., Veer, V. and Singh, L., Detection of human papillomavirus in women attending Pap cervical screening camp at a peripheral hospital of North-Eastern India. *Medical journal armed forces india*, 2015; 71(2): 182-185.
19. Kim, S.M., Human papilloma virus in oral cancer. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*, 2016; 42(6): 327-336.
20. Rautava, J. and Syrjänen, S.,Biology of human papillomavirus infections in head and neck carcinogenesis. *Head and neck pathology*, 2012; 6(1): 3-15.
21. Jain, N., Singh, V., Hedau, S., Kumar, S., Daga, M.K., Dewan, R., Murthy, N.S., Husain, S.A. and Das, B.C., Infection of human papillomavirus type 18 and p53 codon 72 polymorphism in lung cancer patients from India. *CHEST Journal*, 2005; 128(6): 3999-4007.
22. Duensing, S. and Münger, K., Mechanisms of genomic instability in human cancer: insights from studies with human papillomavirus oncoproteins. *International Journal of Cancer*, 2004; 109(2): 157-162.
23. Kojima, A., Maeda, H., Sugita, Y., Tanaka, S. and Kameyama, Y., Human papillomavirus type 38 infection in oral squamous cell carcinomas. *Oral Oncology*, 2002; 38(6): 591-596.
24. Smith, E.M., Ritchie, J.M., Summersgill, K.F., Klussmann, J.P., Lee, J.H., Wang, D., Haugen, T.H. and Turek, L.P., Age, sexual behavior and human papillomavirus infection in oral cavity and oropharyngeal cancers. *International journal of cancer*, 2004; 108(5): 766-772.
25. Herrero, R., Castellsagué, X., Pawlita, M., Lissowska, J., Kee, F., Balaram, P., Rajkumar, T., Sridhar, H., Rose, B., Pintos, J. and Fernández, L., Human papillomavirus and oral cancer: the International Agency for Research on Cancer multicenter study. *Journal of the National Cancer Institute*, 2003; 95(23): 1772-1783.
26. Edelstein, Z.R., Schwartz, S.M., Hawes, S., Hughes, J.P., Feng, Q., Stern, M.E., O’Reilly, S., Lee, S.K., Xi, L.F. and Koutsky, L.A., Rates and determinants of oral human papillomavirus (HPV) infection in young men. *Sexually transmitted diseases*, 2012; 39(11): 860.

27. Cook, R.L., Thompson, E.L., Kelso, N.E., Friary, J., Hosford, J., Barkley, P., Dodd, V.J., Abrahamsen, M., Ajinkya, S., Obesso, P.D. and Rashid, M.H., Sexual behaviors and other risk factors for oral human papillomavirus infections in young women. *Sexually transmitted diseases*, 2014;41(8): 486.
28. Jeon, S. and Lambert, P.F., Integration of human papillomavirus type 16 DNA into the human genome leads to increased stability of E6 and E7 mRNAs: implications for cervical carcinogenesis. *Proceedings of the National Academy of Sciences*, 1995; 92(5): 1654-1658.
29. Syrjänen, S.M., Syrjänen, K.J. and Happonen, R.P., Human papillomavirus (HPV) DNA sequences in oral precancerous lesions and squamous cell carcinoma demonstrated by in situ hybridization. *Journal of Oral Pathology & Medicine*, 1988; 17(6): 273-278.
30. Campisi, G., Panzarella, V., Giuliani, M., Lajolo, C., Di Fede, O., Falaschini, S., Di Liberto, C., Scully, C. and Lo Muzio, L., Human papillomavirus: Its identikit and controversial role in oral oncogenesis, premalignant and malignant lesions. *International journal of oncology*, 2007;30(4): 813-823.
31. Kim, S.M., Human papilloma virus in oral cancer. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*, 2016; 42(6): 327-336.
32. Kulkarni, S.S., Kulkarni, S.S., Vastrad, P.P., Kulkarni, B.B., Markande, A.R., Kadakol, G.S., Hiremath, S.V., Kaliwal, S., Patil, B.R. and Gai, P.B., Prevalence and distribution of high risk human papillomavirus (HPV) types 16 and 18 in carcinoma of cervix, saliva of patients with oral squamous cell carcinoma and in the general population in Karnataka, India. *Asian Pac J Cancer Prev*, 2011; 12(3): 645-648.
33. Premoli-De-Percoco, G. and Ramirez, J.L., High risk human papillomavirus in oral squamous carcinoma: evidence of risk factors in a Venezuelan rural population. Preliminary report. *Journal of oral pathology & medicine*, 2001; 30(6): 355-361.
34. Schwartz, S.R., Yueh, B., McDougall, J.K., Daling, J.R. and Schwartz, S.M., Human papillomavirus infection and survival in oral squamous cell cancer: a population-based study. *Otolaryngology—Head and Neck Surgery*, 2001; 125(1): 1-9.
35. Syrjänen, S.M., Syrjänen, K.J. and Happonen, R.P., Human papillomavirus (HPV) DNA sequences in oral precancerous lesions and squamous cell carcinoma demonstrated by in situ hybridization. *Journal of Oral Pathology & Medicine*, 1988; 17(6): 273-278.
36. Bhurgri, Y., Bhurgri, A., Usman, A., Pervez, S., Kayani, N., Bashir, I., Ahmed, R. and Hasan, S.H., Epidemiological review of head and neck cancers in Karachi. *Asian Pacific Journal of Cancer Prevention*, 2006; 7(2): 195.

37. Solanki, M.C., Gandhi, S., Koshy, G. and Mathew, G.C., Squamous cell carcinoma of maxilla in 10 year old boy: A rare case report. *International Journal of Pediatric Otorhinolaryngology Extra*, 2012;7(1): 33-35.
38. Halboub, E.S., Al-Anazi, Y.M. and Al-Mohaya, M.A., Characterization of Yemeni patients treated for oral and pharyngeal cancers in Saudi Arabia. *Saudi Med J*, 2011; 32(11): 1177-1182.
39. Schwartz, S.M., Daling, J.R., Madeleine, M.M., Doody, D.R., Fitzgibbons, E.D., Wipf, G.C., Carter, J.J., Mao, E.J., Huang, S., Beckmann, A.M. and McDougall, J.K., Oral cancer risk in relation to sexual history and evidence of human papillomavirus infection. *Journal of the National Cancer Institute*, 1998; 90(21): 1626-1636.
40. Rao, S.V.K., Mejia, G., Roberts-Thomson, K. and Logan, R., Epidemiology of oral cancer in Asia in the past decade-an update (2000-2012). *Asian Pacific journal of cancer prevention*, 2013; 14(10): .5567-5577.
41. Zafereo, M.E., Xu, L., Dahlstrom, K.R., Viamonte, C.A., El-Naggar, A.K., Wei, Q., Li, G. and Sturgis, E.M., Squamous cell carcinoma of the oral cavity often overexpresses p16 but is rarely driven by human papillomavirus. *Oral oncology*, 2016; 56: 47-53.
42. Mellin, H., Friesland, S., Lewensohn, R., Dalianis, T. and Munck-Wikland, E., Human papillomavirus (HPV) DNA in tonsillar cancer: clinical correlates, risk of relapse, and survival. *International journal of cancer*, 2000; 89(3): 300-304.
43. Koskinen, W.J., Chen, R.W., Leivo, I., Mäkitie, A., Bäck, L., Kontio, R., Suuronen, R., Lindqvist, C., Auvinen, E., Molijn, A. and Quint, W.G., Prevalence and physical status of human papillomavirus in squamous cell carcinomas of the head and neck. *International Journal of Cancer*, 2003; 107(3): 401-406.
44. Melchers, L.J., Mastik, M.F., Cameron, B.S., Van Dijk, B.A.C., De Bock, G.H., Van Der Laan, B.F.A.M., Van Der Vegt, B., Speel, E.J.M., Roodenburg, J.L.N., Witjes, M.J.H. and Schuurin, E., Detection of HPV-associated oropharyngeal tumours in a 16-year cohort: more than meets the eye. *British journal of cancer*, 2015;112(8): 1349-1357.
45. Balaram, P., Nalinakumar, K.R., Abraham, E., Balan, A., Hareendran, N.K., Bernard, H.U. and Chan, S.Y., Human papillomaviruses in 91 oral cancers from Indian betel quid chewers—high prevalence and multiplicity of infections. *International journal of cancer*, 1995; 61(4): 450-454.
46. Miller, C.S. and White, D.K., Human papillomavirus expression in oral mucosa, premalignant conditions, and squamous cell carcinoma: a retrospective review of the

- literature. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology, 1996; 82(1): 57-68.
47. Kreimer, A.R., Clifford, G.M., Boyle, P. and Franceschi, S., Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. *Cancer Epidemiology and Prevention Biomarkers*, 2005; 14(2): 467-475.
48. Sankaranarayanan, R., Duffy, S.W., Padmakumary, G., Day, N.E. and Nair, M.K., Risk factors for cancer of the buccal and labial mucosa in Kerala, southern India. *Journal of Epidemiology & Community Health*, 1990; 44(4): 286-292.
49. Nagpal, J.K., Patnaik, S. and Das, B.R., Prevalence of high-risk human papilloma virus types and its association with P53 codon 72 polymorphism in tobacco addicted oral squamous cell carcinoma (OSCC) patients of Eastern India. *International journal of cancer*, 2002; 97(5): 649-653.
50. Chatterjee, R., Mukhopadhyay, D., Chakraborty, R.N. and Mitra, R.B., Evaluation of argyrophilic nucleolar organizer regions (AgNORs) in oral carcinomas in relation to human papillomavirus infection and cytogenetics. *Journal of oral pathology & medicine*, 1997; 26(7): 310-314.
51. Pillai, M.R., Phanidhara, A., Kesari, A.L., Nair, P. and Nair, M.K., Cellular manifestations of human papillomavirus infection in the oral mucosa. *Journal of surgical oncology*, 1999; 71(1): 10-15.
52. Kumar, R.V., Kadkol, S.S., Daniel, R., Shenoy, A.M. and Shah, K.V., Human papillomavirus, p53 and cyclin D1 expression in oropharyngeal carcinoma. *International journal of oral and maxillofacial surgery*, 2003; 32(5): 539-543.
53. Walboomers, J.M., Jacobs, M.V., Manos, M.M., Bosch, F.X., Kummer, J.A., Shah, K.V., Snijders, P.J., Peto, J., Meijer, C.J. and Munoz, N., Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *The Journal of pathology*, 1999; 189(1): 12-19.
54. De Martel, C., Plummer, M., Vignat, J. and Franceschi, S., Worldwide burden of cancer attributable to HPV by site, country and HPV type. *International journal of cancer*, 2017; 141: 664-670.
55. Hoots, B.E., Palefsky, J.M., Pimenta, J.M. and Smith, J.S., Human papillomavirus type distribution in anal cancer and anal intraepithelial lesions. *International journal of cancer*, 2009; 124(10): 2375-2383.
56. Shukla, S., Bharti, A.C., Mahata, S., Hussain, S., Kumar, R., Hedau, S. and Das, B.C., 2009. Infection of human papillomaviruses in cancers of different human organ sites.

57. Rosa, M.I., Silva, G.D., de AzedoSimões, P.W.T., Souza, M.V., Panatto, A.P.R., Simon, C.S., Madeira, K. and Medeiros, L.R., The prevalence of human papillomavirus in ovarian cancer: a systematic review. *International Journal of Gynecological Cancer*, 2013; 23(3): 437-441.
58. Ahmed, H.G., Beely, M.A.B.I., Aziz, M.S.A.E., Alshammari, F.D. and Elzaki, S.E.G., Molecular Detection of High Risk Human Papilloma Viruses (HR-HPV 16 and 18) among Patients with Esophageal Cancer. *J Hum VirolRetrovirol*, 2017; 5(3): 00154.
59. Li, X., Gao, L., Li, H., Gao, J., Yang, Y., Zhou, F., Gao, C., Li, M. and Jin, Q., Human papillomavirus infection and laryngeal cancer risk: a systematic review and meta-analysis. *The Journal of infectious diseases*, 2012; 207(3): 479-488.
60. De Villiers, E.M., Sandstrom, R.E., zurHausen, H. and Buck, C.E., Presence of papillomavirus sequences in condylomatous lesions of the mamillae and in invasive carcinoma of the breast. *Breast Cancer Research*, 2004; 7(1): R1.
61. Jørgensen, K.R., Høyer, S., Sørensen, M.M. and Jensen, J.B. Human papillomavirus types 44, 52, 66 and 67 detected in a woman with squamous cell carcinoma of the urinary bladder. *Scandinavian journal of urology*, 2017;51(1): 85-86
62. Zhai, K., Ding, J. and Shi, H.Z. HPV and lung cancer risk: a meta-analysis. *Journal of Clinical Virology*, 2015; 63: 84-90.
63. Ang, K.K., Harris, J., Wheeler, R., Weber, R., Rosenthal, D.I., Nguyen-Tân, P.F., Westra, W.H., Chung, C.H., Jordan, R.C., Lu, C. and Kim, H. Human papillomavirus and survival of patients with oropharyngeal cancer. *New England Journal of Medicine*, 2010; 363(1): .24-35.
64. Chiba, I., Shindoh, M., Yasuda, M., Yamazaki, Y., Amemiya, A., Sato, Y., Fujinaga, K., Notani, K.I. and Fukuda, H. Mutations in the p53 gene and human papillomavirus infection as significant prognostic factors in squamous cell carcinomas of the oral cavity. *Oncogene*, 1996; 12(8): 1663-1668.
65. Ritchie, J.M., Smith, E.M., Summersgill, K.F., Hoffman, H.T., Wang, D., Klussmann, J.P., Turek, L.P. and Haugen, T.H. Human papillomavirus infection as a prognostic factor in carcinomas of the oral cavity and oropharynx. *International journal of cancer*, 2003;104(3): 336-344.
66. Huang, C.G., Lee, L.A., Tsao, K.C., Liao, C.T., Yang, L.Y., Kang, C.J., Chang, K.P., Huang, S.F., Chen, I.H., Yang, S.L. and Lee, L.Y. Human papillomavirus 16/18 E7 viral loads predict distant metastasis in oral cavity squamous cell carcinoma. *Journal of Clinical Virology*, 2014; 61(2): 230-236.

67. Krüger, M., Pabst, A.M., Walter, C., Sagheb, K., Günther, C., Blatt, S., Weise, K., Al-Nawas, B. and Ziebart, T. The prevalence of human papilloma virus (HPV) infections in oral squamous cell carcinomas: a retrospective analysis of 88 patients and literature overview. *Journal of Cranio-Maxillo-Facial Surgery*, 2014; 42(7): 1506-1514.
68. Meyer, M.F., Seuthe, I.M., Drebber, U., Siefer, O., Kreppel, M., Klein, M.O., Mikolajczak, S., Klussmann, J.P., Preuss, S.F. and Huebbers, C.U. Valosin-containing protein (VCP/p97)-expression correlates with prognosis of HPV-negative oropharyngeal squamous cell carcinoma (OSCC). *PloS one*, 2014; 9(12): 114-170.
69. Gan, L.L., Zhang, H., Guo, J.H. and Fan, M.W. Prevalence of human papillomavirus infection in oral squamous cell carcinoma: a case-control study in Wuhan, China. *Asian Pac J Cancer Prev*, 2014;15(14): 5861-5865.
70. Lee, L.A., Huang, C.G., Tsao, K.C., Liao, C.T., Kang, C.J., Chang, K.P., Huang, S.F., Chen, I.H., Fang, T.J., Li, H.Y. and Yang, S.L. Increasing rates of low-risk human papillomavirus infections in patients with oral cavity squamous cell carcinoma: association with clinical outcomes. *Journal of clinical virology*, 2013; 57(4): 331-337.
71. Quintero, K., Giraldo, G.A., Uribe, M.L., Baena, A., Lopez, C., Alvarez, E. and Sanchez, G.I., Human papillomavirus types in cases of squamous cell carcinoma of head and neck in Colombia. *Brazilian journal of otorhinolaryngology*, 2013;79(3): 375-381.
72. González-Ramírez, I., Irigoyen-Camacho, M.E., Ramírez-Amador, V., Lizano-Soberón, M., Carrillo-García, A., García-Carrancá, A., Sánchez-Pérez, Y., Méndez-Martínez, R., Granados-García, M., Ruíz-Godoy, L.M. and García-Cuellar, C.M. Association between age and high-risk human papilloma virus in Mexican oral cancer patients. *Oral diseases*, 2013; 19(8): 796-804.
73. Lingen, M.W., Xiao, W., Schmitt, A., Jiang, B., Pickard, R., Kreinbrink, P., Perez-Ordóñez, B., Jordan, R.C. and Gillison, M.L. Low etiologic fraction for high-risk human papillomavirus in oral cavity squamous cell carcinomas. *Oral oncology*, 2013; 49(1): 1-8.
74. Huang, S.F., Li, H.F., Liao, C.T., Wang, H.M., Chen, I.H., Chang, J.C., Chen, Y.J. and Cheng, A.J. Association of HPV infections with second primary tumors in early-staged oral cavity cancer. *Oral diseases*, 2012; 18(8): 809-815.
75. Duray, A., Descamps, G., Decaestecker, C., Rimmelink, M., Sirtaine, N., Lechien, J., Ernoux-Neufcoeur, P., Bletard, N., Somja, J., Depuydt, C.E. and Delvenne. Human papillomavirus DNA strongly correlates with a poorer prognosis in oral cavity carcinoma. *The Laryngoscope*, 2012;122(7): 1558-1565.

76. Lee, L.A., Huang, C.G., Liao, C.T., Lee, L.Y., Hsueh, C., Chen, T.C., Lin, C.Y., Fan, K.H., Wang, H.M., Huang, S.F. and Chen, I.H. Human papillomavirus-16 infection in advanced oral cavity cancer patients is related to an increased risk of distant metastases and poor survival. *PloS one*, 2012; 7(7): 407-467.
77. Chen, S.F., Yu, F.S., Chang, Y.C., Fu, E., Nieh, S. and Lin, Y.S. Role of human papillomavirus infection in carcinogenesis of oral squamous cell carcinoma with evidences of prognostic association. *Journal of Oral Pathology & Medicine*, 2012; 41(1): 9-15.
78. Kaminagakura, E., Villa, L.L., Andreoli, M.A., Sobrinho, J.S., Vartanian, J.G., Soares, F.A., Nishimoto, I.N., Rocha, R. and Kowalski, L.P. High-risk human papillomavirus in oral squamous cell carcinoma of young patients. *International journal of cancer*, 2012; 130(8): 1726-1732.
79. Laco, J., Nekvindova, J., Novakova, V., Celakovsky, P., Dolezalova, H., Tucek, L., Vosmikova, H., Vosmik, M., Neskudlova, T., Cermakova, E. and Hacova, M. Biologic importance and prognostic significance of selected clinicopathological parameters in patients with oral and oropharyngeal squamous cell carcinoma, with emphasis on smoking, protein p16INK4a expression, and HPV status. *Neoplasma*, 2012; 59(4): 398.
80. Harris, S.L., Thorne, L.B., Seaman, W.T., Neil Hayes, D., Couch, M.E. and Kimple, R.J. Association of p16INK4a overexpression with improved outcomes in young patients with squamous cell cancers of the oral tongue. *Head & neck*, 2011;33(11): 1622-1627.
81. Elango, K.J., Suresh, A., Erode, E.M., Subhadradevi, L., Ravindran, H.K., Iyer, S.K., Iyer, S.K. and Kuriakose, M.A. Role of human papilloma virus in oral tongue squamous cell carcinoma. *Asian Pac J Cancer Prev*, 2011; 12(4): 889-896.
82. St Guily, J.L., Jacquard, A.C., Prétet, J.L., Haesebaert, J., Beby-Defaux, A., Clavel, C., Agius, G., Birembaut, P., Okaïs, C., Léocmach, Y. and Soubeyrand, B. Human papillomavirus genotype distribution in oropharynx and oral cavity cancer in France—the EDiTH VI study. *Journal of Clinical Virology*, 2011; 51(2): 100-104.
83. Zhao, D., Xu, Q.G., Chen, X.M. and Fan, M.W. Human papillomavirus as an independent predictor in oral squamous cell cancer. *International journal of oral science*, 2009; 1(3): 119.
84. Simonato, L.E., Garcia, J.F., Sundefeld, M.L.M.M., Mattar, N.J., Veronese, L.A. and Miyahara, G.I. Detection of HPV in mouth floor squamous cell carcinoma and its correlation with clinicopathologic variables, risk factors and survival. *Journal of oral pathology & medicine*, 2008; 37(10): 593-598.
85. Sugiyama, M., Bhawal, U.K., Kawamura, M., Ishioka, Y., Shigeishi, H., Higashikawa, K. and Kamata, N. Human papillomavirus-16 in oral squamous cell carcinoma: clinical

- correlates and 5-year survival. *British Journal of Oral and Maxillofacial Surgery*, 45(2), pp.116-122.
86. Koyama, K., Uobe, K. and Tanaka, A., 2007. Highly sensitive detection of HPV-DNA in paraffin sections of human oral carcinomas. *Journal of oral pathology & medicine*, 2007; 36(1): 18-24.
87. Boy, S., Rensburg, E.J.V., Engelbrecht, S., Dreyer, L., Heerden, M. and Heerden, W. HPV detection in primary intra-oral squamous cell carcinomas—commensal, aetiological agent or contamination?. *Journal of oral pathology & medicine*, 2006;35(2): 86-90.
88. El-Mofty, S.K. and Patil, S. Human papillomavirus (HPV)-related oropharyngeal nonkeratinizing squamous cell carcinoma: characterization of a distinct phenotype. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 2006; 101(3): 339-345.
89. Kozomara, R., Jović, N., Magić, Z., Branković-Magić, M. and Minić, V. p53 mutations and human papillomavirus infection in oral squamous cell carcinomas: correlation with overall survival. *Journal of cranio-maxillo-facial surgery*, 2005;33(5): 342-348.
90. Zhang, Z.Y., Sdek, P., Cao, J. and Chen, W.T. Human papillomavirus type 16 and 18 DNA in oral squamous cell carcinoma and normal mucosa. *International journal of oral and maxillofacial surgery*, 2004; 33(1): 71-74.
91. Dahlgren, L., Dahlstrand, H.M., Lindquist, D., Högmö, A., Björnestål, L., Lindholm, J., Lundberg, B., Dalianis, T. and Munck-Wikland, E. Human papillomavirus is more common in base of tongue than in mobile tongue cancer and is a favorable prognostic factor in base of tongue cancer patients. *International journal of cancer*, 2004; 112(6): 1015-1019.
92. Correnti, M., Rivera, H. and Cavazza, M.E. Detection of human papillomaviruses of high oncogenic potential in oral squamous cell carcinoma in a Venezuelan population. *Oral diseases*, 2004; 10(3): 163-166.
93. Ritchie, J.M., Smith, E.M., Summersgill, K.F., Hoffman, H.T., Wang, D., Klussmann, J.P., Turek, L.P. and Haugen, T.H. Human papillomavirus infection as a prognostic factor in carcinomas of the oral cavity and oropharynx. *International journal of cancer*, 2003; 104(3): 336-344.
94. Sugiyama, M., Bhawal, U.K., Dohmen, T., Ono, S., Miyauchi, M. and Ishikawa, T. Detection of human papillomavirus-16 and HPV-18 DNA in normal, dysplastic, and malignant oral epithelium. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 2003; 95(5): 594-600.

95. Fregonesi, P.A., Teresa, D.B., Duarte, R.A., Neto, C.B., De Oliveira, M.R. and Soares, C.P. p16 (INK4A) immunohistochemical overexpression in premalignant and malignant oral lesions infected with human papillomavirus. *The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society*, 2003; 51(10): 1291-1297.
96. Chang, J.Y.F., Lin, M.C. and Chiang, C.P. High-Risk Human Papillomaviruses May Have an Important Role in Non-Oral Habits-Associated Oral Squamous Cell Carcinomas in Taiwan. *American journal of clinical pathology*, 2003;120(6): 909-916.
97. Chen, P.C.H., Kuo, C., Pan, C.C. and Chou, M.Y. Risk of oral cancer associated with human papillomavirus infection, betel quid chewing, and cigarette smoking in Taiwan—an integrated molecular and epidemiological study of 58 cases. *Journal of oral pathology & medicine*, 2002; 31(6): 317-322.
98. Tshako, K., Nakazato, I., Miyagi, J., Iwamasa, T., Arasaki, A., Hiratsuka, H., Sunakawa, H., Kohama, G. and Abo, T. Comparative study of oral squamous cell carcinoma in Okinawa, Southern Japan and Sapporo in Hokkaido, Northern Japan; with special reference to human papillomavirus and Epstein-Barr virus infection. *Journal of oral pathology & medicine*, 2000; 29(2): 70-79.
99. Aggelopoulou, E.P., Skarlos, D.I.M.O.S.T.H.E.N.I.S., Papadimitriou, C.O.N.S.T.A.N.T.I.N.A., Kittas, C.H.R.I.S.T.O.S. and TROUNGos, C.O.N.S.T.A.N.T.I.N.O.S. The Human Papilloma Virus DNA Detection in Oral Lesions in the Greek Population. *Anticancer research*, 1999; 19(2): 1391-1396.
100. Premoli-De-Percoco, G., Ramirez, J.L. and Galindo, I. Correlation between HPV types associated with oral squamous cell carcinoma and cervicovaginal cytology: An in situ hybridization study. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 1998;86(1): 77-81.
101. Koh, J.Y., Cho, N.P., Kong, G., Lee, J.D. and Yoon, K. p53 mutations and human papillomavirus DNA in oral squamous cell carcinoma: correlation with apoptosis. *British journal of cancer*, 1998; 78(3): 354.
102. Ibrahim, S.O., Bertelsen, B., Kalvenes, M.B., Idris, A.M., Vasstrand, E.N., Nilsen, R. and Johannessen, A.C. Expression of keratin 13, 14 and 19 in oral squamous cell carcinomas from Sudanese snuff dippers: lack of association with human papillomavirus infection. *Apmis*, 1998; 106(7-12): 959-969.

103. Mao, E.J., Schwartz, S.M., Daling, J.R., Oda, D., Tickman, L. and Beckmann, A.M. Human papilloma viruses and p53 mutations in normal, pre-malignant and malignant oral epithelia. *International journal of cancer*, 1996;69(2): 152-158.
 104. Cruz, I.B.F., Snijders, P.J.F., Steenbergen, R.D.M., Meijer, C.J.L.M., Snow, G.B., Walboomers, J.M.M. and Van der Waal, I. Age-dependence of human papillomavirus DNA presence in oral squamous cell carcinomas. *European journal of cancer part b: oral oncology*, 1996; 32(1): 55-62.
 105. Van Rensburg, E.J., van Heerden, W.F., Venter, E.H. and Raubenheimer, E.J. Detection of human papillomavirus DNA with in situ hybridisation in oral squamous carcinoma in a rural black population. *South African Medical Journal*, 1995;85(9).
 106. Shindoh, M., Chiba, I., Yasuda, M., Saito, T., Funaoka, K., Kohgo, T., Amemiya, A., Sawada, Y. and Fujinaga, K. Detection of human papillomavirus DNA sequences in oral squamous cell carcinomas and their relation to p53 and proliferating cell nuclear antigen expression. *Cancer*, 1995; 76(9): 1513-1521.
 107. Holladay, E.B. and Gerald, W.L. Viral gene detection in oral neoplasms using the polymerase chain reaction. *American journal of clinical pathology*, 1993; 100(1): 36-40.
 108. Zeuss, M.S., Miller, C.S. and White, D.K. In situ hybridization analysis of human papillomavirus DNA in oral mucosal lesions. *Oral surgery, oral medicine, oral pathology*, 1991; 71(6): 714-720.
 109. Greer Jr, R.O., Eversole, L.R. and Crosby, L.K. Detection of human papillomavirus-genomic DNA in oral epithelial dysplasias, oral smokeless tobacco-associated leukoplakias, and epithelial malignancies. *Journal of Oral and Maxillofacial Surgery*, 1990; 48(11): 1201-1205.
 110. Chang, F., Syrjänen, S., Nuutinen, J., Kärjä, J. and Syrjänen, K. Detection of human papillomavirus (HPV) DNA in oral squamous cell carcinomas by in situ hybridization and polymerase chain reaction. *Archives of dermatological research*, 1990; 282(8): 493-497.
 111. Dang, J., Manrique, H.X.Z., Veron, D. and Feng, Q. Oral Human Papillomavirus (HPV) Infection in Healthy Individuals and Patients with Head and Neck Squamous Cell Carcinoma (HNSCC). *Epidemiology (sunnyvale)*, 2015; 5(180): 2161-1165.
 112. Chocolatewala, N.M. and Chaturvedi, P. Role of human papilloma virus in the oral carcinogenesis: an Indian perspective. *Journal of cancer research and therapeutics*, 2009; 5(2): 71.
-