

International Journal of Scientific Research and Reviews

Effect of Foliar Application of Growth Hormones (IAA And GA₃) on Oil Percentage of Certain Oil Seed Crops

Mukundraj B. Patil*

Department of Botany, Late Ramesh Warpukar ACS college, Sonpeth, Dist. Parbhani (MS) INDIA
mukundrajpatil@gmail.com, +919657700237

ABSTRACT

Experiment was carried out during December 2013 to March 2014 to study effect of IAA and GA₃ on oil percentage of Groundnut (TAG-24), Soyabean (JS-335) and sunflower (SS-56). Sowing of seeds was carried out in plots of 1m x 1m. (One for control and 10ppm to 50ppm concentrations of IAA and GA₃). Different concentrations (10ppm to 50ppm) of IAA and GA₃ were applied twice 30 and 60 days after sowing. After harvesting oil percentage of the oil seeds was estimated using Soxhlet oil extractor. It was found that application of 40ppm and 50ppm GA₃ was able to enhance oil percentage in all the oil seeds. In soyabean application of 20ppm and 30ppm GA₃ as well as 40ppm IAA was able to increase oil percentage significantly.

KEYWORDS- IAA, GA₃, Oil percentage, Soxhlet oil extractor.

***Corresponding Author**

Mukundraj B. Patil

Head, Department of Botany

Late Ramesh Warpukar ACS college, Sonpeth, Dist. Parbhani (MS) INDIA

mukundrajpatil@gmail.com, +919657700237

INTRODUCTION

India is a country where main profession is Agriculture. In India variety of crops are sown, oil seeds contribute about fourteen percent of the total area under cultivation in India. India occupies prominent position both in area under cultivation and in production of oilseeds. India is the fourth largest edible oil economy in the world and contributes about ten percent of the world oil seed production. The growth in the domestic production of oilseeds has not been able to keep pace with the growth in the demand in the country. Groundnut, Soyabean and Sunflower are among the major nine edible oilseeds in India.

Auxins and GA₃ are important growth hormones in plants. Besides regulating growth plant hormones affect seed germination, growth, time of flowering, sex of the flowers, senescence of leaves fruits etc. They also affect fruit development, fruit ripening, plant longevity, fruit quality and even plant health^{1,2,3} Most of the workers have studied effect of growth regulators on seed germination^{4,5,6,7} and morphological or yield parameters^{8,9,10} Present work highlights on the effect of two important growth hormones on the oil percentage of three important oil yielding crops Groundnut, Soyabean and Sunflower.

MATERIALS AND METHODS

A Stock solution of 100 ppm GA₃ and IAA was prepared by dissolving 500mg into 500 ml of water (Before preparation of Stock solution GA₃ it was dissolved in small quantity of alcohol). It was diluted to prepare 10ppm, 20ppm, 30ppm, 40ppm and 50ppm IAA and GA₃.

To study effect of growth hormones on oil percentage plots of 1meter X 1meter (Three rows containing 11 plots each) were prepared. Seeds of Groundnut (TAG-24), Soyabean (JS-335) and sunflower (SS-56) were sowed in three separate rows with recommended spacing on 21 December 2013. Application of growth hormones was carried out twice first 30 days after and second 60 days after date of sowing. Spraying of growth hormones was carried out early in the morning with hand spray. Among eleven plots for a crop one was kept as control, ten plots were applied with gradually increasing (10ppm to 50ppm) concentration of IAA and GA₃. All the parameters except application of growth hormones were kept Constant.

Oil percentage was estimated using Soxhlet oil extractor after harvesting the crop.

Statistical Analysis- The mean, standard deviation (SD) and coefficient of variation (CV) has been calculated as described by Mungikar¹¹ SE was calculated as S.D. / \sqrt{n} (n = number of

observations), and Critical difference (CD) was calculated as S.E. multiplied with t value for n-1. (C.D.= S.E. x t value for n-1).

RESULT AND DISCUSSION

When groundnut was applied with different concentrations of GA₃ and IAA individually oil percentage was influenced by higher concentration of GA₃ (40ppm and 50ppm) and IAA (50ppm). Lower concentration of GA₃ and IAA has no significant effect to increase oil percentage of seed. 10ppm IAA and GA₃ showed decrease in the oil percentage which was non-significant. Maximum oil percentage was found in seeds of the plant treated with 50ppm GA₃ that was 39.1% thus 50 ppm GA₃ increased oil content of groundnut seed by 9.8%.

When soyabean was applied with different concentrations of GA₃ or IAA oil percentage was increased. GA₃ is more effective to increase oil percentage in Soyabean. Except 10ppm GA₃ all treatments of GA₃ showed increase in the oil percentage which was statistically significant. Maximum oil percentage was found in seeds of the plant treated with 50ppm GA₃ that was 15.8% thus 50 ppm GA₃ increased oil content of Soyabean seed by 21.5%. Same results were found by Travaglia¹².

When sunflower was applied with GA₃ or IAA individually oil percentage was influenced by higher concentration of GA₃ (40ppm and 50ppm) and 20ppm IAA. Lower concentration of GA₃ has no significant effect to increase oil percentage of sunflower seed. 30ppm IAA showed decrease in the oil percentage which was non-significant. Other treatment showed non-significant increase in the oil percentage. Maximum oil percentage was found in seeds of the plant treated with 50ppm GA₃ that was 34.3% thus seeds of Sunflower plants applied with 50 ppm GA₃ showed 15.5% increase in the oil content in sunflower than the seeds of control plant. The results are more promising than those recorded previously¹³ which showed no significant change in the oil content in Sunflower due to application of plant growth regulators.

Table 1 - Influence of different concentrations of growth hormones (IAA and GA3) on oil percentage of oil yielding crops

Treatment	Oil percentage		
	Groundnut	Soyabean	Sunflower
Control	35.6	13	29.7
GA 10ppm	35.4^	13.2^	29.7^
GA 20PPM	35.7^	13.6*	29.9^
GA 30PPM	36.1^	14.2**	30.5^
GA 40PPM	37.3**	14.9**	31.6**
GA 50PPM	39.1**	15.8**	34.3**
IAA 10PPM	35^	13.2^	29.8^
IAA20PPM	35.8^	13.1^	28.7*
IAA30PPM	36.1^	13.5^	29.3^
IAA40PPM	36.3^	13.6*	29.6^
IAA50PPM	36.7**	13.5^	30.2^
Mean	36.3	13.8	30.3
S.D.	1.1	0.9	1.5
C.V.	3.1	6.3	5
S.E.	0.3	0.3	0.5
C.D. 5 %	0.8	0.6	1
C.D. 1 %	1.1	0.8	1.4

(* Significant at p=0.05; ** Significant at p=0.01; ^ Non-significant)

ACKNOWLEDGEMENT- Author is thankful to UGC (WRO Pune) for financial assistance under Minor Research Projects.

REFERENCES

1. Christodoulou A, Weaver RJ, Pool RM. Relation of gibberellin treatment to fruit-.set, berry development, and cluster compactness in *Vitis vinifera* grapes. Proc. Amer. Soc. Hort. Sci. 1968; 92:301-10 .
2. Morris Justin R. Effects of gibberellic acid (GA₃) on yield and quality of grapes. Proceeding Arkansas State Hort. Soc. 108th Ann. Mtg., 1987; 76-79.
3. Horvitz S, Godoy C, López Camelo AF, Yommi A, Godoy C, . Application of gibberellic acid to ‘Sweet heart’ sweet cherries: Effects on fruit quality at harvest and during cold storage. Acta Hort.2003; 628: 311–316.
4. Chauhan JS, Tomar YK, Badoni Anoop, Singh N. Indrakumar, Ali Seema, Debarati, Rawat AS, Nautiyal VP. Morphology and Influence of Various Plant Growth Substances on Germination and Early Seedling Growth in *Macrotyloma uniflorum* (Lam.): Journal of American Science. 2009; 5(6) :43-50.
5. Patil JG, Ahire ML, Nikam TD. Influence of plant growth regulators on in vitro seed germination and seedling development of *Digitalis purpurea* L. The Asian and Australasian Journal of Plant Science and Biotechnology. 2012; 6 (Special Issue 1): 12-18.
6. Dhoran VS , Gudadhe SP. Effect of Plant Growth Regulators on Seed Germination and Seedling Vigour in *Asparagus sprengeri* Regel. International Research Journal of Biological Sciences. 2012; 1(7): 6-10
7. Patil Mukundraj B, Shailaja B. Bhosale. Effect of GA₃ on germination of certain oilseeds. Bioscience Discovery.2017; 8(3):483-485.
8. Emongor VE, Ndambole CM. Effect of gibberellic acid on performance of cowpea. African Crop Science Conference Proceedings. 2011;10: 87-92.
9. Rastogi Anu , Siddiqui A, Mishra BK, Srivastava M, Pandey R, Misra P, Singh M and Shukla S. Effect of auxin and gibberellic acid on growth and yield components of linseed (*Linum usitatissimum* L.) Crop Breeding and Applied Biotechnology. 2013; 13: 136-143.
10. Khairul Mazed HEM, Md. Hasanuzzaman Akand, Israt Jahan Irin, Jannatul Ferdous Moonmoon, Md. Hafizur Rahman. Effect of gibberellic acid on the growth and yield of cabbage (*Brassica oleracea* var. capitata L.). International Journal of Applied Research. 2015; 1(4):24-29.

11. Mungikar AM. Biostatistical Analysis. Saraswati Publ. Printing press, Aurangabad (M.S.) India 2003.
 12. Travaglia C, Reinoso H, Boltini R. Application of abscisic acid promotes yield in field-cultured soybean by enhancing production of carbohydrates and their allocation in seed. Crop and Pasture Science. 2009; 60: 1131-1135.
 13. Tahsin Nurettin, Tanko Kolev. Investigation on the effect of some plant growth regulators on Sunflower (*Helianthus annuus* L.). Journal of Central European Agriculture. 2005; 6(4): 583-586.
-