

International Journal of Scientific Research and Reviews

Bioaccumulation of heavy metals in tissues of some fishes in Umrar Dam on Umrar River in Umari district, Shahdol division in central India

Vandana Ram^{1*} & Binay Kumar Singh²

¹UGC Post Doctor Fellow

²Professor, Dept. of Zoology, Pandit S.N.Shukla, University, Shahdol (M.P.), India

Email: prayervandana29@gmail.com

ABSTRACT

The concentrations of heavy metals (Cu, Zn, Fe, Pb and Hg) were measured in the Muscles, liver, gills, kidney and gonad of fish species collected from Umrar Dam, Umari in central India. The levels of heavy metals varied significantly among fish species and organs. Muscles possessed the lowest concentration of metals. The essential metals as Cu were accumulated mainly in liver and gonad, Zn accumulated mainly in Gills and Liver, Fe were accumulated in all organs with little bit fluctuation in concentration while Pb accumulated in gill, liver and gonad and Hg levels were below detectable limits. The concentration of metals in the present fish organs within the permissible limits given by WHO and FAO but in case of Pb these are higher than the limits. This is also noticeable that the concentration of metals is higher in summer seasons while lowest concentrations are found in winter. This study reveals that fishes found in this dam are not suitable for human consumption it may cause severe health hazard because of high concentration of lead.

KEYWORDS: Heavy metals, Umrar Dam, Fishes, Health threads.

***Corresponding author**

Vandana Ram

UGC Post Doctor Fellow

Professor of Zoology, Pandit S.N.Shukla, University, Shahdol (M.P.), India

Email: prayervandana29@gmail.com

INTRODUCTION

The pollution of the aquatic environment with heavy metals has become a worldwide problem in recent years, because they are indestructible and most of them have toxic effect on organisms¹⁷. In the recent years, world consumption of fish has increased simultaneously with the growing concern of their nutritional and therapeutic benefits. In addition to its important source of protein, fish typically have rich contents of essential minerals, vitamins and unsaturated fatty acids¹⁹. The American Heart Association recommended eating fish at least twice per week in order to reach the daily intake of omega-3 fatty acids. However, fish are relatively situated at the top of the aquatic food chain; therefore they normally can accumulate heavy metals from food, water and sediments.⁰⁹

In the last few decades, the concentrations of the heavy metals in fish have been extensively studied in different parts of the world.¹³ Most of these studies concentrated mainly on the heavy metals in the edible parts that is fish muscles however other studies reported the distribution of metals in different organs like the liver, kidney, hearts, gonads, bone, digestive tract, gills and brain. The content of toxic heavy metals in fish can counteract their beneficial effects and may cause many adverse effects on human health this may include serious threats like renal failure, liver damage, cardiovascular diseases and even death.^{16, 18}

Heavy metals are implicated in neurological disorders especially in the foetus and in children, which can lead to behavioural changes and impaired performance in intelligent quotient (IQ) test¹⁵. The quality of the ecosystem has been degrading due to agriculture and human activities. Fish is an important component of the human diet in many villages and cities in shahdol division of central India and Umrar Dam is the very enormous source of fish culture and transportation of fishes to different region of shahdol division and other places too, for this reason, the results obtained from the study would provide information on background levels of metals in the fish species of the river Umrar, contributing to the effective monitoring of both environmental quality and the health of organisms inhabiting the river ecosystem.

It is therefore very important for study to be conducted on the concentration of heavy metals in the tissues of fishes of river Umrar in Umrar dam at Umara district in central India and check whether or not the concentration levels are within the permissible limits for human consumption in comparison to safety reference standards for the consumption of fish. Because in this area many people are dependent on fish as a food, especially fisher mans and it may cause severe health hazards.

MATERIALS AND METHODS

FISH SAMPLING

12 water samples and 24 commercial fish samples were used for study in three seasons of the year summer, winter and rainy, during two years (from 2015-16 to 2016-17) from every site. The collected species were *Labeo rohita*, *Rasbora daniconius* and *Ompok bimaculatus*. These fish species represent different biotopes and are economically important. Collected fish were immediately preserved in an ice box and transferred to the laboratory where they were classified, weighed, measured by total length and kept frozen at -20°C until further analysis. The fish and water samples collected from the different sites and analyzed at laboratory. Atomic Absorption Spectrophotometer (AAS) was used for the determination of the heavy metals in the tissue and water samples.

DETERMINATION OF METAL CONCENTRATIONS

Preparation of subsamples and analysis were made for metal analysis, frozen fish were partially thawed and each fish was dissected using stainless steel instruments. Muscles, Liver, Gills, Kidney and Gonad were taken out and dehydrated it, in oven, composite samples of 2–5 g were used for subsequent analysis.

The samples were digested with ultra pure nitric acid at 100°C until the solution become clear. The solution was made up to known volume with deionized distilled water and analyzed for Cu, Zn, Pb, Fe and Hg using the Atomic Absorption Spectrophotometer (AAS model ELICO, SL-168) the obtained results were expressed as mg/kg.

STATISTICAL ANALYSIS

Results were generally expressed as mean \pm standard deviation and one way ANOVA test was used to compare the data among seasons at the level of 0.05.

OBSERVATIONS

Concentrations of heavy metals (Cu, Zn, Pb, Fe and Hg) in the muscles, liver, gill, kidney and gonad of fish collected from the different sites of Umrar Dam.

As shown in Table-01, the contamination levels of these five metals were high in tissues. Specially, the concentration of Pb exceeding, FAO and WHO target values. Consumption of water as well as fish may create health problems related with Pb contamination may occur in human beings.

The accumulation of metals in a single species showed significant inter-specific variations in all metals. However it can be noticed that, different organs exhibited different patterns in metals accumulation. In other words, no single type of fish showed the highest metals in all

organs. Therefore, concentrations of metals between species were analyzed in single organ; all results showed significant variations between species. Furthermore, some fish from the same species collected from different sites also significantly accumulated different concentrations of metals. Variations of metals distribution in the studied fish can be summarized as the following:

Table-I:- Table showing mean (\pm SD) concentrations of heavy metals (mg/kg) in some organs of fish species collected from Umrar Dam.

FISH SPECIES	ORGANS	Metals	Cu	Zn	Fe	Pb
<i>Labeo rohita</i>	Muscles	Summer	0.292 \pm 0.001	0.045 \pm 0.001	0.325 \pm 0.004	0.840 \pm 0.001
		Rainy	0.021 \pm 0.001	0.043 \pm 0.001	0.296 \pm 0.002	0.789 \pm 0.000
		Winter	0.022 \pm 0.001	0.046 \pm 0.000	0.298 \pm 0.002	1.788 \pm 0.000
	Liver	Summer	1.463 \pm 0.001	2.204 \pm 0.096	0.883 \pm 0.003	1.846 \pm 0.001
		Rainy	0.299 \pm 0.001	0.102 \pm 0.000	0.802 \pm 0.001	1.841 \pm 0.001
		Winter	0.297 \pm 0.000	0.101 \pm 0.000	0.786 \pm 0.003	2.841 \pm 0.001
	Gills	Summer	1.298 \pm 0.000	2.873 \pm 0.016	0.228 \pm 0.002	3.081 \pm 0.000
		Rainy	0.998 \pm 0.001	1.852 \pm 0.001	0.202 \pm 0.001	3.068 \pm 0.001
		Winter	0.978 \pm 0.009	2.643 \pm 0.001	0.203 \pm 0.001	2.056 \pm 0.000
	Kidney	Summer	3.440 \pm 0.005	0.198 \pm 0.096	0.837 \pm 0.002	0.043 \pm 0.000
		Rainy	2.452 \pm 0.002	0.120 \pm 0.000	0.824 \pm 0.003	0.099 \pm 0.001
		Winter	2.401 \pm 0.001	0.197 \pm 0.000	0.794 \pm 0.005	0.085 \pm 0.000
	Gonad	Summer	1.177 \pm 0.001	0.218 \pm 0.115	0.903 \pm 0.003	3.050 \pm 0.001
		Rainy	1.065 \pm 0.001	0.122 \pm 0.000	0.853 \pm 0.002	3.046 \pm 0.000
		Winter	1.049 \pm 0.001	0.123 \pm 0.000	0.786 \pm 0.001	2.067 \pm 0.000
<i>Rasbora daniconius</i>	Muscles	Summer	0.226 \pm 0.001	0.201 \pm 0.002	1.224 \pm 0.002	0.788 \pm 0.001
		Rainy	0.201 \pm 0.000	0.200 \pm 0.001	1.204 \pm 0.001	0.776 \pm 0.000
		Winter	0.242 \pm 0.000	0.201 \pm 0.001	1.222 \pm 0.001	0.776 \pm 0.000
	Liver	Summer	1.562 \pm 0.000	1.204 \pm 0.000	1.788 \pm 0.010	0.806 \pm 0.001
		Rainy	1.432 \pm 0.002	0.199 \pm 0.001	0.778 \pm 0.001	1.804 \pm 0.001
		Winter	1.422 \pm 0.001	0.198 \pm 0.000	0.680 \pm 0.002	1.801 \pm 0.001
	Gills	Summer	1.280 \pm 0.001	2.883 \pm 0.001	0.205 \pm 0.004	4.832 \pm 0.000
		Rainy	0.988 \pm 0.001	2.879 \pm 0.001	0.201 \pm 0.001	3.765 \pm 0.001
		Winter	0.989 \pm 0.002	1.852 \pm 0.001	0.202 \pm 0.001	3.752 \pm 0.000
	Kidney	Summer	3.241 \pm 0.002	0.200 \pm 0.002	0.759 \pm 0.005	0.799 \pm 0.000
		Rainy	1.990 \pm 0.001	0.196 \pm 0.001	0.754 \pm 0.001	0.726 \pm 0.001
		Winter	2.984 \pm 0.004	0.194 \pm 0.000	0.752 \pm 0.002	0.725 \pm 0.000
	Gonad	Summer	2.210 \pm 0.002	0.218 \pm 0.000	2.886 \pm 0.003	3.786 \pm 0.001
		Rainy	1.122 \pm 0.001	0.216 \pm 0.001	1.979 \pm 0.002	3.740 \pm 0.000

		Winter	1.124±0.001	0.216±0.001	2.870±0.002	2.755±0.000
<i>Ompak bimaculatus</i>	Muscles	Summer	0.229±0.001	0.268±0.000	0.326±0.004	0.810±0.001
		Rainy	0.122±0.001	0.260±0.002	0.220±0.002	0.806±0.000
		Winter	0.117±0.000	0.260±0.002	0.363±0.002	0.805±0.000
	Liver	Summer	2.010±0.001	1.201±0.000	1.854±0.001	2.837±0.001
		Rainy	2.778±0.001	0.200±0.001	0.853±0.000	1.837±0.001
		Winter	1.763±0.002	0.200±0.001	1.705±0.010	1.826±0.001
	Gills	Summer	0.823±0.002	2.562±0.003	1.490±0.002	2.880±0.002
		Rainy	0.725±0.004	1.558±0.002	0.400±0.004	3.821±0.001
		Winter	0.993±0.003	2.502±0.003	0.339±0.002	2.798±0.002
	Kidney	Summer	4.001±0.001	0.255±0.001	0.685±0.004	0.844±0.000
		Rainy	3.994±0.005	0.253±0.002	0.782±0.002	0.790±0.001
		Winter	2.872±0.003	0.250±0.001	0.699±0.002	0.789±0.000
	Gonad	Summer	2.452±0.002	0.288±0.001	2.738±0.002	3.855±0.001
		Rainy	0.654±0.001	0.270±0.000	0.863±0.002	2.789±0.001
		Winter	1.582±0.002	0.271±0.000	1.764±0.000	2.789±0.001

Table-II:-Table showing maximum permissible limit (MPL) of heavy metals in fish tissues (mg/kg) according to international standards.

	Cu	Zn	Fe	Pb
FAO/WHO limit(2011)	30	40	43	0.5
*FAO(1983)	30	30	---	0.5
**WHO 1989	30	100	100	2
***FSAI(2009)	-	-	-	0.3
****FSSAI(2011)	30	50		2.5

*Food and Agriculture Organization

**World Health Organization

***Food Safety Authority of Ireland

****Food Safety and Standard Authority of India

Copper (Cu)

The copper concentration in the tissues of *Labeo rohita* is highest in the summer season in about all organs taken for observation. Copper concentrations are reached the highest level in kidney in summer (3.440±0.005) whereas lowest concentrations are found in muscles in rainy season (0.021±0.001). In *Rasbora daniconius* copper concentration reached the highest in kidney in summer (3.241±0.002) and lowest are found in muscle in rainy season (0.201±0.000) and in *Ompak*

bimaculatus highest concentration is in kidney (4.001 ± 0.001) in summer whereas lowest is in muscles in rainy season (0.122 ± 0.001). Copper concentrations varied highly significantly ($P < 0.001$) from season to season in organs of all experimental fishes.

Zinc (Zn)

The Zinc concentration in the tissues of *Labeo rohita* is highest in summer whereas lowest concentrations are found in winter. Zn concentrations are reached the highest level in gills in summer (2.873 ± 0.016) whereas lowest concentrations are found in muscles in rainy season (0.043 ± 0.001). Zn concentrations varied highly significantly ($P = 0.001$) from season to season in organs of *Labeo rohita*. In *Rasbora daniconius* Zn concentration reached the highest in gills in summer (0.194 ± 0.000) and lowest are found in Kidney in winter season (2.883 ± 0.001). Zn concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Rasbora daniconius* and in *Ompak bimaculatus* highest concentration is in gills (4.001 ± 0.001) in summer whereas lowest is in liver in rainy and winter seasons (0.200 ± 0.001). Zn concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Ompak bimaculatus*.

Iron (Fe)

The Fe concentration in the tissues of *Labeo rohita* is highest in summer whereas lowest concentrations are found in winter. Fe concentrations are reached the highest level in gonad in summer (0.903 ± 0.003) whereas lowest concentrations are found in gills in rainy season (0.202 ± 0.001). Fe concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Labeo rohita*. In *Rasbora daniconius* Fe concentration reached the highest in gonad in summer (2.886 ± 0.003) and lowest are found in gills in rainy season (0.201 ± 0.001). Fe concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Rasbora daniconius* and in *Ompak bimaculatus* highest concentration is in gonad (2.738 ± 0.002) in summer whereas lowest is in muscle in rainy seasons (0.220 ± 0.002). Fe concentrations varied significantly ($P = 0.05$) from season to season in organs of *Ompak bimaculatus*.

Lead (Pb)

The Pb concentration in the tissues of *Labeo rohita* is highest in summer whereas lowest concentrations are found in winter. Fe concentrations are reached the highest level in gills in summer (3.081 ± 0.000) whereas lowest concentrations are found in kidney in summer season (0.043 ± 0.000). Pb concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Labeo rohita*. In *Rasbora daniconius* Pb concentration reached the highest in gills in

summer (4.832 ± 0.000) and lowest are found in kidney in winter season (0.725 ± 0.000). Pb concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Rasbora daniconius* and in *Ompak bimaculatus* highest concentration is in gonad (3.855 ± 0.001) in summer whereas lowest is in kidney in winter seasons (0.789 ± 0.000). Pb concentrations varied highly significantly ($P < 0.001$) from season to season in organs of *Ompak bimaculatus*.

DISCUSSIONS

Knowledge of heavy metal concentrations in fish is important for both human consumption and nature management in this study; we examined metals in tissue of experimental fishes to evaluate heavy metal concentrations in Umrar dam. It is also aimed to investigate whether metal concentrations varied seasonally in the study.

Present study showed the lowest concentration of metals in muscle. The essential metals as Cu were accumulated mainly in liver, kidney and gonad, Zn accumulated mainly in Gills and Liver, Fe were accumulated in all organs with little bit fluctuation in concentration and Pb were accumulated mainly in gills and gonad.

The accumulation of metals in liver is probably linked to its role in metabolism²³ high levels of Cu and Zn in hepatic tissues are usually related to a natural binding proteins such as metallothioneins¹¹ which act as an essential metal store as Zn and Cu to fulfil enzymatic and other metabolic demands²² while Fe tends to accumulate in hepatic tissues due to the physiological role of the liver in blood cells and haemoglobin synthesis¹¹. On the other hand, the liver also showed high levels of non-essential metals such as Pb to displace the normally metallothioneins associated metals in hepatic tissues⁰². Previous studies also show similar trends to accumulate high level of essential and non-essential metals in liver cells in fishes.^{23,07,03,12,06}

Presence of these metals in gills shows that gills are main route of metal ion exchange from water²⁰ as they have large surface area and facilitate rapid diffusion of toxic metals⁰⁵. Therefore it is suggested that metals accumulated in gills are mainly concentrated from water specially Pb and Zn, previous studies also show the similar things as^{14,04,01,20,07}.

In present study highest concentrations of metals are found in summer season whereas lowest concentrations are in rainy and winter seasons. The increase of heavy metal levels in summer and winter could be related to increasing physiological activity of fish during summer, primarily caused by the increasing water temperature and decrease in water level because of agricultural activities and domestic activities or high temperature as well. For all seasons we found Cu, Zn and Fe are under the limits of WHO, FAO, FSAI and FSSAI while Pb concentrations are higher than the permissible limits.

It is well known that muscles are not active site for metal biotransformation and accumulation⁰⁸ but in polluted aquatic habitats the concentration of metals in fish muscles may exceed the permissible limits for human consumption and imply severe health threats.

CONCLUSIONS

The concentration of metals in the present fish organs within the permissible limits given by WHO and FAO but in case of Pb these are higher than the limits. So health risk analysis of heavy metals in the edible part (muscle) of the fish indicated safe levels for human consumption and concentrations in the muscles are generally accepted by the international legislation limits however, whole fish with all organs specially small size fishes and the ovary is consumed by many people's of this region so study reveals that fishes found in this dam are not suitable for human consumption it may cause severe health threats.

ACKNOWLEDGMENT

Author are thankful to my supervisor Dr. Binay Kumar Singh for his guidance and also thankful to University Grants Commission New Delhi for the award of Post Doctoral Fellowship vide letter No: F./PDFSS-2014-15-SC-MAD-9038.

REFERENCES

1. Abu Hilal AH, Ismail NS. Heavy metals in eleven common species of fish from the Gulf of Aqaba, Red Sea. *Jordan J Biol Sci* 2008; 1(1):13-8.
2. Amiard JC, Amiard-Triquet C, Barka S, Pellerin J, Rainbow PS. Metallothioneins in aquatic invertebrates: their role in metal detoxification and their use as biomarkers. *Aquat Toxicol* 2006; 76:160-202.
3. Amundsen PA, Staldvik FJ, Lukin AA, Kashulin NA, Popova OA, Reshetnikov YS. Heavy metal contamination in freshwater fish from the border region between Norway and Russia. *Sci Total Environ* 1997; 201:211-24.
4. Avenant-Oldewage A, Marx HM. Bioaccumulation of chromium, copper and iron in the organs and tissues of *Clarias gariepinus* in the Olifants River, Kruger National Park. *Water Sanit* 2000; 26: 569-82.
5. Dhaneesh KV, Gopi M, Ganeshamurthy R, Kumar TTA, Balasubramanian T. Bioaccumulation of metals on reef associated organisms of Lakshadweep Archipelago. *Food Chem* 2012; 131: 985-91.

6. Dural M, Goksu MZL, Ozak AA. Investigation of heavy metal levels in economically important fish species captured from the Tuzla lagoon. *Food Chem* 2007; 102:415-21.
7. Eisler R. *Compendium of trace metals and marine biota*. Amsterdam: Vertebrates Elsevier; 2010.
8. Elnabris KJ, Muzyed SK, El-Ashgar NM. Heavy metal concentrations in some commercially important fishes and their contribution to heavy metals exposure in Palestinian people of Gaza Strip (Palestine). *J Assoc Arab Univ Basic Appl Sci* 2013; 13:44-51
9. F. Yilmaz, N. Ozdemir, A. Demirak, A.L. Tuna Heavy metal levels in two fish species *Leuciscus cephalus* and *Lepomis gibbosus*, *Food Chem*, , 2007; 100: 830-835.
10. FAO/WHO; Joint FAO/WHO food standards programme codex committee on contaminants in foods, fifth. Session 2011; 64-89.
11. Gorur FK, Keser R, Akcay N, Dizman S. Radioactivity and heavy metal concentrations of some commercial fish species consumed in the Black Sea Region of Turkey. *Chemosphere* 2012; 87: 356-61.
12. Jose U, Carmen I, Jose M, Ignacio G. Heavy metals in fish (*Solea vulgaris*, *Anguilla anguilla* and *Liza aurata*) from salt marshes on the southern Atlantic coast of Spain. *Environ Int* 2004; 29:949-56.
13. K.J. Elnabris, S.K. Muzyed, N.M. El-Ashgar Heavy metal concentrations in some commercially important fishes and their contribution to heavy metals exposure in Palestinian people of Gaza Strip (Palestine), *J Assoc Arab Univ Basic Appl Sci*, 2013; 13: 44-51
14. Kargin F. Metal concentrations in tissues of the freshwater fish *Capoeta barroisi* from the Seyhan River (Turkey). *Bull Environ Contam Toxicol* 1998; 60:822-8.
15. Landner and Lindstrom: Zinc in society and in the environment. Miljoforskargruppen, Stockholm, 1998;160.
16. M. Al-Busaidi, P. Yesudhasan, S. Al-Mughairi, W.A.K. Al-Rahbi, K.S. Al-Harthy, N.A. Al-Mazrooei, *et al.* Toxic metals in commercial marine fish in Oman with reference to national and international standards, *Chemosphere*, 2011; 85(1):67-73
17. MacFarlane, G.R. & M.D. Burchett. 2000. Cellular distribution of Cu, Pb and Zn in the Grey Mangrove *Avicennia marina* (Forsk.) Vierh. *Aquatic Botany* 2011;68: 45–59.
18. M.S. Rahman, A.H. Molla, N. Saha, A. Rahman Study on heavy metals levels and its risk assessment in some edible fishes from Bangshi River, Savar, Dhaka, Bangladesh, *Food Chem*, 2012; 134(4) :1847-1854.

19. P. Kris-Etherton, W. Harris, L. Appel. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease, *Circulation*, 2002; 106: 2747-2757.
 20. Qadir A, Malik RN. Heavy metals in eight edible fish species from two polluted tributaries (Aik and Palkhu) of the River Chenab, Pakistan. *Biol Trace Elem Res* 2011; 143:1524-40.
 21. R.J. Medeiros, L.M. dos Santos, A.S. Freire, R.E. Santelli, A.M.C.B. Braga, T.M. Krauss, *et al.* Determination of inorganic trace elements in edible marine fish from Rio de Janeiro State, Brazil, *Food Control*, 2012; 23: 535-541.
 22. Roesijadi G. Metallothionein and its role in toxic metal regulation. *Comp Biochem Physiol C* 1996; 113(2):117- 23.
 23. S. Zhao, C. Feng, W. Quan, X. Chen, J. Niu, Z. Shen. Role of living environments in the accumulation characteristics of heavy metals in fishes and crabs in the Yangtze River Estuary, China, *Mar Pollut Bull*, 2012; 64:1163-1171.
 24. Tiimub, Benjamin Makimilua, Mercy Ananga Dzifa Afua , Tiimub & Dzifa Afua © American Determination of Selected Heavy Metals and Iron Concentration in Two common Fish Species in Densu River at Weija District in Greater Accra Region of Ghana. *American International Journal of Biology*, July 2013; 1(1):45-55
-