

## International Journal of Scientific Research and Reviews

### Effect of precursor feeding on quercetin production in callus cultures of *Merremia aegyptia* and *Merremia dissecta*.

Ridhi Joshi<sup>1\*</sup> and Vidya Patni<sup>2</sup>

<sup>1\*</sup> Assistant professor of Botany, Jaipur National University, Jagatpura, Jaipur, Rajasthan, India.

Email: [ridhi.joshi316@gmail.com](mailto:ridhi.joshi316@gmail.com), Ph:8955191724

<sup>2</sup> Associate Professor, Plant Pathology, Tissue Culture and Biochemistry Laboratory, Department of Botany, University of Rajasthan, Jaipur, Rajasthan, India. Email: [vidyapatni@gmail.com](mailto:vidyapatni@gmail.com),

Ph:9829059868

#### ABSTRACT

L-Phenylalanine is an aromatic amino acid and a substrate of phenylalanine ammonia lyase (MW 270-330kDa.) (EC 4.3.1.24) that catalyzes L-phenylalanine into ammonia and trans-cinnamic acid in the biosynthesis of plant phenolic compounds as it is involved in phenylpropanoid pathway of plants producing various flavonoids, phenylpropanoids, and lignin in plants. Optimisation of suitable concentrations of plant growth regulator and harvest time was accomplished with the aim of enhancing the production of quercetin in callus cultures of *Merremiaaegyptia* and *Merremiadissecta* through precursor feeding. *Merremiadissecta* callus cultures showed 1.2 fold increase in quercetin content in precursor fed cultures and stem cultures of *Merremiaaegyptia* showed two fold increase in quercetin content supplemented with 50 mg/100ml PA analysed through high performance thin layer chromatography.

**KEYWORDS:** Phenyl alanine, *Merremiadissecta*, *Merremiaaegyptia*, HPTLC

**\*Corresponding author:**

**Ridhi Joshi**

Assistant professor of Botany,

Jaipur National University,

Jagatpura, Jaipur, Rajasthan, India.

Email: [ridhi.joshi316@gmail.com](mailto:ridhi.joshi316@gmail.com),

Ph:8955191724

## INTRODUCTION:

Plant cell and organ culture has emerged as potential area of secondary metabolites production that are used as pharmaceuticals, agrochemicals, flavors, fragrances, coloring agents, bio pesticides, and food additives. The selection of high-producing cells and optimization of medium parameters such as suitable medium, salt, sugar, nitrogen, phosphate, and plant growth regulator levels; and physical factors such as temperature, illumination, light quality, medium pH, agitation, aeration, and environmental gas (e.g., oxygen, carbon dioxide, and ethylene) is required for their production. Several chemical and physical factors have been recognized, which could influence biomass accumulation and synthesis of secondary metabolites in plant cell and organ cultures such as type of culture medium, suitable salt strength of the medium, types and levels of carbohydrates, nitrate levels, phosphate levels and growth regulator levels<sup>2</sup>.

The production of secondary metabolites by cell suspension culture is not always rewarding; thus, organ culture methods (e.g., root, embryo, and shoot culture methods) have been developed for various plant species as alternatives for production of secondary metabolites<sup>3,4</sup>. Many plant cell cultures have also been used to convert precursors into products by utilizing pre-existing enzyme systems such as the addition of loganin, tryptophan, and tryptamine enhanced the production of secologanin<sup>5</sup> and in dole alkaloids<sup>6</sup> in *Catharanthus roseus* suspension cultures. Effect of phenylalanine on flavonoids has been studied in tissue cultures of *Hydrocotyle bonariensis*<sup>7</sup> and *Silybum marianum*<sup>8</sup>. However for better results, factors such as the timing and concentration of the precursor should be considered prior addition to the cell culture medium. Sophisticated modern quantification techniques such as UV-Vis spectrophotometry, TLC, HPTLC, HPLC, and NMR provide quantitative information on cells and organs under culture conditions. For example, TLC, UV, HPLC and HPTLC analyses have been used for quantification of metabolites in cell suspensions and *in vitro* regenerated organs of *Hypericum perforatum*<sup>9</sup>. Callus cultures were fed with phenylalanine and given stress conditions for a period of 3-7 weeks. The level of enhancement in the augmented callus cultures were determined and compared to controlled callus cultures.

## MATERIALS AND METHODS

Unorganised static callus cultures were established from young leaf and stem explants on Murashige and Skoog's (MS) basal medium supplemented with NAA (1.0 mg/l) in combination with BAP (0.5 mg/l) in *M. aegyptia* and NAA (2.0 mg/l) in combination with BAP (0.5 mg/l) in *M. dissecta*. Obtained callus tissues were then transferred to fresh MS medium with additives singly supplemented with concentrations of phenylalanine (25 mg/100 ml, 50 mg/100 ml, 75 mg/100 ml and 100 mg/100 ml). These cultures were allowed to grow upto their maximum growth age. The

callus tissues were dried, till a constant weight was achieved, powdered, weighed and subjected to quercetin analysis. Five replicates were used in each case and their mean values were taken into consideration. The dried samples were separately soxhlet extracted in 80% methanol (100 ml/gm dry weight) on a water bath for 24 hrs. Each of the extracts was concentrated and reconcentrated in petroleum ether (40°-60°C) (fraction-I), ethyl ether (fraction-II) and ethyl acetate (fraction-III) in succession. Each of the steps was repeated thrice to ensure complete extraction in each case. Fraction I was rejected since it was rich in fatty substances whereas fraction II and fraction III of each of the test samples was hydrolysed by refluxing with 7% H<sub>2</sub>SO<sub>4</sub> (10 ml/gm residue) for 5 hours. The mixture was filtered and the filtrate was extracted with ethyl acetate in a separating funnel. The ethyl acetate layer was washed with distilled water till neutrality and dried. The residues were taken up in small volumes of methanol separately and then subjected to various tests for quercetin.

## RESULTS:

Identification and quantification of bioactive secondary metabolite quercetin was carried out using HPTLC techniques. Results indicated a gradual increase in growth index upto five weeks; after that it declined. Highest GI was reported in callus cultures obtained when fed with phenylalanine (50 mg/100 ml) at fifth week of growth. Further quantification of quercetin was confirmed by high performance thin layer chromatography (HPTLC) and peak identification was carried out on the basis of an authentic sample of quercetin. R<sub>f</sub> value (0.51) of quercetin isolated from the plant samples coincided with the R<sub>f</sub> value of standard quercetin (0.51) in the chromatogram.

## DISCUSSION AND CONCLUSION

Flavonoids are one of the major secondary metabolites imparting medicinal values for human beings. Of these, quercetin is considered as an active ingredient which has many biological roles. In the enhancement study, It has been observed that combination as well as concentration of different plant growth regulators in combination affects the synthesis of secondary metabolites and hence, optimization of medium could enhance the production of secondary metabolites 10,11 . Similarly effect of phenylalanine on quercetin content has been studied in tissue cultures of *Gossypium* cultivars <sup>12</sup> , *Balanites aegyptica* <sup>13</sup> and *Cassia angustifolia* <sup>14</sup> . Enhancement of quercetin via precursor feeding technique in *Pluchea lanceolata* has also been done <sup>15</sup> . According to this study, Phenylalanine <sup>16</sup> and Cinnamic acid <sup>17</sup> are reported to be the precursors of quercetin. Addition of precursors (Phenylalanine and Cinnamic acid) in *in vitro* callus cultures of *Pluchea lanceolata* resulted in multifold increase in quantity of quercetin. In our investigation, callus cultures of *M. aegyptia* and *M. dissecta* were fed with phenyl alanine (25-100 mg/ml) and were given stress

conditions for obtaining better results. Five weeks of feeding and experimental condition resulted in higher GI and produced about 1 to 2 fold higher quantity of quercetin when tested as compared to unelicited controls. *Merremia dissecta* callus cultures showed 1.2 fold increase in quercetin content and stem cultures of *Merremia aegyptia* showed two fold increase in quercetin content in five weeks old callus cultures. 4.79mg/g and 7.44 mg/g dry weight quercetin were observed in callus tissue fed with 50 mg PA/100 ml medium as compared to 3.75mg/g and 3.71 mg/g dry weight of quercetin content observed in control cultures of both the samples. Therefore, the protocol described in this report is expected to contribute to the future studies for large scale production of certain biologically active plant metabolites in various species from their in vitro propagated elite materials. Accumulation of quercetin in callus cultures can be exploited for its large-scale production in a possible array of health promoting benefits as anti-oxidant, anti-inflammatory and anti-cancerous agents in future.

## ACKNOWLEDGEMENTS

Financial support provided by the University Grants Commission (UGC), New Delhi and support from Department of Botany for successful conduction of work is gratefully acknowledged.

## REFERENCES:

1. Rao RS, Ravishankar GA. Plant tissue cultures; chemical factories of secondary metabolites. *Biotechnol. Adv.*, 2002; 20: 101-153.
2. Dornenburg H, Knorr D. Strategies for improvement of secondary metabolite production in plant cell cultures. *Enz Microb Technol* 1995;17:674–684.
3. Stafford A, Morris P, Fowler MW. Plant cell biotechnology: a perspective. *Enzyme Microb Technol* . 1986, 8:578–587.
4. Baque MA, Moh SH, Lee EJ, Zhong JJ, and Paek K. Production of biomass and useful compounds from adventitious roots of high-value added medicinal plants using bioreactor. *Biotech adv.* 2012; 101-105.
5. Verpoorte R, Contin A and Memelink J. Biotechnology for the production of plant secondary metabolites. *Phytochem Reviews*. 2002; 1:13-25.
6. Contin A, Van der Heijden R, Verpoorte R. Effects of alkaloid precursor feeding and elicitation on the accumulation of secologanin in a *Catharanthus roseus* cell suspension culture. *Plant Cell Tiss Org Cult* .1999; 56: 111–119

7. Moreno PRH, Van der Heijden R, Verpoorte R. Effect of terpenoid precursor feeding and elicitation on formation of indole alkaloids in cell suspension cultures of *Catharanthus roseus*. Plant Cell Rep . 1993b; 199(12): 702–705
8. Masoumian M, Arbakariya A, Syahida A, And Maziah M. Effect of precursors on flavonoid production by *Hydrocotyle bonariensis* callus tissues. African J. Biotech., 2011; 10: 6021-6029.
9. Rahimi S, Hasanloo T, Najafi F and Nejad RAK. Enhancement of *Silymarin* accumulation using precursor feeding in *Silybum marianum* hairy root cultures, Plant Omics J., 2011; 4: 34-39.
10. Pasqua G, Avato P, Monacelli B, Santamaria AR, Argentieri MP Metabolites in cell suspension cultures, calli, and *In vitro* regenerated organs of *Hypericum perforatum* cv. Topas. Plant Sci. 2003; 165: 977–982.
11. Karuppusamy S. A review on trends in production of secondary metabolites from higher plants by *Invitro* tissue, organ and cell cultures. J of Medicinal Plants Res. 2009; 3(13): 1222-1239.
12. Ashrafi S, Mofid MR, Otroshi M, Ebrahimi M and Khosroshahli M. Effects of plant growth regulators on the callogenesis and Taxol production in cell suspension of *Taxus baccata* L. Trakia J. Sci., 2010; 8: 36–43
13. Goyal S. Endogenous bioactive and bioregulatory metabolites in *Gossypium* varieties grown in tissue culture. Ph.D. Thesis, MDS University, Ajmer, 1997.
14. Bidawat S. Evaluation of *Balanites aegyptiaca* (Hingota) an arid zone medicinal plant for phytochemically important metabolites. Ph.D. Thesis, M.D.S. University, Ajmer, India. 2006.
15. Reddy A, Rao HR, Goswami A. Effect of phenylalanine on production of flavonoids in tissue culture of medicinal plants *Cassia angustifolia* and *Ailanthus excelsa*. In: Abstract Vol. 30, All India Botanical Conference, Nov. 28- 30, Jiwaji University, Gwalior. 2007; 171-174.
16. Arya D, and Patni V. Comparative analysis of total flavonoids and quercetin content *In vivo* and *Invitro* and enhancement of quercetin *via* precursor feeding in *Pluchea lanceolata* Oliver & Hiern. Int. J. Pharmacy and Pharma. Sci. 2013; 5(3):1-5.
17. Barz W. Plant Tissue Culture and its Biotechnological applications. In : Barz W, Reinhard E, Zenk MH, editors. Catabolism of endogenous and exogenous compounds by plant cell cultures. Springer-Verlag, New York. 1977; 153-171.
18. Grisebach H. and Grambow HJ Biosynthesis of flavonoids XV Occurrence and biosynthesis of flavonoids in *Datisca cannabina*. Phytochem. 1968; 7: 51-56.