

## *International Journal of Scientific Research and Reviews*

### **Antifungal Efficacy of Essential Oils against common bean Anthracnose caused by *Colletotrichum lindemuthianum***

**Bhardwaj Jyoti<sup>1\*</sup>, Seth MK<sup>1</sup>, Akhtar Jameel<sup>2</sup> and Chand Dinesh<sup>3</sup>**

<sup>1</sup>Department of Bio-Science, Himachal Pradesh University, Shimla-171005. India

<sup>2</sup>Plant Quarantine Division, National Bureau of Plant Genetic Resources, New Delhi -110012. India

<sup>3</sup>National Bureau of Plant Genetic Resources, Regional Station, Akola, Maharashtra-444104. India

#### **ABSTRACT**

Fifteen essential oils were evaluated to determine the fungitoxicity against anthracnose of common bean caused by *Colletotrichum lindemuthianum* (Sacc. & Magnus) Briosi & Cavara. For screening of antifungal susceptibility in each essential oil, healthy control and fungicide control were set to verify growth inhibition zone diameters. After bioassay, oils of lemon grass and peppermint were found to be most effective with complete inhibition followed by oils of winter green and geranium against test fungus. If anthracnose is found in the field, spread of disease can be limited by the use of fungicides. To protect agricultural crops enormous amount of synthetic pesticides are used worldwide resulting ill-effects on the life and life supporting systems. Efforts are thus being made world over to replace these synthetic chemicals with safer and eco -friendly alternatives. The plant world comprises of a rich storehouse of biochemical that could be tapped for use as fungicide. Given the abundance of flora in India, botanical fungicides have a great potential in integrated disease management in the context of eco-friendliness and sustainable agro-ecosystem.

**KEYWORDS:** Bean anthracnose, *Colletotrichum lindemuthianum*, essential oil, antifungal activity

#### **\*Corresponding Author:**

**Jyoti Bhardwaj**

Department of Bio-Sciences

Himachal Pradesh University, Summer Hill

Shimla, 171005.

**E-mail:** [redjyoti@yahoo.com](mailto:redjyoti@yahoo.com)

## **INTRODUCTION**

A number of economically significant pests, pathogens are seed-borne and seed transmitted<sup>1</sup>. A list of more than 1500 fungi belonging to 287 genera reported to be seed-borne in 534 crops of cereals, millets, vegetables and ornamental etc<sup>2</sup>. Seed transmission of pathogens is the most important mode of long distance dissemination and carry over from season to season. World over, pests (especially weeds, pathogens and insects) are the largest competitor of agricultural crops and severely reduce the crop production in the range of 25-50%<sup>3</sup>. Seed-borne /transmitted fungi thus play a crucial role in the development of disease in the field and also in the trans-boundary movement of pathogens along with the exchange of plant genetic resources (PGR) to be utilized in crop improvement programmes in the country.

About one third crop losses worldwide are attributed to plant pathogenic micro-organisms including fungi, bacteria and viruses. Bean anthracnose, caused by the fungus *Colletotrichum lindemuthianum*, is a major disease of beans (*Phaseolus vulgaris* L.), causing serious crop loss in many parts of the world. In 1921, M. F. Barrus of Cornell University demonstrated that bean anthracnose is seed borne. This information resulted in the widespread use of plant treated, diseased free seed to minimize the risk of anthracnose infection.

The use of plant-derived pesticides against crop pests both on the field and during post-harvest is now emerging as one of the important means to be used in crop protection under an Integrated Pest Management framework following the multiple global challenges created by synthetic chemical pesticides. Interest in the antifungal activity of essential oils has increased markedly in recent years also because of drug resistance in microbes. The volatile (vapor) components of several essential oils have been demonstrated to have potent antifungal activity. Essential oils from various aromatic plants are known to show a wide spectrum of anti-microbial activity. Antibacterial properties in the floral petals of many higher plants have been reported<sup>4</sup>. Efficacy of essential oils for control of seed-borne pathogens has been evaluated<sup>5</sup>.

Essential oils of some higher plants have recently been proved successful in providing effective control of storage losses and post harvest fungal diseases of plant origin<sup>6, 7, 8</sup>. Very little work has been done in case of *Colletotrichum* spp.<sup>9, 10, 11, 12, 13</sup> and need to explore the scope of utilization of biopesticides to control anthracnose of common bean. Effective management of a plant disease is a key to save plants from diseases caused from microbes, since plants are significant as they are both economical and aesthetic.

## MATERIAL AND METHODS

### *Collection of samples*

Essential oils were procured from Jain super store, Palika Bazar, Connaught place, New Delhi for bio-assay. Seed material from different districts (Chamba, Shimla and Hamirpur) of Himachal Pradesh was collected to get a pure culture of *Colletotrichum lindemuthianum*.

### *Isolation and maintenance of pathogen culture*

Seeds of *Phaseolus vulgaris* were examined for seed health status. After visual and stereobinocular microscopical detection seeds with disease symptoms like discoloration, pigmentation, spores/conidia and fruiting body fruiting bodies (acervuli) and subjected to blotter test by placing them on 4 layers of moistened blotter discs of 110 mm. Plastic Petri plates were incubated for 7 days at 21<sup>o</sup> ( $\pm 1$ ) C under alternate cycle of light and darkness (12h each)<sup>14</sup>. Mycelium were picked up with the help of sterilized needle and cultured on PDA in sterilized Petri plates. Pure culture were prepared and maintained for inhibition studies.

### *Screening of extracts (antifungal assay)*

15 essential oils from 11 plant families were bio-assayed by poisoned food technique<sup>15</sup>. Appropriate quantity of the desired essential oil was added to a pre-autoclaved adequately cooled PDA. Tween 20 was also added to the media to get better suspension. The medium was then dispensed into sterilized petriplates. PDA without any extract served as control, bavistin (0.02%), mancozeb (0.25%) and a binary mixture of bavistin & mancozeb, (1:1 w/w)(B+M) served as fungicide control for comparison. Five replicates were kept for each treatment and incubated for seven days at 21<sup>o</sup> ( $\pm 1$ ) C at alternate cycle of light and darkness. Radial growth of colonies was measured at two points along the diameter of the fungal colony. The growth of the colonies in control sets was compared with that of various treatments and calculated the percent inhibition by using this formula:-

$$\text{Percent Inhibition} = \frac{C-T}{C} \times 100$$

Where, C= radial diameter of the colony in control, T= radial diameter of the colony in treatment

Completely randomized design was used in carrying out ANOVA from which LSD at 5% level was computed for each test concentration.

## RESULT AND DISCUSSION

**Table No.1: In-vitro effect of essential oils with radial growth (mm) and percent inhibition of *Colletotrichum lindemuthianum* on eighth day of incubation**

Botanical name/Family	Common name	Concentration (PPM)	Radial growth (mm)	Percent Inhibition
<i>Pimpinella anisum</i> (Apiaceae)	Anethole Oil	500	72.75	19.00
		1000	66.40	26.22
<i>Cymbopogon winterianus</i> (Poaceae)	Citronella Oil	500	74.60	17.00
		1000	67.40	25.11
<i>Tagetes erecta</i> (Asteraceae)	Tagetes Oil	500	75.40	16.00
		1000	71.20	20.89
<i>Cymbopogon martini</i> (Poaceae)	Palma Rosa Oil	500	75.00	16.67
		1000	66.60	26.00
<i>Juniperus virginiana</i> (Pinaceae)	Cedar Wood Oil	500	74.20	17.56
		1000	68.80	23.56
<i>Laurus nobilis</i> (Lauraceae)	Kranj Oil	500	75.40	16.22
		1000	66.20	26.44
<i>Ocimum basilicum</i> (Lamiaceae)	Basil Oil	500	69.20	23.11
		1000	74.80	16.89
<i>Gossypium.hirsutum</i> (Malvaceae)	Cotton seed Oil	500	72.00	20.00
		1000	67.00	25.56
<i>Cymbopogon citratus</i> (Poaceae)	Lemon Grass Oil	500	0.00	100.00
		1000	0.00	100.00
<i>Gaultheria procumbens</i> (Ericaceae)	Winter Green Oil	500	26.50	70.56
		1000	0.00	100.00
<i>Pogostemon cablin</i> (Lamiaceae)	Patchauli Oil	500	76.00	15.56
		1000	59.60	33.78
		1500	36.33	59.63
		2000	13.67	84.81
<i>Pelargonium graveolens</i> (Geraniaceae)	Geranium Oil	500	77.00	14.44
		1000	53.60	40.44
		1500	0.00	100.00
		2000	12.00	86.67
<i>Eucalyptus globulus</i> (Myrtaceae)	Eucalyptus Oil	500	57.60	36.00
		1000	44.20	50.89
		1500	36.67	59.26
		2000	21.00	76.67
<i>Mentha piperita</i> (Lamiaceae)	Peppermint Oil	500	0.00	100.00
		1000	0.00	100.00
<i>Azadirachta indica</i> (Meliaceae)	Neem Oil	500	55.00	38.89
		1000	48.60	46.00
		1500	29.00	67.78
		2000	71.00	21.11
Bavistin	-	0.02%	0.00	100.00
Mancozeb	-	0.25%	0.00	100.00
Bavistin+mancozeb (B+M)	-	(1:1)	0.00	100.00
Control	-	-	90.00	0.00
LSD (5%)		500	1.9	-
		1000	2.8	-

The average radial growth of pathogen on eight day of incubation and their percent inhibition with various essential oils are mentioned in Table 1. On the basis of percent inhibition of the fungi at different concentrations of the essential oils, the degree of fungi toxicity was clustered in three broad group viz; high active (71-100%). active (51-70%) moderately active (31-50%) and least active (0-30%).

We started the dose dependent bio-assay with 500 & 1000 PPM and found oils of Lemongrass and peppermint were giving complete inhibition on 500 PPM than we proceeded further with lower concentrations (100-500) and found both the oils were exhibiting highest efficacy as that of reference fungicides (Fig.1). Zambonelli *et al.* also found *Mentha piperata* fungicidal against test fungus. Other oils with at least moderately activity on 1000 PPM were selected and screened with higher concentrations (1000, 1500 & 2000) and rest with least activity were rejected for screening. Oils of winter green and geranium both ranked in first category with complete inhibition at 1000 & 1500 PPM.




Fig.1 Percent Inhibition of oils of Lemon grass and Pepper mint

In poaceae family only lemon grass oil is effective with highest fungitoxicity while oils of palma rosa & citronella belonged to fourth category respectively with least inhibition of 26 & 25.11%. At 2000 PPM oils of patchouli & eucalyptus showed highest category with percent inhibition of 84.41&76.67 respectively.

Eucalyptus species not only provide fuel mass and reduce atmospheric carbon dioxide levels directly but also used as pesticidal agents. Eucalyptus oil showed its inhibitory action against *C. lindemuthianum*. In fact eucalyptus oil has been known for hundreds of years as antibacterial, antifungal and antiseptic in nature<sup>16</sup>. Whilst eucalyptus oils have a sturdy toxicity in the vapor form against a wide range of microbes and insects, and they could be commercially exploited as a fumigant for stored products.

Neem, basil and geranium oils after getting their maximum inhibitory action at specific concentration, whereas accelerated the growth of fungal mycelium too. We observed the rest of oils were least fungitoxic with lowest percent inhibition against test fungi.

### ***LD50 determination***

The purpose of estimating different concentrations is to determine the antifungal activity with its LD50 value which is sufficient to kill fungus and to use the estimate to make comparisons. It is generally easiest to estimate the median (50%) response level of the population. Lethal dose (LD50) of the test essential oils were determined by using GW-Bascis LD50 programme.

**Table No. 2. LD 50 value of promising essential oils**

<b>Essential oil</b>	<b>LD 50(ug/ml)</b>
<i>Cymbopogon citratus</i>	380.27
<i>Gaultheria procumbens</i>	1157.85
<i>Pogostemon cablin</i>	951.23
<i>Pelargonium graveolens</i>	951.23
<i>Eucalyptus globulus</i>	438.72
<i>Mentha piperita</i>	110.61
<i>Azadirachta indica</i>	292.23

## CONCLUSION

Present investigation revealed that oils of lemongrass, peppermint, along with wintergreen and geranium were found to be most fungicidal with complete inhibition but lemongrass & peppermint were considered to be more effective and promising to be continue with for *in-vivo* study because they gave complete inhibition at lowest concentration as compared to others. Implications caused by fungi mean there is a constant striving to produce safer food crops and to develop new antifungal agents. It is clear from above discussion that essential oils posses a wide spectrum of antifungal activity and provide a simple, inexpensive, and environmental friendly alternative pest control. The oil of lemon grass and pepper mint also be exploited for seed treatment and under field conditions in ensuring effective disease management. These are the findings with possible potential application in augmenting the crop yields.

## REFERENCES

1. Neergaard, P. Seed Pathology. McMillan Press, London and Basingskote. 1977; 839.
2. Richardson, MJ. An annotated list of seed-borne diseases. Fourth edition, ISTA, Zurich, Switzerland.1990; 387.
3. CAB International. Crop protection compendium. CAB International, Wallingford: UK. 2007.
4. Darokar MP, Mathur A, Dwivedi S, Bhalla R, Khanuja SPS and Sushil K. Detection of antibacterial activity in the floral petals of some higher plants. Current Science. 1998; 75 (3): 187-189.
5. Dev U, Devkumar C, Agarwal PC, Suneja P and Mohan J. Antifungal essential oils: An eco-friendly approach for salvaging germplasm. Indian Journal of Plant Genetic Resources. 2005; 18(1): 144.
6. Yahyazadeh M, Omidbaigi R, Zare R and Taheri H. Effect of some essential oils on mycelial growth of *Penicillium digitatum* Sacc. World Journal of Microbiology and Biotechnology. 2008; 24(8): 1445-1450.
7. Lee Yeon Suk, Kim Jun Heon, Shin Sang Chul, Lee Sang Gil and Park IlK won. Antifungal activity of Myrtaceae essential oils and their components against three phytopathogenic fungi. Flavour and Fragrance Journal.2008; 23(1): 23-28
8. Yang VW and Clausen CA. Antifungal effect of essential oils on southern yellow pine. International Biodeterioration and Biodegradation.2007; 59(4): 302-306

9. Zambonelli A, D'Aulerio AZ, Bianchi A and Albasini A. Effects of Essential Oils on Phytopathogenic Fungi *In Vitro*. Journal of Phytopathology, 1996; 144: 491–494.
  10. Ranasinghe L, Jayawardena B and Abeywickrama K. Fungicidal activity of essential oils of *Cinnamomum zeylanicum* (L.) and *Syzygium aromaticum* (L.) Merr et L.M.Perry against crown rot and anthracnose pathogens isolated from banana. Letters in Applied Microbiology. 2002; 35: 208–211.
  11. Tzortzakis NG and Economakis CD. Antifungal activity of lemongrass (*Cymbopogon citratus* L.) essential oil against key postharvest pathogens. Innovative Food Science and Emerging Technologies. 2007; 253-258
  12. Cosic J, Vrandecic K, Postic J, Jurkovic D and Ravlic M. In vitro antifungal activity of essential Oils on growth of phytopathogenic fungi. Poljoprivreda.2010; 16(2), 25-28
  13. Maqbool M, Alderson AA, Mahmud Tengku PG, Siddiqui MMY and Zahid N. Postharvest application of gum arabic and essential oils for controlling anthracnose and quality of banana and papaya during cold storage. Postharvest Biology and Technology. 2011; 62(10): 71-76.
  14. Kulshrestha DD, Mathur SB and Neergaard P. Identification of seed borne species of *Colletotrichum*. Friesia.1976; 11, 116-125.
  15. Nene YL and Thapliyal PN. Fungicides in plant Disease Control. Universal Oxford and UYF: Calcutta: 1979.
  16. Brooker MIH and Kleinig DA. Field guide to Eucalyptus. South-eastern Australia. Bloomings, Melbourne.2006; 3(1).
-