

International Journal of Scientific Research and Reviews

Influence of Bio and Nano Zinc Fertilization on Growth and Yield of Rice

Ponnmani .P, T. Muthukumararaja* and M.V. Sriramachandrasekharan

Department of Soil Science and Agricultural Chemistry, Faculty of Agriculture,
Annamalai University, Annamalai Nagar- 608002 Email:saradhakirankumar@gmail.com

ABSTRACT

Zinc is the essential plant micronutrients and their importance for crop productivity is similar to that of major nutrients. The field experiment was conducted in zinc deficient soil belonging to Padugai series (VerticUstrophept) at the farmer's holding during the Navarai season of year 2017. The treatments consists of viz., T₁ - Absolute control , T₂ - NPK (RDF), T₃ - RDF + ZnSO₄ @ 25 kg ha⁻¹ , T₄ - RDF + Nano zinc (Granules @ 10 kg ha⁻¹), T₅ - RDF + Nano zinc (Granules @ 15 kg ha⁻¹), T₆ - RDF + Bio zinc (Granules @ 10 kg ha⁻¹), T₇ - RDF + Bio zinc (Granules @ 15 kg ha⁻¹), T₈ - RDF + Foliar spray of 0.5 % as ZnSO₄ at T.S and P.I., T₉-RDF + Foliar spray of 1 ml L⁻¹ as nano zinc at T.S and P.I. and T₁₀- RDF + Foliar spray of 1.5 ml L⁻¹ as bio zinc at T.S and P.I.. The test crop was rice var.CO 51. The results revealed that grain and straw yield was significantly enhanced on addition of different sources of zinc over control. The grain and straw yield was maximum (6070 and 7068 kg ha⁻¹) with application of RDF+ soil application of bio zinc @ 30 kg ha⁻¹ and was on par with RDF + Foliar spray of 0.5 % ZnSO₄ (5983 and 6980 kg ha⁻¹). Similarly, the plant height and DMP was recorded with application of RDF+ soil application of bio zinc @ 30 kg ha⁻¹.

KEY WORD: Zinc, organics, rice, yield, Plant height, DMP.

***Corresponding author:**

T. Muthukumararaja

Department of Soil Science and Agricultural Chemistry,
Faculty of Agriculture,
Annamalai University,
Annamalai Nagar- 608002

INTRODUCTION

Micronutrient deficiencies such as limited zinc (Zn) availability are one of the main problems limiting agricultural productivity, especially in alkaline calcareous soils. Therefore, Zn is often included in macronutrient fertilizers to improve crop quality and productivity¹Zinc is an essential element that plays many important roles in various physiological and metabolic processes in plant. This trace element plays vital function in structural molecules such as DNA and activates different metabolic and regulatory enzymes. It has been reported that nearly 925 proteins in humans and over 500 proteins in plant contain zinc.²It also plays a role in photosynthesis, protein synthesis, cell division, maintaining integrity of the membrane structure³, resistance against the pathogen infection⁴ and add sexual reproduction through affecting production and shape of pollen and changes in the stigma. Plants also may develop symptoms such as interveinal chlorosis, bronzing of leaves, abnormally shaped leaves, stunting or resetting which can effectively reduce crop production and health⁵.According to the World Health Organization, the average prevalence of zinc deficiency in the world population is 31 per cent which may range from 4 per cent to 73 per cent in different countries⁶. Agricultural systems are the main pathway from which nutrients including zinc enter the human food chain. Therefore, zinc malnutrition must be directly dependant on the inability of cropping systems to deliver enough zinc to the food crops⁷.

In India, zinc is considered as the fourth important yield limiting nutrient after nitrogen, phosphorus and potassium respectively. Zinc deficiency affects one third of the world's population. In India, 47 per cent of the soils are zinc deficient. Critical limit of a nutrient in soils refers to a level below which the crops will readily respond to its application. This critical limit varies with soil, crops and varieties. Critical limit of zinc for rice was 0.74 ± 0.18 ppm across the soils and indifferent agro-ecological regions of India⁸. The analysis of DTPA extractable zinc in soils has shown the 40 per cent of soil samples were potentially zinc deficient⁹. It has been postulated that the zinc deficiency is likely to increase from 49- 63 per cent by the year 2025 as most of the marginal soils brought under cultivation are showing the symptoms of zinc deficiency¹⁰.Rice (*Oryza sativa* L.) is an important stable food crop among all the cereals.About 90 per cent of rice grown and consumed in South and South East Asia. In some parts of the world consumption of rice is a high as 990 g per person per day¹¹. India ranks first in the world in terms of area of rice cultivation with 44.6 m ha and second in productivity of 2.96 t ha^{-1} . In Tamil Nadu has produced 79.14 lakh tones of rice from an 18.3 lakh hectares in 2014 -2015¹². The efficiency of applied ZnSO_4 is only 1 to 4% and most of the applied zinc is rendered unavailable to plants due to many factors such as leaching, fixation¹³. Hence it is essential to minimize the nutrient losses in fertilizer application, increase the crop yield through the exploitation of new applications with the help of nano technology and nano materials.The overview

of green synthesis is easily understood when the precursor is added to the leaf extract it changes the color which indicates the formation of nano catalyst^{14,15}. Literally very little information exist on the application of nano zinc both in chemical and bioforms applied to soil and foliar application of these materials on rice crop under field conditions. With this background the present experiment entitled influence of bio and nano zinc fertilization on growth and yield of rice.

MATERIALS AND METHODS

With a view to study the response of growth and yield of rice to bio and nano zinc in deficient soil, The field experiment was conducted in zinc deficient soil belonging to Padugai series (Typic Ustifluvents) at the farmer's holding during the Navarai season of year 2017. The experimental soil was clay loam in texture with pH 7.78, EC 0.84 dS m⁻¹, organic carbon 3.9 g kg⁻¹ (low), low in KMnO₄-N 275 kg ha⁻¹, low in Olsen-P 10.4 kg ha⁻¹, high in NH₄OAc-K 294 kg ha⁻¹ and low in available DTPA-Zn 0.68 mg kg⁻¹. The experiment was laid out in randomized block design. The treatments consists of eight treatments viz., T₁ - Absolute control , T₂ - NPK (RDF), T₃ - RDF + ZnSO₄ @ 25 kg ha⁻¹ , T₄ - RDF + Nano zinc (Granules @ 10 kg ha⁻¹), T₅ - RDF + Nano zinc (Granules @ 15 kg ha⁻¹), T₆ - RDF + Bio zinc (Granules @ 10 kg ha⁻¹), T₇ - RDF + Bio zinc (Granules @ 15 kg ha⁻¹), T₈ - RDF + Foliar spray of 0.5 % as ZnSO₄ at T.S and P.I., T₉ - RDF + Foliar spray of 1 ml L⁻¹ as nano zinc at T.S and P.I. and T₁₀ - RDF + Foliar spray of 1.5 ml L⁻¹ as bio zinc at T.S and P.I. The recommended dose of 150:50:50 N, P₂O₅, K₂O ha⁻¹ through urea, superphosphate and muriate of potash was added uniformly to all the plots. Nitrogen was applied in three split doses i.e., 50% as basal, 25% each at active tillering and 25% panicle initiation stages. The entire dose of P₂O₅ and K₂O were applied basally as per the treatment schedule. The test crop rice CO 51. The zinc was applied through bio and nano zinc formulations. At different stages of crop growth plant height and DMP and harvest stage grain and straw yield were recorded.

RESULTS AND DISCUSSION

Plant height

Analysis of variance on plant height furnished in (Table 1) showed that application of sources and methods caused significant improvement on plant height at all stages over control. Plant height ranged from 36.5 to 56.8 cm at tillering stage, 68.4 to 89.4 cm at panicle initiation stage and 88.1 to 107.2 cm at harvest stage respectively. Plant height at all stage was highest with RDF + soil application of bio zinc @ 30 kg ha⁻¹ compared to rest of the treatments. Among the treatments, tallest plant (56.8, 89.4 and 107.2 cm) at tillering, panicle initiation and harvest stage was noticed with application of RDF + soil application of bio zinc @ 30 kg ha⁻¹ (T₇) and was on par with T₈, T₉ and significantly followed by T₃. At harvest stage application of RDF + soil application of bio zinc @ 30 kg

ha⁻¹ produced 17 % of taller plants as compared to control. Compared to all the treatments the lowest plant height was recorded in the treatment receiving RDF @ 120: 50: 50 kg ha⁻¹ (101.2 cm) followed by control (88.1 cm) at harvest stage. This could be attributed to adequate supply of zinc which accelerates the activity of enzymes and auxin metabolism in plants. It could also be attributed to the increased growth of internodal portion with higher synthesis of IAA and metabolizing gibberellic acid¹⁶. Zinc is known for precursor of growth as they play greater and crucial role during early phase of the plant life. Zinc is precursor of tryptophan, which is supposed to play a decisive role in the synthesis of auxin, the main factor behind apical dominance, growth and development, which is involved in cell division and internode elongation¹⁷. The result obtained in the present study was confirmed by the significant positive correlation between plant height with DTPA-Zn ($r=0.972^{**}$), zinc concentration ($r=0.992^{**}$) and Zn uptake ($r=0.987^{**}$).

Dry matter production (DMP)

Addition of zinc sources and methods caused significant effect on dry matter production at all stages of crop growth over NPK alone (Table 2). DMP at all stage was highest with RDF + soil application of bio zinc @ 30kg ha⁻¹ compared to rest of the treatments. The DMP ranged from 1708 to 2929 kg ha⁻¹ at tillering stage and 4061 to 5802kg ha⁻¹ at panicle initiation stage respectively. Among the treatments, the treatment receiving RDF + soil application of bio zinc @ 30kg ha⁻¹ recorded significantly highest dry matter production at tillering stage (2929 kg ha⁻¹) and panicle initiation stage (5802 kg ha⁻¹) respectively. This was followed by RDF + foliar spray of 0.5 % Zn SO₄ (T₈), RDF + foliar spray of 0.1% nano zinc (T₉) which were on par with each other. These treatments were followed by RDF + soil application of Zn SO₄ @ 25 kg ha⁻¹. The lowest dry matter was recorded in the treatment absolute control and was followed by the treatment receiving RDF @ 150: 50: 50 kg ha⁻¹. Increase in DMP due to zinc might be due to role of zinc for the activities of various types of enzymes such as those required for CO₂ assimilation pathway and chlorophyll biosynthesis¹⁸. This was confirmed by significant positive correlation between DMP with DTPA-Zn ($r=0.90^{**}$; $r=0.87^{**}$) and Zn uptake ($r=0.94^{**}$; $r=0.95^{**}$) at tillering and panicle initiation stages respectively. Increase in DMP due to zinc was reported by earlier workers^{19,20,21}.

Rice yield

A significant increase in grain and straw yield of rice was noticed due to application of various sources and methods of zinc application over control (Table 3). Grain and straw yield ranged from 3319 to 6070 and 4032 to 7068 kg ha⁻¹. The highest grain and straw yield was obtained with RDF + soil application of bio zinc @ 30kg ha⁻¹ (6070 and 7068 kg ha⁻¹) and was on par with RDF + foliar spray of 0.5 % ZnSO₄ (5522 kg ha⁻¹), T₉. These treatments were significantly followed by T₃, T₁₀ and

T₆. The percentage increase in grain and straw yield (45 and 42 %) was noticed with RDF + soil application of bio zinc @ 30kg ha⁻¹(T₇) compared to over control (T₁).The lowest grain and straw yield was recorded in the treatment receiving RDF @150: 50: 50 kg ha⁻¹ followed by control(3319 and 4032 kg ha⁻¹).Increase in grain yield due to zinc was the logical result due to increase in yield components. In the present study, number of panicles m⁻², number of grains panicle⁻¹, panicle length and 1000 grain weight increased with Zn levels and highest value was obtained with RDF + soil application of bio zinc @ 30kg ha⁻¹.The above argument was ably supported by linear relationship between grain yield with number of panicles m⁻² ($Y = 3628 - 0.703x + 0.010x^2$, x^2 , $R^2 = 99^{**}$), numberof grains panicle⁻¹ ($Y = 4688- 83.49 x - 9.322 x^2$, $R^2=99^{**}$), panicle length ($Y = -353.6 + 393.6x - 4.473 X^2$, $R^2= 0.99^{**}$) which showed that 99 per cent variation in grain yield are brought out by different yield attributes. ^{22,23}reported increase in grain yield due to improvement in yield components.

CONCLUSION

In conclusion, Bio zinc application to zinc deficient soil significantly increased plant height, DMP and grain yield of rice. Application of Bio zinc proved better over application of alone zinc sulphate (ZnSO₄) indicating the positive role of organic matter in increasing plant height, DMP and grain yield on soils affected with zinc deficient.

Table 1. Effect of zinc sources and methods on plant height (cm) at different growth stage.

Treatments	Tillering stage	Panicle initiation stage	Harvest stage
T ₁ -Absolute control	36.5	68.4	88.1
T ₂ - NPK (RDF)	41.5	74.0	93.0
T ₃ - RDF + Zn SO ₄ (25 kg/ha)	50.10	83.6	101.4
T ₄ -RDF + Nano Zinc (granules @ 10 kg /ha)	43.1	75.6	94.4
T ₅ -RDF + Nano Zinc (granules @ 15 kg /ha)	44.5	76.8	95.7
T ₆ - RDF + Bio zinc (granules @ 15 kg/ha)	48.0	80.5	98.8
T ₇ - RDF + Biozinc (granules @ 30 kg/ha)	56.8	89.4	107.2
T ₈ - RDF + Foliar spray of 0.5% Zn SO ₄ @ TS and PIS	55.1	88.7	106.4
T ₉ - RDF + Foliar spray of 0.1% Nano Zinc @ TS and PIS	53.5	87.6	104.8
T ₁₀ - RDF + Foliar spray of 0.15% Bio Zinc @ TS and PIS	49.2	82.1	100.3
SEd	1.25	1.09	1.28
CD (P = 0.05)	2.64	2.29	2.70

Table 2. Effect of zincsources and methods on DMP (kg ha⁻¹) at different growth stages

Treatments	Tillering stage	Panicle initiation stage
T ₁ – Absolute control	1708	4061
T ₂ – NPK (RDF)	1937	4600
T ₃ – RDF + Zn SO ₄ (25 kg/ha)	2487	5304
T ₄ –RDF + Nano Zinc (granules @ 10 kg /ha)	2034	4800
T ₅ –RDF + Nano Zinc (granules @ 15 kg /ha)	2122	4891
T ₆ – RDF + Bio zinc (granules @ 15 kg/ha)	2323	5206
T ₇ – RDF + Biozinc (granules @ 30 kg/ha)	2929	5802
T ₈ – RDF + Foliar spray of 0.5% Zn SO ₄ @ TS and PIS	2854	5706
T ₉ - RDF + Foliar spray of 0.1% Nano Zinc @ TS and PIS	2765	5601
T ₁₀ - RDF + Foliar spray of 0.15% Bio Zinc @ TS and PIS	2412	5303
SEd	55.42	73.35
CD (P = 0.05)	116.43	154.11

Table 3. Effect of zinc sources and methods on grain and straw yield of rice (kg ha⁻¹)

Treatments	Grain Yield (kg ha ⁻¹)	Straw Yield (kg ha ⁻¹)
T ₁ – Absolute control	3319	4032
T ₂ – NPK (RDF)	4483	5365
T ₃ – RDF + Zn SO ₄ (25 kg/ha)	5522	6572
T ₄ –RDF + Nano Zinc (granules @ 10 kg /ha)	4657	5551
T ₅ –RDF + Nano Zinc (granules @ 15 kg /ha)	4758	5708
T ₆ – RDF + Bio zinc (granules @ 15 kg/ha)	5338	6377
T ₇ – RDF + Biozinc (granules @ 30 kg/ha)	6070	7068
T ₈ – RDF + Foliar spray of 0.5% Zn SO ₄ @ TS and PIS	5983	6980
T ₉ - RDF + Foliar spray of 0.1% Nano Zinc @ TS and PIS	5962	6909
T ₁₀ - RDF + Foliar spray of 0.15% Bio Zinc @ TS and PIS	5455	6459
SEd	131.17	106.23
CD (P = 0.05)	281.88	223.19

REFERENCE

1. Khalivand Behrouzfar E, M .Yarnia , FR . khoii, M .Mogaddam and M.N. Safarzadehvishkaii. The effect of methanol and some micro – macronutrients foliar application on Maize (*Zea mays* L.,) maternal plant on some of morph physiological characteristics in a subsequent generation. International journal of Agronomy and plant production. 2012; 3 (12): 618 -624.
2. Graham, R.D. Micronutrient deficiencies in crops and their global significance. In : micronutrient deficiencies in global crop production(Ed Alloway BJ). Springer. 2008; 41 -61.
3. Marschner, H. Mineral nutrition of higher plants. 2nd ed. Academic press pub., New York (USA), 1995; 559.
4. Sarwar, M. Effects of zinc fertilizers application on the incidence of rice stem borers (*Scirpophagasp.*) (Lepidoptera: Pyralidae) in rice (*Oryza sativa*) crop. **J. Cereals Oilseeds**, 2011;2(5): 61-65.

5. Alloway, B.J. Soils factors associated with zinc deficiency in crops and humans. **Environ. Geochem. Hlth.**, 2009; **31**(5): 537-548.
6. Caulfield L.E and R.E .Black . Zinc deficiency, In M. Ezzati, et al., eds. Comparative Quantification of Health Risks: Global and Regional Burden of Disease Attributable to Selected major Risk Factors. 2004. World Health Organization, Geneva.
7. Welch R.M.. Linkages between trace elements in food crops and human health. 2008; 41-61. In B.J. Alloway, ed. Micronutrient Deficiencies in Global crop Production. Springer Science.
8. Muthukumararaja, T. and M.V. Sriramachandrasekharan. Critical limit of zinc for rice soils of Veeranam command area, Tamilnadu, India. *ARPN J. Agrl. Biol. Sci.* 2012; **7**(1): 1-12.
9. Shukla, A.K., Tiwari, P.K and Chandra Prakash. Micronutrients deficiencies vis-a- vis food and nutritional security of India. *Indian J. Fertilizers.* 2014; **10**(12): 94-112.
10. Arunachalam, P., Kannan, P., Prabukumar, G and Govindaraj, M. Zinc deficiency in Indian soils with special focus to enrich zinc in peanut. *African J. Agricultural research.* 2013;**8**(50): 6681 -6688.
11. Sharma, B.D., Arora, H., Kumar, R and Nayyar, V.K.. Relationships between soil characteristics and total DTPA –extractable Zn, Cu, Mn and Fe in Indo-Gangetic plains. *Communication in soil science and plant analysis.* 2015;**35**: 799 -818.
12. Vaithilingam, R. 2015. Policy Note – Department Of Agriculture, Government Of Tamil Nadu
13. Nair, R., S. H. Varghese, B. G. Nair, T. Maekawa, Y. Yoshida and D. S. Kumar. Nanoparticulate material delivery to plants. **J. Plant Sci.** 2010; **179**, 154:163.
14. Yuvaraj, M and Subramanian, K.S. Fabrication of zinc nano fertilizer on growth parameter of rice. *Trends in biosciences.* 2014;**7**(17): 2564-2565.
15. Kandasamy, S and Prema, R.S. Methods of synthesis of nano particles and its applications . *Journal of Chemical and Pharmaceutical research.* 2015; **7**(3): 278-285
16. Flaig, W., N. Beutolspacher and E. Rietz. Chemical composition and physical chemical properties of humic substances. In: Soil components (J.E. Gieseking, ed.), Springer-Verlag, Berlin, New York 1975; 1.
17. Shanmugam, P.M. and R.Veeraputhiran. Effect of organic and inorganic N and zinc application on soil fertility and nutrient uptake of rabi rice. **Madras Agric. J.**, 2000; **89**(1-3): 90- 92.
18. Devarajan, R. and G. Ramanathan. Direct, residual and cumulative effect of applied zinc for rice in red soil. **Madras Agric. J.**, 1995; **82**(2): 102-103.
19. Obrador, A., J. Novillo and J.M. Alvarez. Mobility and availability to plants of two sources zinc applied for a calcareous soil. **Soil Sci. Soc. Amer. J.**, 2003;**67**: 564-572.

20. Shehu, H.E. and G.Y. Jamala. Available Zn distribution, response and uptake of rice (*Oryza sativa* L.) to applied Zn along a Toposequence of Lake Gerio Fadama soils at Yola, North Eastern Nigeria. **J. Amer. Sci.**, 2010; **6**(11): 1013-1016.
21. Fageria, N.K., A.B. Das Santos and T. Cobucci. Zinc nutrition of lowland rice. **Commun. Soil sci. Plant Ann.** 2011;**42**(4): 1719-1727.
22. Abbas, G., G. Hassan, M.A., Ali, M. Aslam and Z. Abbas.. Response of wheat to different doses of ZnSO₄ under Thal desert environment. **Pak. J. Bot.**, 2010; **42**(6): 4079-4085.
23. Rahman, K.M.H., Md. Abdul Khan Chowdhury, F. Sharmeen and A. Sarkar. Effect of zinc and phosphorus on yield of *Oryza sativa* (cv. BR 11). **Bang. Res. Pub. J.**, 2011; **5**(4): 351-358.