

## *International Journal of Scientific Research and Reviews*

### **Immobilization of Partially Purified Fungal Dye Degrading Laccase Enzyme**

**S. Sudhapriyadharsini\* and N. Hema Shenpagam**

\*PG and Research Department of Microbiology, Hindusthan College of Arts and Science  
Coimbatore, Tamilnadu, [Indai.shareatsudha@gmail.com](mailto:Indai.shareatsudha@gmail.com), Ph.8637698978.

Associate professor, PG and Research Department of Microbiology,  
Hindusthan College of Arts and Science, Bharathiyar University, Coimbatore, Tamilnadu, India.  
[drhemashenpagam@gmail.com](mailto:drhemashenpagam@gmail.com), Ph 9443672644.

#### **ABSTRACT**

In the modern world many industries different kinds of polluting agents are spoiling our mother nature. 100000 dyes have been generated worldwide with an annual production of over  $7 \times 10^5$  metric tons. Laccase is a copper-containing polyphenol oxidase that acts on a wide range of substrates. This enzyme is found in many plant species and is widely distributed in fungi. Because of its importance in bioremediation, potential dye degrading fungal cultures were screened for laccase enzyme production by plate test method using the indicator compound crystal violet. Alginate-gelatin mixed gel was applied to immobilized laccase for decolonization of some industrial dyes including crystal violet. The immobilization procedure was accomplished by adding alginate to a gelatin solution containing the enzyme and the subsequent dropwise addition of the mixture into a stirred  $\text{CaCl}_2$  solution. The obtained data showed that both immobilized and free enzymes acted optimally at  $45^\circ\text{C}$  for removal of industrial dye, but the entrapped enzyme showed higher thermal stability compared to the free enzyme. The immobilized enzyme represented optimum decolorization at pH 7. Reusability of the entrapped laccase was also studied and the results showed that 71% activity was retained after five successive cycles. The best removal condition was applied for decolorization of several other industrial dyes. Results showed that the maximum and minimum dye removal was related to gentian violet and direct black 19, respectively. Potential fungal dye degrading laccase enzyme is as an important source for bioremediation of toxic dye removal.

**KEYWORDS:** Industrial dye, *Aspergillus* spp, immobilized enzyme bioremediation.

#### **\*Corresponding author**

**S. Sudhapriyadharsini**

Ph. D Research scholar

PG and Research Department of Microbiology

Hindusthan college of Arts and Science

Coimbatore, Tamil Nadu, INDIA.-641028

Email: [shareatsudha@gmail.com](mailto:shareatsudha@gmail.com), Ph. 8637698978

## **INTRODUCTION**

Dyes are released in to the environmental through industrial effluents from three major sources such as textiles, dyestuff manufacturing and paper industrial<sup>1</sup>. Green technologies to deal with this problem include adsorption of dyestuffs on bacterial and fungal biomass<sup>2, 3</sup> or low-cost non-conventional adsorbents<sup>4, 5</sup>. In the recent years, enzymes have gained great importance in Industries; Laccases are one among them which are widely present in the nature. Laccases are the oldest and most studied enzymatic systems. These enzymes contain 15–30% carbohydrate and have a molecule mass of 60–90kDa<sup>6</sup>. Laccases (EC 1.10.3.2; benzenediol: oxygen oxidoreductases) are multicopper enzymes belonging to the group of blue oxidases that catalyses oxidation of a wide variety of organic and inorganic compounds including diphenols, polyphenols, diamines, and aromatic amines. One electron at a time is removed from the substrate and molecular oxygen is used as the electron acceptor<sup>7</sup>. Enzymes action at mild conditions of pH, pressure and temperature proposes them as candidates for suitable catalysts in industries where low cost, energy savings and simplicity are important<sup>8, 9</sup>. Enzyme entrapment uses natural and synthetic polymers such as agarose, agar and gelatin through thermo reversal polymerization alginate, polyvinyl acetate, acrylic acid and  $\beta$ -carrageenan by ionotropic gelation<sup>10</sup>. Laccases are an interesting group of multicopper enzymes produced by higher plants and fungi that catalyze the oxidation of a wide range of organic compounds such as phenols in the presence of molecular oxygen<sup>11, 12</sup>. The most important applications discussed for laccases include pharmaceutical and food industries, textile effluent transformation and wastewater detoxification biosensors. The unique properties of laccases such as high stability in solution, mild reaction conditions and selectivity for phenolic structure make them attractive for use in chemical synthesis<sup>13</sup>. Alginategelatinmixed gel to immobilize laccase which is further employed in the decolorization of some synthetic dyes, such as crystal violet in aqueous solutions. Optimum pH, temperature, and proper enzyme content for decolorization by the immobilized laccase were also studied. To our knowledge, the use of alginate-gelatin mixed gel has not been examined previously for the purpose of enzyme entrapment<sup>14, 15</sup>. The aim of the present paper is to screen and isolate potential dye degrading fungi from dye accumulated soil and water samples and identifying a new source of extracellular laccase. The extracellular fungal laccase in the culture medium was subjected to purification, characterization and immobilization studies.

## **MATERIALS & METHODS**

All other reagents and chemicals were of the highest purity available. Various industrial dye accumulated soil and water samples were collected for isolate a potential dye degrading microorganisms (fungi). Different types of industrial dyes are collected from the industry from 4

different places in Thirupur {Tamil nadu,India} in 2 summer and winter climatic conditions. Isolation of dye decolourizing fungal strain was selected and further Production of Extracellular laccase enzyme was partially purified from the following of Mikiashvili et al., (2006) <sup>16</sup> procedure was applied for the potential dye degrading fungal isolate.

#### **A. *Enzymatic plate assay***

Laccase activity was determined by the oxidation of ABTS method <sup>17</sup>. The laccase plate assay allowed rapid determination of the presence of laccase in the extracellular fluid. 15mL of sterile agarose (0.5%) medium containing 0.5mM of ABTS per mL in sodium acetate buffer (pH 4.5, 0.1M) was placed on a petriplate (5 × 5 cm). The development of an intense bluish-green color around the wells was considered as a positive test for laccase activity <sup>18</sup>.

#### **B. *Preparation of mixed gels, immobilization of laccase, and decolorization of crystal violet***

The gelatin-alginate mixed gel was prepared according to the method of Panouille and Larreta-Garde <sup>19</sup>, with some modifications. Briefly, entrapment of laccase in gelatin-alginate mixed gel was performed by adding 0.1 g sodium alginate to 10mL of a solution containing 10% (w/v) gelatin and laccase in the range of 5–50mg (1mg enzyme is equal to 1.25 U) under continuous stirring at room temperature. The mixture was syringed into a stirring CaCl<sub>2</sub> solution (200 mM), and the resulting beads were left to be hardened for 1 h under the same conditions, washed three times by deionized water, and then stored at 4°C prior to being used in the decolorization study. The amount of bound protein was determined by Lowry's method.

#### **C. *Optimum temperature, pH, and enzyme concentration of the immobilized laccase***

The effect of pH on the enzymatic decolorization was monitored with a dye concentration of 0.25mg/ml at a 3–9 pH range adjusted by citrate or ammonia buffers. To determine the effect of temperature on the enzymatic decolorization, the reaction mixture was incubated at a thermal range of 30–70°C, in steps of 10°C. In order to assess the effect of enzyme quantity on decolorization, the reaction was started with different enzyme amounts, from 0.25 to 0.5mg/ml, in 0.05mg/ml increments. The experiments were performed in triplicate and the results shown are means standard deviation.

#### **D. Reusability**

Reusability of the gelatin-alginate immobilized enzyme was investigated in acetate buffer solution 0.1M at pH 8 and 45°C. The used beads were filtered at the end of each cycle and washed three times with the same buffer to treat in the next fresh colored solution. The reusability study was performed in triplicate.

Relative Decolorization (%) =  $(A_{\text{initial}} - A_{\text{final}}) / A_{\text{initial}} \times 100$ , where  $A_{\text{initial}}$  was the initial absorbance and  $A_{\text{observed}}$  was the final absorbance at the given wavelength.

### **RESULT AND DISCUSSION**

The potential dye degrading fungal (*Aspergillus* spp) isolate to produce a partially purified laccase enzyme was applied for further analysis of ABTS plate assay and immobilized laccase was optimized.

#### **A. Enzymatic Plate Assay**

The laccase plate assay showed the presence of laccase activity in the partially purified sample. The heat-treated sample showed less intense green color than that of the partially purified sample as shown in (Fig. I). The development of an intense bluish-green color zone the wells was considered as a positive test for laccase activity




**Figure: I –ABTS plate assay for demonstration of laccase activity in potential fungal sample were compared with control and test petriplates.**

#### **B. Preparation of mixed gels, immobilization of laccase, and decolorization of crystal violet**

Spherical and regular-shaped gelatin-alginate beads were obtained by using ionotropic gelation; the diameters of the beads were 2.0–5mm (Fig. II). The results indicated the rupture force increase with the rise in alginate concentration. The prepared beads containing 5% alginate exhibited

the highest mechanical stability. Alginate could provide a very strong network that required 0.980kgf (kilogram-force) forces to rupture and the bound protein level 0.149  $\mu\text{g/ml}$ .



**Figure: II – Preparation of Mixed Gels, Immobilization of Laccase enzyme**

### ***C. Optimum temperature, pH, and enzyme Concentration of decolorization of industrial dyes of the immobilized laccase enzyme***

Among this analysis the effect of temperature on dye decolorization was investigated by changing the reaction mixture temperatures in a 30–70°C range. The results obviously showed that decolorization increased as the temperature increased up to 45°C; at higher temperatures, decolorization efficiency remarkably decreased, which normally attributed to denaturation of enzyme. However, the comparison of decolorization results indicated high thermal stability of immobilized enzyme and confirmed the protective role of alginate-gelatin mixed gel for maintaining enzyme activity. At 45°C, the crude and immobilized enzymes exhibited more activity. Similarly, Forootanfar and colleagues (2011) reported that the optimum temperature for laccase activity was 45°C. Other research studies have reported an optimum temperature range of 45–50°C for (Table.I)laccase activity immobilized by both covalent binding and adsorption<sup>19,20</sup>.

Extremely high or low pH values generally result in complete loss of activity for most enzymes. pH is also a factor in the stability of enzymes. As with activity for each enzyme there is also a region of pH optimal stability. pH plays the key role in dye decolorization. The effect of pH on dye decolorization was examined at pH ranging from 3 to 9, using buffers. The free enzyme significantly displayed higher decolorization activity in pH 5 (table. II). Immobilized enzyme beads are sharp increase at pH values above 7 to 9. The optimal pH for the immobilized enzyme activity was 8. In addition to temperature and pH there are other factors, such as ionic strength, which can affect the enzymatic reaction. Each of these physical and chemical parameters must be considered and optimized in order for an enzymatic reaction to be accurate and reproducible.

To determine the proper amount of enzymes required for maximum decolorization, the effect of enzyme quantity on dye decolorization was also studied. As shown in table.III, decolorization increased as enzyme quantity increased, from 0.5 to 5 mg/mL. However, the results demonstrated that the minimum enzyme quantity to obtain maximum decolorization was 0.35 µg/mL.

**TABLE: I Optimum temperature for dye degradation of the immobilized laccase enzyme**

S.No	20°C.	30°C.	40°C.	45°C.	50°C.	60°C.	70°C.
Laccase	1.256	1.038	0.947	0.307	0.529	1.077	1.298

**TABLE: II Optimum pH for dye degradation of the immobilized laccase enzyme**

S.No	3	5	7	8	9
Laccase	1.298	0.934	0.541	0.299	0.359


**TABLE: III Optimum enzyme concentration for dye degradation of the immobilized laccase enzyme**

S.No	0.5 mg/ml	0.3 mg/ml	0.35 mg/ml	0.4 mg/ml	0.45 mg/ml
Laccase	1.042	0.563	0.187	0.185	0.187

#### D. Reusability

Reusability of immobilized enzymes exhibits the most important aspect for industrial applications, because immobilization of enzymes decreases the cost of production due to their repeated continuous uses. In this work, reusability of the immobilized enzyme was investigated up to five cycles; the residual activities (Figure. III). During enzymatic reactions, the alginate-gelatin mixture may cause a decrease in the pore sizes of the network, leading to difficulties in the diffusion of the substrate and product in the matrix of the gel. This restriction may cause a decrease in the efficient activity of laccase entrapped in gel after repeated use. In the literature, there are reports of successful reuses of various immobilized laccase systems, such as 60% activity after ten cycles for covalently immobilized laccase on activated polyvinyl alcohol<sup>21, 22, 23.</sup>

**Figure: III– Reusability of immobilized enzyme for dye degradation**



Hence, this study/analysis bears significance for the dye decolourization and this potential isolate *aspergilluspp* has a high potential to decolourize the different types of industrial dyes. The use of these fungi, thus could offer a much cheaper and efficient alternative treatment of waters contaminated heavily with textile dyes<sup>24</sup>.

## CONCLUSIONS

The potential fungal isolate *Aspergilluspp* was capable of decolorizing the different industrial dyes. But the % of decolourization will be varying from one to another. Minimum 55.7% to maximum 90.9% of dye decolourization was detected. That responsible fungal laccase enzyme was monitored further<sup>25</sup>. Immobilization of partially purified laccase enzyme has received increasing attention in dye removal of different types of industrial dyes. Immobilized laccase enzyme Gelatin-alginate beads were prepared as a matrix system for laccase entrapment; the immobilized enzymes exhibited more stability during operation compared to free enzymes, a result that can be considered an advantage in wastewater treatment. To determine the proper amount of enzymes required for maximum decolorization, the effect of immobilized laccase enzyme beads quantity on dye decolorization was also detected. In addition, the reusability of gelatin-alginate beads provides economic benefits when used in largescale applications. Future investigations may focus on the decolorization of different types of dyes that are widely used in the textile industries.

## ACKNOWLEDGEMENT

First and foremost I thank THE ALMIGHTY for being the unfailing source of support, strength and comfort throughout the completion of my research.

My sincere thans to Principal, Hindusthan college of arts and science, Dr. Ponnusamy and I express my deep sense of gratitude and profound thans to Mrs. Saraswathikhannaiyan, secretary Mrs. Priya Satheeshprabhu and joint secretary of hindusthan college of arts and science, coimbatore for their support.

I am highly indebted to **Dr. Laligrowther**, Head of the Department, I wish to express my deep sense of gratitude and heartfelt thanks to my guide **Dr., N. HemashenpagamM.Sc, M.Phil, Ph.D**, Associate Professor, PG and Research Department of Microbiology, Hindusthan College Arts and Science, Coimbatore, for guiding and helping me wholeheartedly in every situation to complete my thesis and made it a successful one and thankful to management also.I express my deep sense of gratitude to all my family members for their love, support, interest and good wishes rendered during the period of my study.

## REFERENCES

1. S. Camarero, D. Ibarra, M. J. Martínez, T. M. Angel. "Lignin-derived compounds as efficient laccase mediators for decolourization of different types of recalcitrant dyes". *Appl Environ Microbiol.* 2005. 71(4):1775–1784.
2. Y. Fu and T. Viraraghavan. Removal of Congo red from an aqueous solution by fungus *Aspergillus niger*. *Advance Environ Res.* 2002; 7:239-247.
3. X.Q Yang, X.X Zhao, C. YLiu, Y. Zheng and S.J Qian. Decolorization of azo, triphenylmethane and anthraquinone dyes by a newly isolated *Trametes* sp. SQ01 and its laccase. *Process Biochem.* 2009; 4: 1185-1189.
4. G. Crini. Non-conventional low-cost adsorbents for dye removal: a review. *Bioresource Technol.* 2006; 97(9):1061-85.
5. Ferrero. Dye removal by low cost adsorbents: Hazelnut shells in comparison with wood sawdust. *J Hazardous Mat.* 2007. 142:144-152.
6. E .Abadulla, T. Tzanov, S. Costa, K.H Robra, A. Cavaco-Paulo and Gubitzi GM. Decolorization and detoxification of textile dyes with a laccase from *Trametes hirsuta*. *Appl Environ Microbiol.* 2000; 66(8):3357–3362.
7. F Gianfreda, Xu, and J.M. Bollag, "Laccases: a useful group of oxidoreductive enzymes," *Bioremediation Journal*, 1999; 3(1): 1–25.
8. Z Konsoula and M. Liakopoulou- Kyriakides. "Thermostable  $\alpha$ -amylase production by *Bacillus subtilis* entrapped in calcium alginate gel capsules," *Enzyme and Microbial Technology*, 2006; 39(4): 690–696.
9. C. J. Silva, Silva, Zille, G. M. Guebitz, and A. Cavaco- Paulo, "Laccase immobilization on enzymatically functionalized polyamide 6,6 fibres," *Enzyme and Microbial Technology*, 2007; 41(6-7): 867–875.
10. S. M Kotwal. And V Shankar. "Immobilized invertase," *Biotechnology Advances*, 2009; 27(4): 311–322.
11. M Forootanfar, A. Faramarzi, A. R. Shahverdi, and M. T. Yazdi, "Purification and biochemical characterization of extracellular laccase from the ascomycete *Paraconiothyrium variable*," *Bioresource Technology*, 2011; 102(2): 1808– 1814.
12. M Forootanfar. M. Movahednia, S. Yaghmaei et al., "Removal of chlorophenolic derivatives by soil isolated ascomycete of *Paraconiothyrium variable* and studying the role of its extracellular laccase," *Journal of Hazardous Materials*, 2012; 209-210 199–203.
13. R Couto. and J. L. T. Herrera, "Industrial and biotechnological applications of laccases: a review," *Biotechnology Advances*, 2006; 24(5): 500–513.

14. M Chivukula. And V. Renganathan. "Phenolic azo dye oxidation by laccase from *Pyriculariaoryzae*," *Applied and Environmental Microbiology*, 1995; 61(12): 4374–4377.
15. L. L. Kiiskinen., MR"att"o and K Kruus."Screening for novel laccase-producing microbes," *Journal of Applied Microbiology*, 2004; 97(3): 640–646.
16. N Mikiashvili, S.P, Wasser, E Nevo, V. Elisashvili. Effects of carbon and nitrogen sources on *Pleurotusostreatus*ligninolytic enzyme activity. *World J. Microbiol. Biotechnol.* 2006. 22, 999-1002.
17. C Srinivasan., D'Souza., K. Boominathan and C. A. Reddy. "A Demonstration of laccase in the white rot basidiomycete *Phanerochaetechrysosporium*," *Applied and Environmental Microbiology*, 1995; 61(12): 4274–4277.
18. M Panouill'e. and V Larreta-Garde. "Gelation behaviour of gelatin and alginate mixtures," *Food Hydrocolloids*, 2009; 23(4): 1074–1080.
19. X. Hu, X. Zhao, and H. M. Hwang, "Comparative study of immobilized *Trametesversicolor*laccaseon nanoparticles and kaolinite," *Chemosphere*, 2007; 66(9): 1618–1626.
20. Hublik and F. Schinner, "Characterization and immobilization of the laccase from *Pleurotusostreatus* and its use for the continuous elimination of phenolic pollutants," *Enzyme and Microbial Technology*, 2000; 27(3–5): 330–336.
21. Yinghui, W. Qiuling, and F. Shiyu, "Laccase stabilization by covalent binding immobilization on activated polyvinyl alcohol carrier," *Letters in Applied Microbiology*, 2002; 35(6): 451–456.
22. N. Vishal Gupta, D. V. Gowda, V. Balamuralidhara, and S. Mohammed Khan, "Formulation and evaluation of olanzapine matrix pellets for controlled release," *DARU, Journal of Pharmaceutical Sciences*, 2011; 19(4): 249–256.
23. G. Mutti, R. Pievo, M. Sgobba, M. Gullotti, and L. Santagostini, "Biomimetic modeling of copper complexes: a study f enantioselective catalytic oxidation on *D-(+)*-catechin and *L-(–)*-epicatechin with copper complexes," *Bioinorganic Chemistry and Applications*, Article ID 762029, 2008; 9.
24. W.M Abd-El-Rahim., H Moawad., and M Khalafallah,. Microflora involved in textile dye waste removal. *J. Basic Microbiol.*,2003; 43: 167-174.
25. Sudhspriyadharsini. S and Hemashenpagam. N. isolation and screening of dye degrading fungi from the industrial effluent soil and water. *IJRAR* Oct, 2018. Volume 5, Issue 4, www.ijrar.org. E-ISSN 2348-1256, P-ISSN 2349-5138.