

## Path Union and Cycle of Graphs with Mean Labeling

Makadia H.M\*

Lukhdhirji Engineering College, MORBI-363642.

E-mail: [makadia.hardik21@gmail.com](mailto:makadia.hardik21@gmail.com)

---

### ABSTRACT

In this paper we investigate mean labeling for path union of  $K_{2,m}$ ,  $P_n$ ,  $P_n \times P_m$ ,  $C_n$ . Also we prove that the mean labeling for cycle of  $P_n$ ,  $C_n$ ,  $P_n \times P_m$ . Path unions of any mean graph are mean graph for that were call Step grid graphics mean graph.

**KEY WORDS:** Cycle, Complete bipartite graph, Grid graph, Step grid graph, Path union of graphs, Cycle of graphs and mean labeling.

---

**\*Corresponding author**

**H. M. Makadia**

Lukhdhirji Engineering College,

Morbi, Gujrat -363642.

E-mail: [makadia.hardik21@gmail.com](mailto:makadia.hardik21@gmail.com)

## 1: INTRODUCTION

We begin with a simple, undirected and finite graph  $G=(V,E)$  with  $|V|=p$  vertices and  $|E|=q$  edges. For all terminology, notations and basic definitions we follow Harary<sup>1</sup>. First of all we give brief summary of definitions which are used in this paper.

**Definition – 1.1:** If the vertices of the graph are assigned values subject to certain conditions then it is known as *graph labeling*.

**Definition – 1.2:** A function  $f$  is called *mean labeling* for a graph  $G = (V, E)$  if  $f: V \rightarrow \{0, 1, \dots, q\}$  is injective and the induced function  $f^*: E \rightarrow \{1, 2, \dots, q\}$  defined as  $f^*(e) = \left\lfloor \frac{f(u)+f(v)}{2} \right\rfloor$  is objective for every edge  $e=(u,v) \in E$ . A graph  $G$  is called *mean graph* if it admits a mean labeling.

**Definition – 1.3:** For a cycle  $C_n$ , each vertex of  $C_n$  is replaced by connected graphs  $G_1, G_2, \dots, G_n$  is known as *cycle of graphs* and we shall denote it by  $C(G_1, G_2, \dots, G_n)$ . If we replace each vertex by a graph  $G$  i.e.  $G_1 = G, G_2 = G, \dots, G_n = G$ , such cycle of a graph  $G$ , we shall denote it by  $C(n \cdot G)$ .

Above definition 1.3 was introduced by Kaneria et. al.<sup>4</sup>

**Definition – 1.4:** Let  $G$  be a graph and  $G_1, G_2, \dots, G_n, n \geq 2$  be  $n$  copies of graph  $G$ . Then the graph obtained by adding an edge from  $G_i$  to  $G_{i+1}$  (for  $i = 1, 2, \dots, n - 1$ ) is called *path union of  $G$* , we shall denote it by  $P(G_1, G_2, \dots, G_n)$ . If we replace each graph  $G_1, G_2, \dots, G_n$  by a graph  $G$  i.e.  $G_1 = G = G_2 = \dots = G_n$ , such path union of  $n$  copies of  $G$ , we shall denote it by  $P(n \cdot G)$ .

For detail survey of various graph labelings and bibliographic references we refer to Gallian [2]. Labelled graphs have many diversified applications. In<sup>3</sup> Somasunderam and Ponraj have introduced the notion of mean labeling of graphs in 2003. They proved that  $P_n, C_n, P_n \times P_m, K_{2,m}$  are mean graphs and  $K_n, K_{1,n}$  are mean graphs iff  $n \leq 3$ . They also prove that  $W_n$  is not a mean graph for  $n > 3$ .

In<sup>5</sup> Kaneria et.al. prove that the step grid graph  $St_n$  where  $n \geq 3$ , is a mean graph with size  $n$ .

**Definition 1.5** Take  $P_n, P_n, P_{n-1}, \dots, P_2$  paths on  $n, n, n-1, n-2, \dots, 3, 2$  vertices and arrange them vertically.

A graph obtained by joining horizontal vertices of given successive paths is known as a step grid graph of size  $n$ , where  $n \geq 3$ . It is denoted by  $St_n$ .

Obviously  $|V(St_n)| = (n^2 + 3n - 2)$  and  $|E(St_n)| = n^2 + n - 2$ .

A Step grid graph  $St_8$  with its mean labeling shown in figure-1.




Figure-1 Mean labeling of  $St_8$ .

They also proved that path union of step grid graph, cycle of step grid graph  $C(r \cdot St_n)$  where  $r \equiv 0 \pmod{2}$  are mean graphs.

A mean graph  $G$  will always have vertices with labels  $q, q - 1$  and  $0$ , where  $q \geq 2$ .

Also two vertices with labels  $q$  and  $q - 1$  are adjacent in the mean graph  $G$ .

In this paper we have proved that path union of any mean graph is also a mean graph and cycle of  $C_n, P_n$  and  $P_n \times P_m$  are mean graphs as well.

## 2: MAIN RESULTS

**Theorem-2.1:** Path union of  $t$  copies of a mean graph  $G$  is also a mean graph.

**Proof :** Let  $G$  be a mean graph with injective mean labeling function  $f: V(G) \rightarrow$

$\{0, 1, \dots, q\}$  and bijective induced function  $f^* : E(G) \rightarrow \{1, 2, \dots, q\}$ .

Let  $V(G) = \{v_i / i = 1, 2, \dots, p\}$ . Since  $\exists v_i, v_j \in V(G)$  such that  $f(v_i) = q$  and  $f(v_j) = 0$ , for some  $i, j \in \{1, 2, \dots, p\}$ , without loss of generality we may assume that  $f(v_1) = q$  and  $f(v_p) = 0$ .

Let  $H$  be the path union of  $t$  copies of the mean graph  $G$ . Let  $u_{i,j} (1 \leq j \leq p)$  be vertices of  $i^{th}$  copy  $G^{(i)}$  of path union  $H$ ,  $\forall i=1,2,\dots,t$ . Now join  $u_{i,1}$  and  $u_{i+1,p}$  by an edge when  $i$  is odd, join  $u_{i,p}$  and  $u_{i+1,1}$  by an edge when  $i$  is even,  $\forall i=1,2,\dots,t-1$  to form the graph  $H$ .

We define the labeling function  $g: V(H) \rightarrow \{0, 1, \dots, Q\}$ , where  $Q = t \cdot q + t - 1$  as follows.

$$\begin{aligned} g(u_{1,j}) &= f(v_j) + (Q - q), & \forall j = 1, 2, \dots, p; \\ g(u_{i,j}) &= g(u_{i-1,j}) - (q + 1), & \forall j = 1, 2, \dots, p, \forall i = 2, 3, \dots, t. \end{aligned}$$

Above labeling pattern give rise a mean labeling to the given  $H$  and so  $H$  is a mean graph.

**Corollary-2.2:** Path union of  $t$  copies of  $K_{2,m}$  is a mean graph.

**Proof :** Let  $H$  be a path union of  $t$  copies of  $K_{2,m}$ . We see that the number of vertices in

$H$  is  $|V(H)| = P = t(m + 2)$  and the number of edges in  $H$  is  $E(H) = Q = 2tm + t - 1$ .

Let  $u_{i,1}, u_{i,2}, v_{i,j} (1 \leq j \leq m)$  be vertices of  $i^{th}$  copy  $K_{2,m}^{(i)}$  of  $H$ ,  $\forall i=1,2,\dots,t$ . Now join

$u_{i,1}$  and  $u_{i+1,2}$  by an edge when  $i$  is odd, join  $u_{i,2}$  and  $u_{i+1,1}$  by an edge when  $i$  is even,

$\forall i=1,2,\dots,t-1$  to form path union of  $t$  copies of  $K_{2,m}$ .

We know that the labeling function  $f: V(K_{2,m}^{(1)}) \rightarrow \{0, 1, \dots, q = 2m\}$  defined by

$$\begin{aligned} f(u_{1,1}) &= q, f(u_{1,2}) = 0 & \text{and} \\ f(u_{1,j}) &= q - (2j - 1), & \forall j = 1, 2, \dots, m \end{aligned}$$

is a mean labeling to the graph  $K_{2,m}$ . Now according to *Theorem-2.1*, we shall define

$g: V(H) \rightarrow \{0, 1, \dots, Q\}$  as follows.

$$\begin{aligned} g(u_{1,j}) &= f(u_{1,j}) + (Q - q), & \forall j = 1, 2; \\ g(v_{1,j}) &= f(v_{1,j}) + (Q - q), & \forall j = 1, 2, \dots, m; \\ g(u_{i,j}) &= g(u_{i-1,j}) - (q + 1), & \forall j = 1, 2, \forall i = 2, 3, \dots, t; \\ g(v_{i,j}) &= g(v_{i-1,j}) - (q + 1), & \forall j = 1, 2, \dots, m, \forall i = 2, 3, \dots, t. \end{aligned}$$

Above labeling pattern give rise mean labeling to the path union of  $t$  copies of  $K_{2,m}$  and so it is a mean graph.

**Illustration – 2.3:** Path union of 4 copies of  $K_{2,3}$  and its mean labeling shown in figure-2.




Figure-2 Path union of 4 copies of  $K_{2,3}$  and its mean labeling.

**Corollary – 2.4:** Path union of  $t$  copies of  $C_n$  is a meangraph.

**Proof :** Let  $H$  be path union of  $t$  copies of  $C_n (n \in \mathbb{N})$ . We see that number of vertices in  $H$  is  $|V(H)| = P = tn$  and number of edges in  $H$  is  $|E(H)| = Q = tn + t - 1$ . Let  $u_{i,j} (1 \leq j \leq n)$  be vertices of  $i^{th}$  copy  $C^{(i)}$  of  $H, \forall i = 1, 2, \dots, t$ . Now join  $u_{i,1}$  and  $u_{i+1,n}$  by an edge when  $i$  is odd, join  $u_{i,n}$  and  $u_{i+1,1}$  by an edge when  $i$  is even,  $\forall i = 1, 2, \dots, t-1$  to form path union of  $t$  copies of  $C_n$ .

We know that the labeling function  $f : V(C_n^{(1)}) \rightarrow \{0, 1, \dots, q = n\}$  defined by

$$f(u_{1,j}) = \begin{cases} q + 1 - j, & \text{when } j \leq \left\lceil \frac{n+1}{2} \right\rceil \\ q - j, & \text{when } j > \left\lceil \frac{n+1}{2} \right\rceil, \forall j = 1, 2, \dots, n \end{cases}$$

is a mean labeling to the graph  $C_n$ . Now according to *Theorem – 2.1*, we shall define


$g : V(H) \rightarrow \{0, 1, \dots, Q\}$ , as follows.

$$g(u_{1,j}) = f(u_{1,j}) + (Q - n), \quad \forall j = 1, 2, \dots, n;$$

$$g(u_{i,j}) = g(u_{i-1,j}) - (n + 1), \quad \forall j = 1, 2, \dots, n, \forall i = 2, 3, \dots, t.$$

Above labeling pattern give rise mean labeling to the graph  $H$  and so  $H$  is a mean graph.

**Illustration–2.5:** Path union of 5 copies of  $C_7$  and its mean labeling shown in figure–3.



**Figure–3** Path union of 5 copies of  $C_7$  and its mean labeling.

**Corollary–2.6:** Path union of  $t$  copies of  $P_n$  is a mean graph.

**Proof :** Let  $H$  be a path union of  $t$  copies of  $P_n (n \in \mathbb{N})$ . We see that the number of vertices in  $H$  is  $tn$  and the number of edges in  $H$  is  $tn-1$ . Let  $u_{i,j} (1 \leq j \leq n)$  be vertices of  $i^{th}$  copy  $P^{(i)}$  of  $H$ ,  $\forall i=1,2,\dots,t$ . Now join  $u_{i,1}$  and  $u_{i+1,n}$  by an edge when  $i$  is odd, join  $u_{i,n}$  and  $u_{i+1,1}$  by an edge when  $i$  is even,  $\forall i=1,2,\dots,t-1$  to form path union of  $t$  copies of  $P_n$ .

We know that the labeling function  $f : V(P_n^{(1)}) \rightarrow \{0, 1, \dots, q=n-1\}$  defined by

$$f(u_{1,j}) = n - j, \quad \forall j=1,2,\dots,n$$

is a mean labeling to the graph  $P_n$ . Now according to *Theorem–2.1*, we shall define


$g : V(H) \rightarrow \{0, 1, \dots, Q\}$ , where  $Q = tn - 1$  as follows.

$$g(u_{1,j}) = f(u_{1,j}) + (Q - q), \quad \forall j=1,2,\dots,n;$$

$$g(u_{i,j}) = g(u_{i-1,j}) - (q+1), \quad \forall j=1,2,\dots,n, \forall i=2,3,\dots,t.$$

Above labeling pattern give rise mean labeling to the graph  $H$  obtained by path union of  $t$  copies of  $P_n$  and so it is a mean graph.

**Illustration – 2.7:** Path union of 7 copies of  $P_5$  and its mean labeling shown in figure-4.



**Figure-4** Path union of 7 copies of  $P_5$  and its mean labeling.

**Corollary – 2.8:** Path union of  $t$  copies of  $P_n \times P_m$  is a mean graph.

**Proof :** Let  $H$  be a path union of  $t$  copies of  $P_n \times P_m$  ( $m, n \in \mathbb{N} - \{1\}$ ). We see that the number of vertices in  $H$  is  $|V(H)| = P = tmn$  and the number of edges in  $H$  is  $|E(H)| = Q = t(q + 1) - 1$ , where  $q = 2mn - (m + n)$ . Let  $u_{i,j,k}$  ( $1 \leq j \leq n, 1 \leq k \leq m$ ) be vertices of  $i^{th}$  copy  $(P_n \times P_m)^{(i)}$  of  $H, \forall i = 1, 2, \dots, t$ . Now join  $u_{i,n,m}$  with  $u_{i+1,1,1}$  by an edge  $\forall i = 1, 2, \dots, t-1$  to form the graph  $H$ .

We know that the labeling function  $f: V((P_n \times P_m)^{(1)}) \rightarrow \{0, 1, \dots, q\}$ , where  $q = 2mn - (m + n)$  defined by

$$f(u_{1,j,k}) = q - (2m - 1)(j - 1) - (k - 1), \quad \forall j = 1, 2, \dots, n, \forall k = 1, 2, \dots, m$$


is a mean labeling to the graph  $(P_n \times P_m)^{(1)}$ . Now define  $g: V(H) \rightarrow \{0, 1, \dots, Q\}$  as follows.

$$g(u_{i,j,k}) = Q - (q + 1)(i - 1) - (2m - 1)(j - 1) - (k - 1),$$

$\forall i = 1, 2, \dots, t, \forall j = 1, 2, \dots, n, \forall k = 1, 2, \dots, m$ .

Above labeling pattern give rise mean labeling to the graph  $H$  obtained by path union of  $t$  copies of grid graph  $P_n \times P_m$  and so  $H$  is a mean graph.

**Illustration–2.9** :Path union of 5 copies of  $P_3 \times P_3$  and its mean labeling shown in figure–5.



**Figure–5** Path union of 3 copies of  $P_3 \times P_4$  and its mean labeling.

**Theorem–2.10:**  $C(t \cdot P_n)$  is a mean graph, where  $t \equiv 0 \pmod{2}$ .

**Proof :**Let  $G = C(t \cdot P_n)$ , where  $n \in \mathbb{N}$ . It is obvious that  $P = |V(G)| = tn = Q =$

$|E(G)|$ . Let  $u_{i,j}(1 \leq j \leq n, 1 \leq i \leq t)$  be vertices of graph  $G$ . We shall join  $u_{i,1}$  with  $u_{i+1,n}$ ,

When  $i + \frac{t}{2}$  is odd and  $u_{i,n}$  with  $u_{i+1,1}$ , When  $i + \frac{t}{2}$  is even to form the cycle graph  $G = C(t \cdot P_n)$ .

Now define the labeling function :  $V(G) \rightarrow \{0, 1, \dots, Q\}$  as follows.

$$g(u_{i,j}) = Q - n(i - 1) - (j - 1), \forall j=1,2,\dots,n, \forall i=1,2,\dots,\frac{t}{2};$$

$$g\left(u_{\frac{t}{2}+1,1}\right) = \frac{Q}{2};$$


$$g\left(u_{\frac{t}{2}+1,j}\right) = \frac{Q}{2} - j, \quad \forall j=1,2,\dots,n;$$

$$g(u_{i,j}) = Q - n(i - 1) - j, \forall j=1,2,\dots,n, \forall i = \frac{t}{2} + 2, \frac{t}{2} + 3, \dots, t.$$

Above labeling pattern give rise mean labeling to the graph  $G$  obtained by taking cycle of path  $P_n$  and so  $G$  is a mean graph.



**Illustration–2.11 :**  $C(6 \cdot P_5)$  and its mean labeling shown in figure–6.



**Figure–6** Cycle graph  $C(6 \cdot P_5)$  and its mean labeling.

**Theorem–2.12:**  $C(t \cdot C_n)$  is a mean graph, where  $n \in \mathbb{N}$  and  $t \equiv 0 \pmod{2}$ .

**Proof :** Let  $G = C(t \cdot C_n)$ , where  $n \in \mathbb{N}$ . It is obvious that  $P = |V(G)| = tn$  and  $Q = |E(G)| = t(n + 1)$ . Let  $u_{i,j} (1 \leq j \leq n, 1 \leq i \leq t)$  be vertices of graph  $G$ . We shall join  $u_{i,1}$  with  $u_{i+1,n}$ , when  $i + \frac{i}{2}$  is odd and  $u_{i,n}$  with  $u_{i+1,1}$ , when  $i + \frac{i}{2}$  is even to form the cycle graph  $G = C(t \cdot C_n)$ .

Now define the labeling function :  $V(G) \rightarrow \{0, 1, \dots, Q\}$  as follows.

$$g(u_{i,j}) = Q - (n + 1)(i - 1) - (j - 1), \text{ when } j \leq \left\lfloor \frac{n+1}{2} \right\rfloor$$

$$= Q - (n+1)(i-1) - j, \quad \text{when } j > \left\lfloor \frac{n+1}{2} \right\rfloor,$$

$$\forall j = 1, 2, \dots, n, \forall i = 1, 2, \dots, \frac{t}{2};$$

$$g\left(u_{\frac{t}{2}+1,1}\right) = \frac{Q}{2};$$

$$g\left(u_{\frac{t}{2}+1,1}\right) = \frac{Q}{2} - j, \quad \forall j = 1, 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor;$$

$$= \frac{Q}{2} - (j + 1), \forall j = \left\lceil \frac{n+2}{2} \right\rceil, \left\lceil \frac{n+4}{2} \right\rceil, \dots, n;$$

$$g(u_{i,j}) = Q - (n + 1)(i - 1) - j, \text{ when } j \leq \left\lceil \frac{n+1}{2} \right\rceil$$


$$= Q - (n + 1)(i - 1) - (j + 1), \quad \text{when } j > \left\lceil \frac{n+1}{2} \right\rceil$$

$$\forall j = 1, 2, \dots, n, \forall i = \frac{t}{2} + 2, \frac{t}{2} + 3, \dots, t.$$

Above labeling pattern give rise mean labeling to the cycle graph  $G$  obtained by  $C_n$

and so  $G$  is a mean graph.

**Illustration**–2.13 :  $C(4 \cdot C_7)$  and its mean labeling shown in figure–7.



Figure–7 Cycle graph  $C(4 \cdot C_7)$  and its mean labeling.

**Theorem**–2.14:  $C(t \cdot P_n \times P_m)$  is a mean graph, where  $m, n \in \mathbb{N}$  and  $t \equiv 0 \pmod{2}$ .

**Proof:** Let  $G = C(t \cdot P_n \times P_m)$ , where  $n, m \in \mathbb{N}$ . It is obvious that  $P = |V(G)| = tmn$  and  $Q = |E(G)| = t(2mn - (m + n) + 1)$ . Let  $u_{i,j,k}$  ( $1 \leq j \leq n, 1 \leq k \leq m, 1 \leq i \leq t$ ) be vertices of graph  $G$ . We shall join  $u_{i,1,1}$  with  $u_{i+1,n,m}, \forall i = 1, 2, \dots, t-1$  to form the cycle graph  $G = C(t \cdot P_n \times P_m)$ .

Now define the labeling function:  $V(G) \rightarrow \{0, 1, \dots, Q\}$ , where  $Q = t(q+1)$  and  $q = 2mn - (m + n)$  as follows.

$$g(u_{i,j,k}) = Q - (q + 1)(i - 1) - (2m - 1)(j - 1) - (k - 1),$$

$$\forall j = 1, 2, \dots, n, \forall k = 1, 2, \dots, m, \forall i = 1, 2, \dots, \frac{t}{2}.$$

$$g\left(u_{\frac{t}{2}+1,1,1}\right) = \frac{Q}{2};$$

$$g\left(u_{\frac{t}{2}+1,j,k}\right) = Q - (q + 1)(i - 1) - (2m - 1)(j - 1) - k,$$

$$\forall j = 1, 2, \dots, n, \forall k = 1, 2, \dots, m,$$

$$g(u_{i,j}) = Q - (n + 1)(i - 1) - j,$$


$$\forall j = 1, 2, \dots, n, \forall k = 1, 2, \dots, m;$$

$$g(u_{i,j,k}) = Q - (q + 1)(i - 1) - (2m - 1)(j - 1) - k,$$

$$\forall j = 1, 2, \dots, n, \forall k = 1, 2, \dots, m, \forall i = \frac{t}{2} + 2, \frac{t}{2} + 3, \dots, t.$$

Above labeling pattern give rise mean labeling to the graph  $G$  obtained by taking cycle of grid graph  $(P_n \times P_m)$  and so  $G$  is a mean graph.

**Illustration**–2.15 :  $C(6 \cdot P_3 \times P_3)$  and its mean labeling shown in figure–8.



Figure–8 Cycle graph  $C(6 \cdot P_3 \times P_3)$  and its mean labeling.

### 3: CONCLUDING REMARKS

Here we have discussed mean labeling for path union of  $C_n$ ,  $P_n$ ,  $K_{2,m}$  and  $P_n \times P_m$ . Also we proved that cycle of  $C_n$ ,  $P_n$ ,  $P_n \times P_m$  are mean graphs. These results contribute some new topics to the families of mean graphs. The labeling pattern is demonstrated by means of illustrations.

*Theorem-2.1* is a strong result of general nature, as it shows  $P(t_1 \cdot P(t_2 \cdot K_{2,m}))$ ,  $P(t_1 \cdot P(t_2 \cdot P_n))$ ,  $P(t_1 \cdot P(t_2 \cdot C_n))$ ,  $P(t_1 \cdot P(t_2 \cdot (P_n \times P_m)))$ ,  $P(t_1 \cdot C(t_2 \cdot P_n))$ ,  $P(t_1 \cdot C(t_2 \cdot C_n))$  and  $P(t_1 \cdot C(t_2 \cdot (P_n \times P_m)))$  are mean graphs. We raise an open question to get mean labeling for the graphs  $C(t_1 \cdot C(t_2 \cdot P_n))$ ,  $C(t_1 \cdot C(t_2 \cdot C_n))$ ,  $C(t_1 \cdot C(t_2 \cdot (P_n \times P_m)))$ .

### REFERENCES

1. F. Harary, Graph theory *Addition Wesley, Massachusetts*, 1972.
  2. J. A. Gallian, *The Electronics Journal of Combinatorics*, 19, #DS6(2013).
  3. S. Somasunderam and R. Ponraj, Mean labelings of graphs, *Natl. Acad. Sci. Let.*, 26 (2003) pp. 210 – 213.
  4. V. J. Kaneria, H. M. Makadia and M. M. Jariya, Graceful labeling for cycle of graphs, *Int. J. of Math. Res.*, vol-2014;6(2): 173 – 178.
  5. V. J. Kaneria, Meera Meghpara and H. M. Makadia, Mean labeling for Step grid graph, *Advances and Applications in Mathematical Sciences*, 2015;14(4): 111 – 119.
-