

International Journal of Scientific Research and Reviews

Effect of Long Term Tobacco Use on Unstimulated Salivary Ph And Buffering Capacity

L Krishna Prasada

Professor, Head of the Department
Dept of Conservative Dentistry & Endodontics
KVG Dental College & Hospital Kurunjibhag, Sullia D.K
Karnataka, India 574327

ABSTRACT

Background: It is firmly established that tobacco use is a primary cause of many oral diseases and adverse oral conditions. The negative impact relates not only to smoking but use of smokeless tobacco. Saliva is the first biological fluid that is exposed to tobacco which contains numerous toxic compositions responsible for structural and functional changes in saliva. Alterations in salivary pH and buffering capacity have a significant impact on oral and dental health. There are several studies concerning the effect of chewing tobacco and smoking on salivary secretion, though, long-term effect of tobacco use on pH and buffering capacity is still not clear.

Aims and Objectives: The aim of this study is to analyze and compare the long- term effects of tobacco on salivary pH and buffering capacity between tobacco chewers, smokers, and controls.

Materials and Methods: Subjects will be divided equally into 4 groups; tobacco smokers (group A), chewers (group B), chewers+ smokers (group C) and controls (group D). Saliva of each subject will be collected under resting condition. The salivary pH and buffering capacity will be measured using GC Saliva-Check Buffer kit.

Statistical Analysis: Data will be analyzed using the Statistical Package for Social Service (SPSS) computer software. Unpaired Student's t-test, one-way ANOVA will be applied to assess the difference between groups.

Results: Within its limitations, the results of the present study suggest that pH and buffering capacity is lower (acidic) in tobacco chewers, chewers+ smokers and tobacco smokers. While comparing the 3 groups, groups in which tobacco chewing habit is present showed more acidic pH.

KEY WORDS: Tobacco, salivary pH, buffering capacity

Corresponding Author:

Dr Krishnan Venugopal*

Post Graduate student

Dept of Conservative Dentistry & Endodontics, KVG Dental College & Hospital

Kurunjibhag, Sullia D.K, Karnataka, India 574327

Ph: +91 9895828996

INTRODUCTION

The epidemic of tobacco use is one of the greatest threats to global health today ¹. Approximately one-third of the adult population in the world use tobacco in some form and of whom half will die prematurely ². According to the World Health Organization (WHO) estimates, globally, there were 100 million premature deaths due to tobacco in the 20th century, and if the current trends of tobacco use continue, this number is expected to rise to 1 billion in the 21st century ¹. In addition to several other chronic diseases, tobacco use is a primary cause of many oral diseases and adverse oral conditions. Various clinical and epidemiological evidence have been documented regarding the adverse effects of tobacco on oral health ³. The adverse effects of cigarette smoking and other smokeless forms are numerous and use of tobacco has been associated with oral mucosa, gingival diseases and dental alterations ⁴.

Saliva is a complex and important body fluid which is very essential for oral health ⁵. It plays a critical role in oral homeostasis because it modulates the ecosystem within the oral cavity ⁶. Taking this into account, quantitative and/or qualitative alterations in salivary secretion may lead to local (caries, oral mucositis, candidiasis, oral infections, chewing disorders) or extraoral (dysphagia, halitosis, weight loss) adverse effects ^{7,8,9}. Saliva is the first biological fluid that is exposed to cigarette smoke, containing numerous toxic compositions responsible for structural and functional changes in saliva. Alterations in salivary pH and buffering capacity have a significant impact on oral and dental health ¹⁰. Several studies of resting salivary pH estimate a range of 6.5–7.9 ¹².

Approximately 600 million people use arecanut worldwide in some form and is the fourth most commonly used psychoactive substance ¹². Arecanut contains four major alkaloids: Arecaidine, arecoline, guvacine and guvacoline. In the presence of lime (calcium oxide which turns to alkali calcium hydroxide in aqueous form), arecoline and guvacoline are largely hydrolyzed into arecaidine and guvacine, respectively ¹³. The main ingredient of tobacco is nicotine and nicotine acts on certain cholinergic receptors in the brain and other organs causing neural activation leading to altered salivary secretion ¹⁴.

There are several studies concerning the effect of chewing tobacco and smoking on salivary secretion, though, long-term effect of tobacco use on pH is still not clear. Due to the paucity of literature on the influence of tobacco use on pH, this study was undertaken to analyze and compare the long-term effect of tobacco on pH in tobacco chewers, tobacco smokers and control.

AIM & OBJECTIVE

The aim of this study is to analyze and compare the long-term effects of tobacco on salivary pH and buffering capacity between tobacco chewers, smokers, and controls.

MATERIALS AND METHODS

The present study consisted of 64 subjects (males and females) within the age group of 25-40 years referred to KVG Dental College & Hospital were randomly selected and equally divided into four groups of 16 subjects each .

Group A: Tobacco smokers

Group B: Tobacco chewers

Group C: Tobacco chewer + smokers

Group D: Control

INCLUSION CRITERIA

- Males and females of age between 25 and 40 years
- Consumption of tobacco (smoked and smokeless form) for minimum period of around 5 years.
- Subjects volunteering to take part in study.

EXCLUSION CRITERIA

- Age over 40 years.
- Alcohol consumption.
- History of any other habits (tongue thrusting, mouth breathing, bruxism).
- Denture wearers
- Pregnant and postmenopausal women.
- History of radiotherapy.
- Uncooperative patients.
- Patients with systemic or salivary gland diseases or under any drug therapy.

Informed consent was obtained from each subject for saliva collection. Thorough history was taken along with detailed oral examination. Collection of saliva each subject was done under resting condition.

The pH and buffering capacity of saliva was measured using GC Saliva Buffer Kit (GC India).

SALIVA COLLECTION

Saliva collection was done under resting conditions to obtain unstimulated saliva. Saliva collection was carried out between 9.00 am and 1.00 pm to avoid diurnal variations. Subjects were requested not to drink, eat or perform oral hygiene or chew or smoke 60 min before procedure. Subjects are then seated on the dental chair and asked to spit 2–3 times into an 1 min in a disposable

container. The patient was instructed not to speak, chew or perform any other activities during the procedure (figure 1, 2).

Testing of salivary pH and buffering capacity was done immediately after collection of the samples using GC Saliva Buffer Kit.



Fig 1: Collection of saliva sample



Fig 2: Collected saliva sample

TESTING OF SALIVARY pH

A drop of saliva sample collected is transferred to a litmus paper. After 10 seconds the color of the paper is compared with the value in the pH scale and the value is recorded (figure 3, 4).



Fig 3: Testing salivary pH



Fig 4: Comparing value with pH indicator

TESTING OF SALIVARY BUFFERING CAPACITY

3 drops of saliva sample collected is transferred to the 3 segments of the testing paper. After 2 minutes the color in the 3 segments is recorded (figure 5). Values are added and buffering capacity is recorded as Normal or Low.



Fig 3: Testing salivary buffering capacity


STATISTICAL ANALYSIS

Data was analyzed using the Statistical Package for Social Service (SPSS) computer software. Unpaired Student's *t*-test, one-way ANOVA was applied to assess the pH difference between different groups. $P < 0.05$ was considered as statistically significant. The confidence of 95% was considered.

RESULTS

The subjects in our study were present in the age group of 25–40 years. Group A, B and C subjects consume tobacco for minimum of around 5 years. The mean pH scores of saliva in four distinct groups showed that pH scores were maximum in the control group while it was least in tobacco chewers group (graph 1).

When unpaired *t*-test was applied for comparison of pH scores of saliva between different groups, results showed a significant difference between pairs of groups at 0.05 level of significance, i.e. ($P < 0.05$) (table 1).



Graph 1- mean pH scores of different groups

GROUP	N	t value	p value
TOBACCO SMOKERS	16	3.03	P = 0.0413**
CONTROL	16		
TOBACCO CHEWERS	16	6.7524	P < 0.0001**
CONTROL	16		
TOBACCO SMOKERS+ CHEWERS	16	4.52	P < 0.0001**
CONTROL	16		


Table 1- comparing values of different groups to control statistically**

When one way ANOVA test was applied for comparing the pH scores of saliva among four groups, it showed a significant difference in pH scores of saliva among four groups (table 2).

Source	SS	df	MSS	F	P
Between groups	2.0469	3	0.6823	19.61	< 0.0001**
Within Group	2.0875	60	0.0348		
Total	4.1344	63			

Table 2- intergroup comparison of pH values ** statistically significant

The mean scores for buffering capacity in four distinct groups showed normal buffering capacity in the control group while it was significantly low in tobacco using groups (graph 2).



Graph 2- mean buffering capacity scores of different groups

When one way ANOVA test was applied for comparing the buffering capacity of saliva among four groups, it showed a significant difference in buffering capacity of saliva among four groups (table 3).

Source	SS	df	MSS	F	P
Between groups	8.1875	3	2.729167	34.47	< 0.0001**
Within Group	4.75	60	0.079167		
Total	12.9375	63			

Table 3- intergroup comparison of buffering capacity statistically significant**

DISCUSSION

Saliva is a complex and important body fluid which is very essential for oral health ⁵. Saliva is required for protecting the oral mucosa, teeth remineralization, digestion, taste sensation, pH balance and phonation. It includes a variety of electrolytes, peptides, glycoproteins and lipids which have antimicrobial, antioxidant, tissue repair and buffering properties ¹⁵. Saliva is the first biological

fluid that is exposed to cigarette smoke, which contains numerous toxic compositions responsible for structural and functional changes in saliva¹⁰.

From the inferences of the present study it is clear that tobacco users have more acidic pH and lower buffering capacity as compared to non tobacco users which may make them more prone dental caries. On comparing tobacco chewers to smokers, the chewers had more acidic pH and lower buffering capacity as compared to tobacco smokers & the difference were extremely statistically significant.

The decrease in salivary pH and buffering capacity is mostly attributed to the decreased salivary flow rate in smokers which reduces the bicarbonate ion concentration in saliva and hence reducing pH and buffering capacity, in addition to this Lime present in mostly all forms of chewing tobacco also reacts with the available bicarbonate ions leading to loss of bicarbonate ions available. The alteration in electrolytes and ions alters the pH as they interact with the buffering systems of saliva.

Voelker et al in his study found relationship between caries risk and smoking, buffering capacity and smoking, and stimulated salivary pH and smoking were concluded. No significance difference between *S. mutans* and smoking were noted from the preliminary results¹⁶.

Grover et al in 2015 in his study observed that a lower (acidic) salivary pH was observed in tobacco users as compared with control. These alterations in pH due to the long-term effect of tobacco use can render oral mucosa vulnerable to various oral and dental diseases⁴. Khan *et al.* in 2010 also observed a lower salivary pH in smokers than in nonsmokers which was consistent with the findings of the present study¹⁷. Rooban *et al.* 2008 observed a mean pH of 6.77 in nonchewers and those who chew tobacco, the mean pH turns acidic¹⁸.

But in contrast Reddy *et al.* and Alpana Kanwar *et al.* observed no difference in salivary pH between the chewers and nonchewers. This difference could be due to the amount of tobacco, lime and other components. The role of lime has been a source of concern. Lime (calcium oxide in aqueous forms calcium hydroxide) could cause a free radical injury or the high alkaline content probably reacts with the salivary buffering systems and alters the pH^{19,20}.

A salivary pH of 7.0 usually indicates a healthy dental and periodontal situation. At this pH, there is a low incidence of dental decay and little or no calculus. A saliva pH below 7.0 usually indicates acidemia (abnormal acidity of the blood). If a chronic condition exists, the mouth is more susceptible to dental decay, halitosis and periodontitis. Chronic acidemia can be a causative factor for a multitude of diseases affecting the whole body²¹.

CONCLUSION

Within its limitations, the results of the present study suggest that pH and buffering capacity is lower (acidic) in tobacco chewers, chewers+ smokers and tobacco smokers. While comparing the 3 groups, groups in which tobacco chewing habit is present showed more acidic pH. This is, to the best of our knowledge, the first study which has reported on the pH & Buffering capacity of saliva in tobacco users..

With a high prevalence of tobacco consumption in different forms, the oral health of the population is at risk, and oral health programmes to increase the awareness of the public to the health hazards of tobacco consumption need to be implemented.

Moreover, due to the high prevalence of smokeless tobacco use in this population, there is a need for further studies with an improved study design to better understand the effects of the different forms of tobacco products on pH, buffering capacity and various other parameters.

Declaration of patient consent

Consent from the patient is obtained before the collection of samples. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be recorded. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Petersen PE. Tobacco and oral health-the role of the World Health Organization. *Oral Health and Preventive Dentistry*. Dec, 2003; 1(4):309-16.
2. Vellappally S, Fiala Z, Šmejkalová J, Jacob V, Shriharsha P. Influence of tobacco use in dental caries development. *Central European journal of public health*. Sep 1, 2007; 15(3).
3. Millar WJ, Locker D. Smoking and oral health status. *J Can Dent Assoc*. 2007;73:155.
4. Grover N Sharma J, Sengupta S, Singh S, Kaur H. Long-term effect of tobacco on unstimulated salivary pH. *J Oral Maxillofac Pathol*. Jan-Apr, 2016; 20(1): 16–19.

5. Rad M, Kakoie S, Niliye Brojeni F, Pourdamghan N. Effect of long-term smoking on whole-mouth salivary flow rate and oral health. *J Dent Res Dent Clin Dent Prospects*. 2010;4:110–4.
6. Atkinson JC, Baum BJ. Salivary enhancement: Current status and future therapies. *J Dent Educ*. 2001;65:1096–101.
7. Atkinson JC, Wu AJ. Salivary gland dysfunction: Causes, symptoms, treatment. *J Am Dent Assoc*. 1994;125:409–16
8. Ship JA, Pillemer SR, Baum BJ. Xerostomia and the geriatric patient. *J Am Geriatr Soc*. 2002;50:535–43.
9. Valdez IH, Fox PC. Interactions of the salivary and gastrointestinal systems. II. Effects of salivary gland dysfunction on the gastrointestinal tract. *Dig Dis*. 1991;9:210–8.
10. Fox PC, Ship JA. Salivary gland diseases. In: Greenberg MS, Glick S, editors. *Burket's Oral Medicine: Diagnosis and Treatment*. 11th ed. Hamilton: BC Decker, 2008; 191–223.
11. Choo RE, Huestis MA. Oral fluid as a diagnostic tool. *Clin Chem Lab Med*. 2004;42:1273–87.
12. Trivedy CR, Craig G, Warnakulasuriya S. The oral health consequences of chewing areca nut. *Addict Biol*. 2002;7:115–25.
13. Chu NS. Effects of betel chewing on the central and autonomic nervous systems. *J Biomed Sci*. 2001;8:229–36.
14. Bouquot DJ, Schroeder K. Oral effect of tobacco abuse. *J Am Dent Inst Contin Educ*. 1992;43:3–17.
15. Zappacosta B, Persichilli S, De Sole P, Mordente A, Giardina B. Effect of smoking one cigarette on antioxidant metabolites in the saliva of healthy smokers. *Arch Oral Biol*. 1999;44:485–8.
16. Voelker M A, Simmer-Beck M, Cole M, Keeven E, Tira D. Preliminary findings on the correlation of saliva pH, buffering capacity, flow, Consistency and *Streptococcus mutans* in relation to cigarette smoking. *J Dent Hyg*. Feb, 2013; 87(1):30-7. Epub 2013 Feb 5.
17. Khan GJ, Javed M, Ishaq M. Effect of smoking on salivary flow rate. *J Med Sci*. 2010;8:221–4.
18. Rooban T, Mishra G, Elizabeth J, Ranganathan K, Saraswathi TR. Effect of habitual arecanut chewing on resting whole mouth salivary flow rate and pH. *Indian J Med Sci*. 2006;60:95–105.
19. Reddy MS, Naik SR, Bagga OP, Chuttani HK. Effect of chronic tobacco-betel-lime “quid” chewing on human salivary secretions. *Am J Clin Nutr*. 1980;33:77–80.

20. Kanwar A, Sah K, Grover N, Chandra S, Singh RR. Long-term effect of tobacco on resting whole mouth salivary flow rate and pH: An institutional based comparative study. *Eur J Gen Dent.* 2013;2:296–9.
 21. Baliga S, Muglikar S, Kale R. Salivary pH: A diagnostic biomarker. *J Indian Soc Periodontol.* 2013;17:461–5.
-