

## *International Journal of Scientific Research and Reviews*

### **A note on Cubic Difference Prime Labeling**

**Sunoj B.S.<sup>1\*</sup> and Mathew Varkey T. K.<sup>2</sup>**

<sup>1</sup>Department of Mathematics, Government Polytechnic College, Attingal, Kerala

<sup>2</sup>Department of Mathematics, T K M College of Engineering, Kollam 5, Kerala

[spalazhi@yahoo.com](mailto:spalazhi@yahoo.com), [mathewvarkeytk@gmail.com](mailto:mathewvarkeytk@gmail.com)

#### **ABSTRACT**

Cubic difference prime labeling of a graph is the labeling of the vertices with  $\{0,1,2,\dots,p-1\}$  and the edges with absolute difference of the cubes of the labels of the incident vertices. The greatest common incidence number of a vertex (*gcin*) of degree greater than one is defined as the greatest common divisor of the labels of the incident edges. If the *gcin* of each vertex of degree greater than one is one, then the graph admits cubic difference prime labeling. Here we identify wheel graph, gear graph, fan graph, friendship graph, prism graph, key graph, helm graph and umbrella graph for cubic difference prime labeling.

**KEYWORDS:** Graph labeling, greatest common incidence number, cubic difference, prime labeling.

#### **\*Corresponding Author**

**B.S. Sunoj**

Department of Mathematics, Government Polytechnic College,

Attingal, Kerala

E Mail - [spalazhi@yahoo.com](mailto:spalazhi@yahoo.com),

## 1. INTRODUCTION

All graphs in this paper are simple, finite, connected and undirected. The symbol  $V(G)$  and  $E(G)$  denotes the vertex set and edge set of a graph  $G$ . The graph whose cardinality of the vertex set is called the order of  $G$ , denoted by  $p$  and the cardinality of the edge set is called the size of the graph  $G$ , denoted by  $q$ . A graph with  $p$  vertices and  $q$  edges is called a  $(p,q)$ - graph.

A graph labeling is an assignment of integers to the vertices or edges. Some basic notations and definitions are taken from <sup>1,2,3,4</sup>. Some basic concepts are taken from <sup>1</sup> and <sup>2</sup>. In this paper we investigated cube difference prime labeling of wheel graph, gear graph, fan graph, friendship graph, prism graph, key graph, helm graph and umbrella graph.

**Definition: 1.1** Let  $G$  be a graph with  $p$  vertices and  $q$  edges. The greatest common incidence number (*gcin*) of a vertex of degree greater than or equal to 2, is the greatest common divisor (gcd) of the labels of the incident edges.

## 2. MAIN RESULTS

**Definition: 2.1**  $G = (V(G), E(G))$  be a graph with  $p$  vertices and  $q$  edges. Define a bijection  $f : V(G) \rightarrow \{0, 1, 2, 3, \dots, p-1\}$  by  $f(v_i) = i-1$ , for every  $i$  from 1 to  $p$  and define a 1-1 mapping  $f_{cdpl}^* : E(G) \rightarrow \text{set of natural numbers } N$  by  $f_{cdpl}^*(uv) = | \{f(u)\}^3 - \{f(v)\}^3 |$ . The induced function  $f_{cdpl}^*$  is said to be cubic difference prime labeling, if for each vertex of degree at least 2, the greatest common incidence number of the labels of the incident edges is 1.

**Definition: 2.2** Graph which admits cubic difference prime labeling is called a cubic difference prime graph.

**Theorem 2.1** Wheel graph  $W_n$  admits cubic difference prime labeling.

Proof: Let  $G = W_n$  and let  $v_1, v_2, \dots, v_{n+1}$  are the vertices of  $G$

Here  $|V(G)| = n+1$  and  $|E(G)| = 2n$

Define a function  $f : V \rightarrow \{0, 1, 2, 3, \dots, n\}$  by

$$f(v_i) = i-1, i = 1, 2, \dots, n+1$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$f_{cdpl}^*(v_i v_{i+1}) = i^3 - (i-1)^3, \quad i = 1, 2, \dots, n-1$$

$$f_{cdpl}^*(v_{n+1} v_i) = n^3 - (i-1)^3, \quad i = 1, 2, \dots, n$$

$$f_{cdpl}^*(v_1 v_n) = (n-1)^3$$

Clearly  $f_{cdpl}^*$  is an injection.

$$\begin{aligned} gcin \text{ of } (v_{i+1}) &= \gcd^2 \text{ of } \{ f_{cdpl}^*(v_i v_{i+1}), f_{cdpl}^*(v_{i+1} v_{i+2}) \} \\ &= \gcd \text{ of } \{ 3i^2 - 3i + 1, 3i^2 + 3i + 1 \} \\ &= \gcd \text{ of } \{ 6i^2, 3i^2 - 3i + 1 \} \\ &= \gcd \text{ of } \{ 3i^2, 3i^2 - 3i + 1 \} \\ &= \gcd \text{ of } \{ 3i - 1, 3i^2 - 3i + 1 \} \\ &= \gcd \text{ of } \{ i, 3i - 1 \} \\ &= \gcd \text{ of } \{ i, i - 1 \} = 1, \quad i = 1, 2, \dots, n-1 \end{aligned}$$

$$\begin{aligned} gcin \text{ of } (v_1) &= \gcd \text{ of } \{ f_{cdpl}^*(v_1 v_2), f_{cdpl}^*(v_1 v_n) \} \\ &= \gcd \text{ of } \{ 1, (n-1)^3 \} = 1. \end{aligned}$$

$$gcin \text{ of } (v_{n+1}) = 1.$$

So,  $gcin$  of each vertex of degree greater than one is 1.

Hence  $W_n$ , admits cubic difference prime labeling. ■

**Theorem 2.2** Gear graph<sup>3</sup>  $G_n$  admits cubic difference prime labeling.

Proof: Let  $G = G_n$  and let  $v_1, v_2, \dots, v_{2n+1}$  are the vertices of  $G$

Here  $|V(G)| = 2n+1$  and  $|E(G)| = 3n$

Define a function  $f : V \rightarrow \{0, 1, 2, 3, \dots, 2n\}$  by

$$f(v_i) = i-1, \quad i = 1, 2, \dots, 2n+1$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$f_{cdpl}^*(v_i v_{i+1}) = i^3 - (i-1)^3, \quad i = 1, 2, \dots, 2n$$

$$f_{cdpl}^*(v_{2n+1} v_{2i}) = (2n)^3 - (2i-1)^3, \quad i = 1, 2, \dots, n-1$$

$$f_{cdpl}^*(v_1 v_{2n}) = (2n-1)^3,$$

Clearly  $f_{cdpl}^*$  is an injection.

$$gcin \text{ of } (v_{i+1}) = 1, \quad i = 1, 2, \dots, 2n-1$$

$$gcin \text{ of } (v_1) = \gcd \text{ of } \{ f_{cdpl}^*(v_1 v_2), f_{cdpl}^*(v_1 v_{2n}) \}$$

$$= \gcd \text{ of } \{1, (2n-1)^3\} = 1$$

$$\mathbf{gcin} \text{ of } (v_{2n+1}) = 1.$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence  $G_n$ , admits cubic difference prime labeling. ■

**Theorem 2.3** Fan graph<sup>4</sup>  $F_n$  admits cubic difference prime labeling.

Proof: Let  $G = F_n$  and let  $v_1, v_2, \dots, v_{n+1}$  are the vertices of  $G$

Here  $|V(G)| = n+1$  and  $|E(G)| = 2n-1$

Define a function  $f : V \rightarrow \{0, 1, 2, 3, \dots, n\}$  by

$$f(v_i) = i-1, \quad i = 1, 2, \dots, n+1$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$f_{cdpl}^*(v_i v_{i+1}) = i^3 - (i-1)^3, \quad i = 1, 2, \dots, n$$

$$f_{cdpl}^*(v_{n+1} v_i) = (n)^3 - (i-1)^3, \quad i = 1, 2, \dots, n-1$$

Clearly  $f_{cdpl}^*$  is an injection.

$$\mathbf{gcin} \text{ of } (v_{i+1}) = 1, \quad i = 1, 2, \dots, n-1$$

$$\mathbf{gcin} \text{ of } (v_1) = 1.$$

$$\mathbf{gcin} \text{ of } (v_{n+1}) = 1.$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence  $F_n$ , admits cubic difference prime labeling. ■

**Theorem 2.4** Prism graph<sup>3</sup>  $Y_n$  admits cubic difference prime labeling.

Proof: Let  $G = Y_n$  and let  $v_1, v_2, \dots, v_{2n}$  are the vertices of  $G$

Here  $|V(G)| = 2n$  and  $|E(G)| = 3n$

Define a function  $f : V \rightarrow \{0, 1, 2, 3, \dots, 2n-1\}$  by

$$f(v_i) = i-1, \quad i = 1, 2, \dots, 2n$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$f_{cdpl}^*(v_i v_{i+1}) = i^3 - (i-1)^3, \quad i = 1, 2, \dots, 2n-1$$

$$f_{cdpl}^*(v_i v_{2n+i-1}) = (2n-i)^3 - (i-1)^3, \quad i = 1, 2, \dots, n-1$$

$$f_{cdpl}^*(v_1 v_n) = (n-1)^3,$$

$$f_{cdpl}^*(v_{n+1} v_{2n}) = (2n-1)^3 - n^3.$$

Clearly  $f_{cdpl}^*$  is an injection.

$$\begin{aligned} \mathbf{gcin} \text{ of } (v_{i+1}) &= 1, & i = 1, 2, \dots, 2n-2 \\ \mathbf{gcin} \text{ of } (v_1) &= 1. \\ \mathbf{gcin} \text{ of } (v_{2n}) &= \gcd \text{ of } \{ f_{cdpl}^*(v_{2n-1} v_{2n}), f_{cdpl}^*(v_1 v_{2n}) \} \\ &= \gcd \text{ of } \{ 12n^2 - 18n + 7, (2n-1)^3 \} \\ &= \gcd \text{ of } \{ 12n^2 - 18n + 7, (2n-1) \} \\ &= \gcd \text{ of } \{ (2n-1)(6n-6)+1, (2n-1) \} = 1 \end{aligned}$$

So,  $\mathbf{gcin}$  of each vertex of degree greater than one is 1.

Hence  $Y_n$ , admits cubic difference prime labeling. ■

**Theorem 2.5** Friendship graph<sup>3</sup>  $F_n$  admits cubic difference prime labeling.

Proof: Let  $G = F_n$  and let  $u, v_1, v_2, \dots, v_{2n}$  are the vertices of  $G$

Here  $|V(G)| = 2n+1$  and  $|E(G)| = 3n$

Define a function  $f : V \rightarrow \{0, 1, 2, 3, \dots, 2n\}$  by

$$f(v_i) = i, \quad i = 1, 2, \dots, 2n$$

$$f(u) = 0.$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$\begin{aligned} f_{cdpl}^*(u v_{2i-1}) &= (2i-1)^3, & i = 1, 2, \dots, n \\ f_{cdpl}^*(u v_{2i}) &= 8i^3, & i = 1, 2, \dots, n \\ f_{cdpl}^*(v_{2i-1} v_{2i}) &= 12i^2 - 6i + 1, & i = 1, 2, \dots, n \end{aligned}$$

Clearly  $f_{cdpl}^*$  is an injection.

$$\begin{aligned} \mathbf{gcin} \text{ of } (v_{2i-1}) &= \gcd \text{ of } \{ f_{cdpl}^*(u v_{2i-1}), f_{cdpl}^*(v_{2i-1} v_{2i}) \} \\ &= \gcd \text{ of } \{ (2i-1)^3, 12i^2 - 6i + 1 \} \\ &= \gcd \text{ of } \{ 2i-1, 6i(2i-1)+1 \} \\ &= 1, & i = 1, 2, \dots, n \\ \mathbf{gcin} \text{ of } (v_{2i}) &= \gcd \text{ of } \{ f_{cdpl}^*(u v_{2i}), f_{cdpl}^*(v_{2i-1} v_{2i}) \} \\ &= \gcd \text{ of } \{ (2i)^3, 12i^2 - 6i + 1 \} \end{aligned}$$

$$= \gcd \text{ of } \{2i, 2i(6i-3)+1\}$$

$$= 1, \quad i = 1,2,\dots,n$$

$$\mathbf{gcin} \text{ of } (u) = 1.$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence  $F_n$ , admits cubic difference prime labeling. ■

**Theorem 2.6** Umbrella graph<sup>3</sup>  $U(n,m)$  admits cubic difference prime labeling.

Proof: Let  $G = U(n,m)$  and let  $u, v_1, v_2, \dots, v_{m+n-1}$  are the vertices of  $G$

Here  $|V(G)| = m+n$  and  $|E(G)| = 2n+m-2$

Define a function  $f : V \rightarrow \{0,1,2,3,\dots,m+n-1\}$  by

$$f(v_i) = i, \quad i = 1,2,\dots,m+n-1$$

$$f(u) = 0.$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$f_{cdpl}^*(u v_i) = i^3, \quad i = 1,2,\dots,n$$

$$f_{cdpl}^*(u v_{n+1}) = (n+1)^3,$$

$$f_{cdpl}^*(v_i v_{i+1}) = 3i^2+3i+1, \quad i = 1,2,\dots,n-1.$$

$$f_{cdpl}^*(v_{n+i} v_{n+i+1}) = (n+i+1)^3 - (n+i)^3, \quad i = 1,2,\dots,m-2.$$

Clearly  $f_{cdpl}^*$  is an injection.

$$\begin{aligned} \mathbf{gcin} \text{ of } (v_n) &= \gcd \text{ of } \{f_{cdpl}^*(v_{n-1} v_n), f_{cdpl}^*(u v_n)\} \\ &= \gcd \text{ of } \{3n^2-3n+1, n^3\} \\ &= \gcd \text{ of } \{n, n(3n-3)+1\} \\ &= 1, \end{aligned}$$

$$\begin{aligned} \mathbf{gcin} \text{ of } (v_{n+1}) &= \gcd \text{ of } \{f_{cdpl}^*(v_{n+1} v_{n+2}), f_{cdpl}^*(u v_{n+1})\} \\ &= \gcd \text{ of } \{3n^2+9n+7, (n+1)^3\} \\ &= \gcd \text{ of } \{n+1, (n+1)(3n+6)+1\} \\ &= 1 \end{aligned}$$

$$\mathbf{gcin} \text{ of } (v_{i+1}) = 1, \quad i = 1,2,\dots,n-2$$

$$\mathbf{gcin} \text{ of } (v_{n+i+1}) = 1, \quad i = 1,2,\dots,m-3$$

$$\mathbf{gcin} \text{ of } (v_1) = 1.$$

$$\mathbf{gcin} \text{ of } (u) = 1.$$

So, *gcin* of each vertex of degree greater than one is 1.

Hence  $U(n,m)$ , admits cubic difference prime labeling. ■

**Theorem 2.7** Helm graph<sup>3</sup>  $H_n$  admits cubic difference prime labeling.

Proof: Let  $G = H_n$  and let  $v_1, v_2, \dots, v_{2n+1}$  are the vertices of  $G$

Here  $|V(G)| = 2n+1$  and  $|E(G)| = 3n$

Define a function  $f : V \rightarrow \{0,1,2,3, \dots, 2n\}$  by

$$f(v_i) = i-1, i = 1, 2, \dots, 2n+1$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$\begin{aligned} f_{cdpl}^*(v_i v_{i+1}) &= i^3 - (i-1)^3, & i = 1, 2, \dots, n \\ f_{cdpl}^*(v_{n+1} v_i) &= n^3 - (i-1)^3, & i = 1, 2, \dots, n-1 \\ f_{cdpl}^*(v_i v_{n+i+1}) &= (n+i)^3 - (i-1)^3, & i = 1, 2, \dots, n \\ f_{cdpl}^*(v_1 v_n) &= (n-1)^3 \end{aligned}$$

Clearly  $f_{cdpl}^*$  is an injection.

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= 1, & i = 1, 2, \dots, n-1 \\ \text{gcin of } (v_1) &= 1. \\ \text{gcin of } (v_{n+1}) &= 1. \end{aligned}$$

So, *gcin* of each vertex of degree greater than one is 1.

Hence  $H_n$ , admits cubic difference prime labeling. ■

**Theorem 2.8** Key graph<sup>3</sup>  $K(n,m)$  admits cubic difference prime labeling.

Proof: Let  $G = K(n,m)$  and let  $v_1, v_2, \dots, v_{n+2m-2}$  are the vertices of  $G$

Here  $|V(G)| = n+2m-2$  and  $|E(G)| = n+2m-2$

Define a function  $f : V \rightarrow \{0,1,2,3, \dots, n+2m-3\}$  by

$$f(v_i) = i-1, i = 1, 2, \dots, n+2m-2$$

Clearly  $f$  is a bijection.

For the vertex labeling  $f$ , the induced edge labeling  $f_{cdpl}^*$  is defined as follows

$$\begin{aligned} f_{cdpl}^*(v_i v_{i+1}) &= i^3 - (i-1)^3, & i = 1, 2, \dots, m+n-1 \\ f_{cdpl}^*(v_{n+i} v_{n+2m-i-1}) &= (n+2m-i-2)^3 - (n+i-1)^3, & i = 1, 2, \dots, m-2 \\ f_{cdpl}^*(v_1 v_n) &= (n-1)^3 \end{aligned}$$

Clearly  $f_{cdpl}^*$  is an injection.

$$gcin \text{ of } (v_{i+1}) = 1, \quad i = 1, 2, \dots, m+n-2$$

$$gcin \text{ of } (v_1) = 1.$$

So,  $gcin$  of each vertex of degree greater than one is 1.

Hence  $K(n,m)$ , admits cubic difference prime labeling. ■

## REFERENCES

1. F Harary. Graph Theory. 2<sup>nd</sup> ed. Addison-Wesley: Massachuset: 1972.
  2. Apostol. Tom M. Introduction to Analytic Number Theory. 2<sup>nd</sup> ed. Narosa: New Delhi: 1998.
  3. Joseph A Gallian. A Dynamic Survey of Graph Labeling. The Electronic Journal of Combinatorics 2016; #DS6: 1 – 408.
  4. Mathew Varkey T. K. Some Graph Theoretic Generations Associated with Graph Labeling. PhD Thesis. University of Kerala: Kariavattam: 2000.
-