

International Journal of Scientific Research and Reviews

Isolation and screening of yeast from sources such as grapes, corn, curd, water sample 1 & 2

Govind P. Misale¹, S.E.Neelgund^{2*}, Vijendra Singh³, S.V. Patil⁴

¹Associate Vice President (Ethanol), ²Professor, ³Executive Director, ⁴Head Alcohol Technology
¹Renuka sugars limited, Belagavi.

²Department of Biochemistry, Kuvempu University, Jnanasahyadri, Shankaraghatta, Road leading to
Kuvempu University, Shimoga, Karnataka 577451

³ShreeRenuka Sugars limited, Mumbai.

⁴Vasant Dada Sugar Institute Pune.

ABSTRACT

Ethanol fermentation, is a organic process where glucose, fructose and sucrose are transformed into mobile energy and thereby produce ethanol and carbondioxide as metabolic waste products. Considering that yeast does this in absence of oxygen, alcoholic fermentation is viewed as anaerobic procedure. The time period alcohol is most usually used to consult the fermented made from sugars. Such products will also be obtained from one of a sources like molasses, fruits, grains etc. There are countless stories on alcohol creation from fruits such as apple, pear, banana, jack fruit, mango, plum, litchi, and strawberry. The environment impact of the distillery industry could be reduced to extents which would be significant in making them Eco-friendly. By discovery of such novel organsims the potential diversification of nature would be realised. Thus the present work contributes in a large way to achieve “sustainable development”.

The present study was done on sources such as corn, curd, grapes, tap water sample. Isolation and screening of yeasts was done on malt extract glucose yeast extract peptone (MGYP) media. The method utilized in research instructed that high alcohol produced immediately from grape, that allows you to be priceless for ethanol creation in many locations. Outcome indicate that grape pattern yeast used to be determined as perfect ethanol producing yeast after optimization as easiest ethanol producing yeast after optimization in jaggery broth. Improved production of ethanol, temperature, nutrient and pH on grape yeast have to be optimized.

The study indicates that grape yeast used to determined as a perfect ethanol producing yeast after optimization. utcome indicate that grape pattern yeast used to be determined as perfect ethanol producing. For better production of ethanol, temperature, nutrient and pH on grape yeast have to be optimized.

KEYWORD: Ethanol, fermentation, isolation, MGYP media.

***Corresponding author:**

Prof. S. E. Neelgund

Department of Biochemistry,

Kuvempu University, Shimoga.

Email: neelgund@gmail.com

INTRODUCTION:

Throughout the 17th century alcohol was once not used for the consumption but additionally as a constituent of clinical drugs and manufacture of paint pigments and other industrial chemical substances. It used to be initiated in the 19th century and the trade became an industry with gigantic construction.¹Ethanol is an essential industrial chemical with rising competencies as a bio-gasoline to exchange fossil fuels. Ethanol is produced by chemical synthesis utilizing hydration of ethylene. Biosynthesis is the fermentation procedure and yeast mono-saccharides as a carbon source and then converts to ethanol through glycolysis beneath anaerobic conditions.^{2,3}

The priceless physiological of yeast have led to their use within the discipline of biotechnology, fermentation of sugars by yeast being the oldest and greatest software of this science. Many varieties of yeasts are used for making many meals: baker's yeast in bread creation; brewer's yeast in beer fermentation; yeast in wine fermentation and for xylitol creation.⁴A worldwide interest within the utilization of bio-ethanol as a vigour source has motivated reports on the cost and effectively of business strategies for ethanol construction. Severe research has been implemented for acquiring effective fermentative organisms, low-priced fermentation substrates and top of the line environmental conditions for fermentation to occur.⁵Frequently, ethanol production is usually comprehensive by using microbial alteration of carbohydrates present in agricultural yields. As, few yeast traces were observed to own considerable traits for ethanol production.⁶

Carbohydrate wealthy raw materials suitable for ethanol construction are labeled into three organizations of agricultural products: sugars, starch and lignocellulose. The primary uncooked fabric workforce is from sugar like sugar-beet, sugarcane and molasses. The 2nd product is starch from crops includes cassava, cereals and potatoes. The last staff is lignocellulose which duvet waste substances from the agriculture crops reminiscent of rice, straw, corn and sugarcane waste. There are a number of wild loved ones of grapes recounted in the literature. However, the yeast is used best in grape.^{7,8}In the present study, several local samples of corn, curd, grapes, water sample 1 & 2 were analyzed for isolation and subsequent characterization of yeast strains, which may further be utilized in alcohol production.

MATERIALS AND METHOD:

Isolation and screening of yeasts:

Yeasts had been remoted from the sources such as corn, curd, grapes, ap water sample and paneer.

1. Sample was once taken to cleaned laminar air flow cabinet (Figure 1)

Figure I: Slant preparation and culture inoculation in media



2. Underneath aseptic, 1 gm of homogenised sample used to be brought to 20 ml of MGYB broth.
3. Sample used to be incubated at 26⁰C for 2 days.
4. Corn sample was washed in sterile D/W after which beaten in sterilised pestle and mortal. Then roughly 22 gm of overwhelmed sample was inoculated in 10 ml MGYB broth aseptically.
5. Curd; 1 ml homogenised curd sample was inoculated in 9 ml of MGYB broth aseptically.
6. Grapes; Grape sample was washed utilizing sterile D/W and crushed. Approximately 1 g of sample used to be aseptically inoculated in 10 ml of sterile MGYB broth (Figure 2).

Figure II: Developed slant of micro organisms.



7. Water sample 1 (dirty water from Hotel): roughly 1 ml of water sample was aseptically inoculated in 9 ml of sterile MGYB broth.
8. Water sample 2 (dirty water from station): roughly 1 ml of dirty water sample was aseptically inoculated in 9 ml of sterile MGYB broth.⁹
9. Starter culture was prepared from Grapes (Figure 3).

Figure III: Lab scale preparation of starter and batch culture.



RESULT AND DISCUSSION:

The striking pattern on MGY media and have been picked single colony to determine the morphology attribute. All samples showed growth in the type of turbidity. Table I indicates morphology characteristics of yeast isolates.

Table I: Morphology characteristics of yeast isolates.

S.No	Name of sample	Morphology observed
1	Grapes	round shaped violet colored cells
2	Corn	rod shaped violet colored cells
3	Curd	round shaped violet colored cells
4	Water-1	round shaped violet colored cells
5	Water-2	oval shaped violet colored cells

After morphology characteristics of yeast isolates. After incubation isolates were got, colony characters have been checked and stain had been done. Table II confirmed results of colony traits of yeast isolates.

Table II: Colony characteristics of yeast isolates.

Sl.No	Colony Characteristics	Grape	Curd	Water -1	Water -2
1	Size	0.8 mm	0.4 mm	0.7 mm	0.5 mm
2	Shape	Spherical	Spherical	spherical	spherical
3	Color	White	White	White	White
4	Margin	Entire	Entire	Entire	Entire
5	Elevation	Concave	Concave	Concave	Concave
6	Opacity	Opaque	Opaque	Opaque	Opaque
7	Consistency	Moisture	Dry	Dry	Dry

All colonies were exclusive whilst shapes of colonies had been spherical with white color. Margin of all colonies have been complete additionally colonies elevation had been concave. Imperviousness of all colonies were showed opaque with dry steadiness excluding colony of grape that was moisture.

Table III confirmed outcome of alcohol influence of alcohol produced by yeast isolates.

Table III: Percent of alcohol produced by yeast isolates.

Sl.No	Name of sample	Specific Gravity	A.B.W	A.B.V
1	Grape	1.084	5.98%	7.21%
2	Corn	1.035	5.2%	6.21%
3	Curd	1.042	4.67%	4.52%
4	Water- 1	1.018	3.98%	5.21%
5	Water- 2	1.015	2.58%	3.98%

NOTE: A.B.W = alcohol by weight and A.B.V= alcohol by volume.

The yeast removed from curd and corn have been 4.6% alcohol through quantity (a.b.v), 5.25% (a.b.v) respectively. Grape yeast isolate used to be 6.03% (a.b.v) alcohol after fermentation. Among the different sample taken of yeast isolates, grapes gave the large and good quality alcohol production.

Ethanol fermentation is responsible for the rising of bread dough. Yeast organisms consume sugars in the dough and produce ethanol and carbon dioxide as waste products. The carbon dioxide types bubbles within the dough, increasing it into some thing of foam. Practically the entire ethanol evaporates from the dough when the bread is baked.

All alcoholic beverages, together with those produced with the aid of carbonic maceration are produced via ethanol fermentation by way of yeast. Wine and brandy are produced by way of fermentation of the typical sugars gift in fruits, especially grapes. The ferment potential of grape is involving cultivar, atmosphere, fertility of soil, stipulations of maturity at harvest and medication of grapes within the winery.^{10,11}

Within the gift study, curd, corn and grape were yeast removed however best yeast gave 6.03% (a.b.v) in grape alcohol after fermentation. This isolate discovered that the yeast are quite often huge in size and so they may be circular or rod shaped. Grape yeast used to be round in shape whose colonies have been dry and their measurement is about 0.7 mm. The method utilized in research instructed that high alcohol produced immediately from grape, that allows you to be priceless for ethanol creation in many locations. Outcome indicate that grape pattern yeast used to be determined as perfect ethanol producing yeast after optimization as easiest ethanol producing yeast after optimization in jaggery broth. Better production of ethanol, temperature, nutrient and pH on grape yeast have to be optimized.

CONCLUSION:

The study indicates that grape yeast used to be determined as a perfect ethanol producing yeast after optimization. Outcome indicates that grape yeast used to be determined as perfect ethanol producing. For better production of ethanol, temperature, nutrient and pH on grape yeast have to be optimized.

REFERENCES:

1. Tanimura A, Kikukawa M, Yamaguchi S, Kishino S, Ogawa J, Shima J. Direct ethanol production from starch using a natural isolate, *Scheffersomyces castellii*: toward consolidated bioprocessing. *Scientific reports*. 2015;5:9593.
2. Oke MA, Annur MS, Simarani K. Mixed feedstock approach to lignocellulosic ethanol production—prospects and limitations. *BioEnergy Research*. 2016;9(4):1189-203.
3. Ruyters S, Mukherjee V, Verstrepen KJ, Thevelein JM, Willems KA, Lievens B. Assessing the potential of wild yeasts for bioethanol production. *Journal of industrial microbiology & biotechnology*. 2015;42(1):39-48.
4. Baeyens J, Kang Q, Appels L, Dewil R, Lv Y, Tan T. Challenges and opportunities in improving the production of bio-ethanol. *Progress in Energy and Combustion Science*. 2015;47:60-88.
5. C. J. Panchal, I. Peacock and G. G. Stewart, *Biotechnol. Lett.* 1982;4, 639-644.
6. Lalou S, Capece A, Mantzouridou FT, Romano P, Tsimidou MZ. Implementing principles of traditional concentrated grape must fermentation to the production of new generation balsamic vinegars. Starter selection and effectiveness. *Journal of food science and technology*. 2016;53(9):3424-36.
7. Morris JR, Main G and Threfall R. Fermentations: Problems, solutions and prevention. *Vitic. Enol. Sci.* 1996; 51: 210-213.
8. Lalou S, Capece A, Mantzouridou FT, Romano P, Tsimidou MZ. Implementing principles of traditional concentrated grape must fermentation to the production of new generation balsamic vinegars. Starter selection and effectiveness. *Journal of food science and technology*. 2016;53(9):3424-36.
9. Goold HD, Kroukamp H, Williams TC, Paulsen IT, Varela C, Pretorius IS. Yeast's balancing act between ethanol and glycerol production in low-alcohol wines. *Microbial biotechnology*. 2017;10(2):264-78.

10. Corbin KR, Hsieh YS, Betts NS, Byrt CS, Henderson M, Stork J, DeBolt S, Fincher GB, Burton RA. Grape marc as a source of carbohydrates for bioethanol: chemical composition, pre-treatment and saccharification. *Bioresource technology*. 2015;193:76-83.
11. Lalou S, Capece A, Mantzouridou FT, Romano P, Tsimidou MZ. Implementing principles of traditional concentrated grape must fermentation to the production of new generation balsamic vinegars. Starter selection and effectiveness. *Journal of food science and technology*. 2016;53(9):3424-36.