

International Journal of Scientific Research and Reviews

Membrane Perturbations in ageing Jamun (*Syzygium cuminii*) Seeds

Jyoti Bakshi*

Department of Botany, St. Thomas College, Bhilai- 490006, C.G. INDIA

e-mail taranjyot2007@rediffmail.com

ABSTRACT

Recalcitrant seeds of *Syzygium cuminii* (Jamun) exhibit 100% viability upto 3 dh after maturity. The rapid loss of viability after 3dh is associated with reduction in moisture content below lowest safe moisture content (47.71% MC). Seed became nonviable after slow drying to 20% MC within a very short period of 27 dh during storage at ambient condition. Decline of percent germination in dehydrating Jamun seeds was strongly and negatively correlated with increased leakage loss of electrolyte and MDA accumulation during drying. Jamun seed of low vigor or quality can be recognized as increased leakage loss and MDA content. Thus the result suggests that Jamun seeds deterioration during natural ageing is closely related to lipid peroxidation which results in the accumulation of MDA.

KEY WORDS: Recalcitrant seeds, *Syzygium cuminii*, Membrane perturbation, Lipid peroxidation.

ABBREVIATIONS: Moisture content (MC), Malondialdehyde (MDA), CWC (Critical Water Content), dh (days after harvesting)

***Corresponding Author**

Dr. Jyoti Bakshi

Department of Botany,

St. Thomas College,

Ruabandha, Bhilai-490006, C.G., India

E-mail taranjyot2007@rediffmail.com,

Mobile No. 9993198870

1. INTRODUCTION

Jamun (*Syzygium cuminii*) seeds are categorized as tropical recalcitrant as they are shed with high water content ($0.93 \text{ g H}_2\text{O g}^{-1} \text{ DM}$), have a short viability (27 days), high CWC ($0.86 \text{ g H}_2\text{O g}^{-1} \text{ DM}$), and desiccation sensitive. In most if not all, membrane are the primary target of desiccation injury and play a key role in maintaining seed viability and vigor^{1,2,3,4,5}. Seed imbibition invariably accompanies solute leakage during the process of membrane reorganization following rehydration in the dry seeds. The rate of leakage is directly correlated with the cell damage and repair in response to ageing^{6, 7}. Enhanced electrolyte leakage loss estimated in the damaged seeds of *Lotus corniculatus*⁸, *Artocarpus heterophyllus*⁹, *Euterpe edulis*¹⁰, *Fagus sylvatica*¹¹ and *Hopea ponga*¹² was considered an indicator of loss in membrane integrity as a result of dehydration and is accountable for reduced germination and vigor of seeds. Initial viability loss has been associated with loss of cellular constituents mainly due to oxidation of protein and lipid; major components of membrane^{13, 14, 15}. The lipid peroxidation plays an important role in initiating and mediating ageing process as the lipids are major component of semi permeable membrane and probably the first easy target of free radicals. During seed imbibition, leakage loss of solutes as a result of altered membrane integrity is closely associated with the accumulation of malondialdehyde (MDA), a final product of lipid peroxidation^{16, 17, 18}.

The objective of this research was to follow the changes in membrane integrity in desiccating recalcitrant Jamun seeds. The relative significance and contribution of axis and cotyledon was also determined during loss of viability by monitoring the amount of MDA formation in these tissues.

2. MATERIALS AND METHODS

2.1 Site of Fruit Collection

Jamun (*Syzygium cuminii* L.) fruits were collected from avenue trees nearby Bhilai (Chhattisgarh), India. Nearly 80 plus trees were marked for collection of fruits.

2.2 Fruit Collection

Freshly mature fruits of Jamun were plucked manually during June-July. The collected fruits were transported to the laboratory within one hour of collection. The healthy fruits were sorted out. The fruit is a drupe, variable in size, oblong or sub-globose, crowned with a persistent truncated first pink, then black with pink juicy mesocarp.

2.3 Seed Extraction

Seeds were extracted by rubbing the fruit with sand to remove pulp. Three replicates of 50 seeds each were used for determination of moisture content. Remaining pulp free seeds were then washed thoroughly with tap water to remove traces of pulp and allowed to air-dry to initial moisture contents. Each fruit contains one seed which is 1-2 cm long, oblong in shape green or brown in colour.

2.4 Seed Drying and Storage

The mature Jamun seeds collected were for conducting experiments related to slow (natural drying at laboratory conditions RH 30% and Temperature (27-30°C). The seed was subjected to slow drying by spreading them in one layer in a perforated basket at ambient conditions (27-30°C at RH 30%).

2.5 Moisture Content and Estimation of Water Content

Moisture contents of slow dried seeds were determined by oven drying the seeds for 72 hours at 103°C¹⁹. Five replicates with 10 seeds each were used to determine the moisture content, on a fresh weight basis. The seed moisture content was determined by the following formula and was expressed in percentage:

$$\text{Seed Fresh Weight} - \text{Seed Dry Weight}$$

$$\% \text{ Moisture Content} = \frac{\text{Seed Fresh Weight} - \text{Seed Dry Weight}}{\text{Seed Fresh Weight}} \times 100$$

$$\text{Seed Fresh Weight}$$

Water content of slow dried seeds was determined by the method given by²⁰.

$$\text{Seed Fresh Weight} - \text{Seed Dry Weight}$$

$$\text{Water Content} = \frac{\text{Seed Fresh Weight} - \text{Seed Dry Weight}}{\text{Seed Dry Weight}} \times \text{g H}_2\text{O g}^{-1}\text{DM}$$

$$\text{Seed Dry Weight}$$

2.6 Leachate Conductivity

Solute leakage loss was measured in terms of electrolytic conductivity by²¹. The leachates were collected after 24 hours imbibition of seed in distilled water and the specific conductivity was estimated using a conductivity meter (Elico). Results were expressed as mS seed⁻¹.

2.7 Lipid Peroxidation

Lipid peroxidation was measured as the concentration of thiobarbituric acid (TBA) reactive products equated with malondialdehyde (MDA), the end product of lipid peroxidation. The method given by²² was followed to determine lipid peroxidation. To 0.05 g of axis and 0.1 g of cotyledons add 3 ml of 0.5% TBA prepared in 20% TCA and 1 ml of BHT. The reaction mixture was incubated in boiling water bath (97°C) for 30 minutes to permit formation of pink colour of the product when the lipid peroxidation products (i.e. MDA) react with TBA. The reaction tubes were plunged into the pail of ice to stop the reaction. The reaction mixture was then centrifuged at 10,000 rpm for 30 minutes, to precipitate all the suspending particles. The clear supernatant was collected and the absorbance was recorded at 440, 532, and 660 nm using spectrophotometer. The following formula given by²³ was used to calculate Lipid peroxidation and expressed as ΔA_{540} /gm FW of axis or cotyledon:

$$[A_{532} - A_{660}] - [(A_{440} - A_{660}) \times 0.0571] \times 10$$

$$\text{Lipid peroxidation} = \frac{\text{-----}}{157000}$$

3. RESULTS

3.1 Desiccation and Loss of Viability

Absolute germination (100%) was recorded in Jamun seeds during slow drying of seed water content from 0.93 to 0.86 g H₂O g⁻¹ DM. Thereafter, with quick decline in water content, a corresponding decline in %germination was also registered, i.e. 95% germination was recorded at water content 0.81 g H₂O g⁻¹ DM and 80% at water content 0.42 g H₂O g⁻¹ DM. Complete loss of germination was noticed at the time when seed was desiccated to 0.23 g H₂O g⁻¹ DM. Along with the per cent germination, the germination index of the seed also decreased considerably with loss of water content during seed storage.




Figure 1. The Decline in Percentage of Germination and Germination Index of *Syzygium cuminii* Seeds During Storage (Slow Drying) Under Ambient Conditions.

3.2 Electrolyte Leakage

The specific conductivity of the seed leachates estimated in the imbibing seeds showed gradual promotion with the loss of seed water content during slow drying. Drying of seeds initially from 0.93 to 0.26 g H₂O g⁻¹ DM, resulted in rapid increase in electrolyte leakage from 0.12 to 0.44 mS seed⁻¹ but further loss of seed water content to 0.23 g H₂O g⁻¹ DM did not much alter the specific conductivity




Figure 2. Loss of Electrolytes with Decline in Water Content of *Syzygium cuminii* Seeds During Storage Under Ambient Conditions (Expressed as Specific Conductivity of Leachates).

3.3 Lipid Peroxidation

The lipid peroxidation product MDA (also called as Thiobarbituric Acid Reactive Substances-TBARS) increased in the embryonic axis and cotyledon of Jamun seeds with the decrease in water content during slow drying at ambient storage condition. The TBARS levels remained significantly low when the water content of the seed declined from 0.93 to 0.86 g H₂O g⁻¹ DM during slow drying both in axis and cotyledon. Slight promotion in MDA content from 2.22 to 2.62 A₅₄₀ g⁻¹ FW was observed in the axis, whereas it increased sharply in the cotyledon from 2.53 to 5.73 A₅₄₀ g⁻¹ FW (more than 2-fold) in response to loss of seed water content from 0.86 to 0.81 g H₂O g⁻¹ DM. Further, slow drying of seeds from 0.81 to 0.23 g H₂O g⁻¹ DM resulted in sharp increase in MDA levels both in axis (2.62 to 4.20 A₅₄₀ g⁻¹ FW) and cotyledon (5.73 to 6.06 A₅₄₀ g⁻¹ FW). In general, the levels of MDA were comparatively higher in the cotyledons than the axis throughout the period of analysis.




Figure 3. Accumulation of TBA Reactive Substances in the Embryonic Axis and Cotyledon of *Syzygium cuminii* Seeds with Decline in Water Content during Slow Drying.

4. DISCUSSION

The present study has confirmed that drying of seeds during maturation phase of seed development and later during storage generally leads to structural and functional alteration in membrane and its components. The integrity of membranes must be repaired during early phase of imbibition^{24, 25}. The reorganization and permeability of the membrane is monitored by measuring the specific conductivity of solutes leaked from a dry seed upon rehydration or imbibition. Dehydration, induced loss of percent germination in recalcitrant Jamun seeds was accompanied by membrane deterioration as indicated by

increased rates of, 1-electrolyte leakage from imbibing seeds and, 2-MDA accumulation. Membrane damage as a result of increased peroxidation of membrane phospholipids caused loss of selective permeability²⁶ that, in turn, is responsible for increased leakage loss. Increased leakage that is invariably associated with ageing is due to enhanced membrane permeability^{27, 28}. In the present study, loss of percent germination in dehydrating Jamun seeds was strongly and negatively correlated with increased leakage loss of electrolyte and MDA accumulation during slow drying.

Loss of germinability, viability and vigor during storage of orthodox and recalcitrant seeds is frequently related to potential damage of membranes due to uncontrolled peroxidation of membrane lipid^{29, 30, 31, 32}. Lipid peroxidation estimated by recording MDA content, an end and volatile product of lipid peroxidation, in the cotyledon and axis indicated promotion in MDA accumulation in response to desiccation of Jamun seeds. Slow drying of fresh Jamun seeds from 0.93 to 0.23 g H₂O g⁻¹ DM resulted in net increase in the estimated MDA by 2.6-fold in cotyledon and 2.2-fold in axis. Differential response of cotyledon and axis in terms of lipid peroxidation to drying perhaps implies the differential sensitivity of these tissues.

Taking together the results of leakage loss and MDA in dehydrating Jamun seeds, it is concluded that constant drying of recalcitrant Jamun seeds activated the autocatalytic reaction of lipid peroxidation resulting in the accumulation of MDA. MDA accumulation, an indicator of membrane damage, is consequently responsible for membrane perturbations in the dehydrating Jamun seeds that exhibited severe leakage loss. Leaky membranes due to extensive lipid peroxidation in desiccating Jamun seeds may be a prime cause of reduced germination and vigor (GI). Enhanced MDA contents mediated by organic radicals and peroxides that are produced during lipid peroxidation of polyunsaturated fatty acids (PUFA) are considered to be one of the likely explanations for lipid peroxidation^{33,34,35,36} in desiccating seeds during germination.

5. ACKNOWLEDGEMENTS

The author wish to acknowledge thanks to the Head, School of Life Sciences, Pt. Ravishankar Shukla University, Raipur for providing the necessary facilities.

6. REFERENCES

1. Ferguson M, Tekrony DM and Egli DE. Changes during early seed and axes deterioration, II Lipids. Crop Science 1990; 30: 179-182.

2. Thornton JM, Powell AA and Mathews S. Investigation of the relationship between seed leachate conductivity and germination of *Brassica* seeds. *Annals of Applied Biology* 1990; 117: 129-135.
3. Chaitanya KSK and Naithani SC. Role of superoxide, lipid peroxidation and superoxide dismutase in membrane perturbation during loss of viability in seeds of (*Shorea robusta* Gaertn. f.). *New Phytologist* 1994; 126: 623-627.
4. Chaitanya KSK and Naithani SC. Kinetin-mediated prolongation of viability in recalcitrant sal (*Shorea robusta* Gaertn. f.) seeds at low temperature: Role of kinetin in delaying membrane deterioration during desiccation-induced injury. *Journal of Plant Growth Regulation* 1998; 17: 63-69.
5. Al-Maskri AY, Khan MM, Khan IA et al. Effect of accelerated ageing on viability, vigor (RGR), lipid peroxidation and leakage in carrot (*Daucus carota* L.) seeds. *International Journal of Agriculture and Biology* 2003; 5(4): 580-584.
6. Larson LA. Effect of soaking pea seeds with or without seed coats on seedling growth. *Plant Physiology* 1968; 43: 255-259.
7. Simon EW. "Membranes in dry and imbibing seeds". In: Crowe JH and Clegg JS. (eds) *Dry Biological Systems*, Academic Press: New York; 1978: 205-224.
8. McKersie BD and Tomes D. Effects of dehydration treatments on germination, seedling vigor and cytoplasmic leakage of wild oats and birds foot trefod. *Canadian Journal of Botany* 1980; 58: 471-476.
9. Smith JW, Pammenter NW, Berjak P et al. The effect of two drying rates on the desiccation tolerance of embryonic axes of recalcitrant jackfruit (*Artocarpus heterophyllus* Lamk.) seeds. *Annals of Botany* 2001; 88: 653-664.
10. Panza V, Lainez V, MalDonado S et al. Effects of desiccation on *Euterpe edulis* Martius seeds. *Biocell* 2007; 31(3): 383-390.
11. Pukacka S and Ratajczak E. Production and scavenging of reactive oxygen species in *Fagus sylvatica* seeds during storage at varied temperature and humidity. *Journal of Plant Physiology* 2005; 162: 873-885.
12. Sukesh SH and Chandrashekhar KR. Biochemical changes during storage of seeds of *Hopea ponga* (Dennst.) Mabberly: an academic species of Western Ghats. *Research Journal of Seed Science* 2011; 4: 106-116.

13. Vertucci CW and Farrant JM. "Acquisition and loss of desiccation tolerance". In: Kigel J and Galili G (eds) Seed Development and Germination. Marcel Dekker, Inc. New York, Basel, Hong Kong; 1995: 237-271.
14. Sacandé M, Hoekstra FA, van Aelst AC et al. Is oxidative stress involved in the loss of neem (*Azadirachta indica*) seed viability? Seed Science Research 2000; 10: 381-392.
15. Varghese B and Naithani SC. Oxidative metabolism related to changes in cryogenically stored neem (*Azadirachta indica* A. Juss.) seeds. Journal of Plant Physiology 2008; 165: 755-765.
16. Berjak P and Pammenter NW. From *Avicennia* to *Zizania*: seed recalcitrance in perspective. Annals of Botany 2008; 101: 213-228.
17. Li YI, Jian-Jun QU, Wei-Min ZHANG et al. Impact of ultra-dry storage on vigor capacity and antioxidant enzyme activities in seed of *Ammopiptanthus mongolicus*. Botanical Studies 2010; 51: 465-472.
18. Parkhey S, Naithani SC and Sahu KK. ROS production and lipid catabolism in desiccating *Shorea robusta* seeds during ageing. Plant Physiology and Biochemistry 2012; 57: 261-267.
19. ISTA. International Rules for Seed Testing. Seed Science and Technology, International Seed Testing Association, Zurich, Switzerland, 1993; 31: 1-288.
20. ISTA. International Rules for Seed Testing. Seed Science and Technology, International Seed Testing Association, Zurich, Switzerland, 1996; 24.
21. Varghese B and Naithani SC. Desiccation-induced changes in lipid peroxidation, superoxide level and antioxidant enzymes activity in neem (*Azadirachta indica* A. Juss.) seeds. Acta Physiologia Plantarum 2002; 24: 79-87.
22. Heath RL and Packer L. Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. Archives of Biochemistry and Biophysics 1968; 125: 189-198.
23. Hodges DM, DeLong JM, Forney CF et al. Improving the TBA reactive substance assay for estimating lipid peroxidation in plant tissue containing anthocyanin and other interfering compounds. Planta 1999; 604-611.
24. Simon EW. Phospholipids and plant membrane permeability. New Phytologist 1974; 73: 377-420.
25. Senaratna T, Gusse JF and McKersie BD. Age induced changes in cellular membranes of imbibed soybean seed axes. Plant Physiology 1988; 73: 85-90.

26. Tatipata A. Effect of seed moisture content packaging and storage period on mitochondria inner membrane of soybean seed. *Journal of Agricultural Technology* 2009; 5(1): 51-64.
27. Copeland LO and McDonald MB. "Principles of Seed Science and Technology". Burgess Publishing Company, Minneapolis, Minnesota, USA; 1985.
28. McDonald MB. Seed deterioration physiology, repair and assessment. *Seed Science and Technology* 1999; 27: 177-237.
29. Fletcher BL, Dillard CJ and Tappel AL. Measurement of fluorescent lipid peroxidation products in biological systems and tissues. *Analytical Biochemistry* 1973; 52: 1-9.
30. Dhindsa RS and Matowe W. Drought tolerance in two mosses correlated with enzymatic defence against lipid peroxidation. *Journal of Experimental Botany*; 1981; 32: 79-91.
31. Kumar GNM and Knowles NR. Changes in lipid peroxidation and lipolytic and free-radical scavenging enzyme activities during ageing and sprouting of potato (*Solanum tuberosum*) seed tubers. *Plant Physiology* 1993; 102: 115-124.
32. Smith MT and Berjak P. "Deteriorative changes associated with the loss of viability of stored desiccation-tolerant and sensitive seeds". In: Kigel J and Galili G (eds) *Seed Development and Germination*, Marcel Dekker In: New York; 1995: 701-704.
33. Roubal WT. Trapped radicals in dry lipid-protein systems undergoing oxidations. *Journal of the American Oil Chemists Society* 1970; 47: 141-144.
34. Funes J and Karel M. Free radical polymerization and lipid binding of lysozyme reacted with peroxidizing linoleic acid. *Lipid* 1981; 16(5): 347-350.
35. Pauls KP and Thompson JE. Effect of *in vitro* treatment with ozone on the physical and chemical properties of membrane. *Physiologia Plantarum* 1981; 53: 255-262.
36. Gutteridge JM and Halliwell B. The measurement and mechanism of lipid peroxidation in biological systems. *Trends in Biological Sciences* 1990; 15: 129-135.