

International Journal of Scientific Research and Reviews

Fixed Point Theorem for Quadruple Self-Mappings in Digital Metric Space

Chauhan M.S.¹, Shrivastava Rajesh², Verma Rupali^{3*}, Kabir Qazi Aftab⁴

^{1,3} Department of Mathematics, Institute for Excellence in Higher Education, Bhopal, M P India.

² Department of Mathematics, Govt. Science and Commerce College Benazir, Bhopal, M P India.

⁴ Department of Mathematics, Saifia Science College Bhopal, M P India.

Email: - ¹dr.msc@rediffmail.com, ²rajeshraju0101@rediffmail.com, ³rupali.varma1989@gmail.com,
⁴qaziaftabkabir@gmail.com.

ABSTRACT:

In this paper, we have established a common fixed point theorem for quadruple self mappings satisfying the continuous function in digital metric space. Our result is the extension of the results obtained by Jain for complete digital metric spaces. We have concluded an example to support our main result.

KEYWORDS: Common Fixed point, Digital metric space, Quadruple mappings, continuous functions.

***Corresponding author**

Rupali Verma

Research Scholar, Department of Mathematics,

Institute for Excellence in Higher Education(IEHE)

Bhopal-462016, M.P. INDIA

Email id- rupali.varma1989@gmail.com Mobile No. 8358906402

INTRODUCTION:

It was the dawn of the fixed point theory when in 1912 Brouwer proved a fixed point result for continuous self maps on a closed ball then later in 1922, Banach¹ gave a very useful result in form of Banach Contraction Principle. At first it was Rosenfeld² who considered digital topology as a tool to study digital images. The digital version of the topological concepts was produced by Boxer³ and later he studied digital continuous functions⁴. Ege and Karaca⁵ established relative and reduced Lefschetz fixed point theorem for digital images, the notion of a digital metric space was also proposed by them. Further, the famous Banach Contraction Principle for digital images was proved by Ege⁶. In digital topology the notion of digital continuity was developed by Rosenfeld⁷ to study 2D and 3D digital images. Further in 2013, Ege and Karaca⁵ described the digital continuous functions.

Fixed point theory has applications in field of mathematics, computer science, engineering, game theory, fuzzy theory, image processing and so forth. In metric space it begins with the Banach fixed-point theorem providing a constructive method to find fixed points and is an essential tool to find solution of some problems in mathematics and engineering and consequently has been generalized in many ways. Till now, several developments have occurred in this area. In 1976 a major shift in the arena of fixed point theory came when Jungck^{8,9,10}, defined the concept of commutative and compatible maps and proved the common fixed point results for such maps. Later on, Sumitra et.al¹¹ proved common fixed point theorems for compatible map in digital metric space with its applications. Certain alterations of commutativity and compatibility can also be found in references^{9,10,12}. The notion of compatible mappings of type(R) in digital metric space was introduced by Jain¹³. In this paper we have generalized and extended the results obtained by Jain¹³ for quadruple mappings in complete digital metric space.

2. DEFINITIONS:

Definition 2.1.⁴ Let $(X, \rho_0) \subset \mathbb{Z}^{n_0}, (Y, \rho_1) \subset \mathbb{Z}^{n_1}$ be digital images and $f: X \rightarrow Y$ be a function.

- (i) If for every ρ_0 -connected subset U of X , $f(U)$ is a ρ_1 -connected subset of Y , then f is said to be (ρ_0, ρ_1) -continuous.
- (ii) f is (ρ_0, ρ_1) -continuous for every ρ_0 -adjacent points $\{x_0, x_1\}$ of X , either $f(x_0) = f(x_1)$ or $f(x_0)$ and $f(x_1)$ are ρ_1 -adjacent in Y .
- (iii) If f is (ρ_0, ρ_1) -continuous, bijective and f^{-1} is (ρ_0, ρ_1) -continuous, then f is called (ρ_0, ρ_1) -isomorphism and denoted by $\cong (\rho_0, \rho_1) Y$.

Proposition 2.2.⁶ Every digital contraction map $T: (X, d, \rho) \rightarrow (X, d, \rho)$ is digitally continuous.

Definition 2.3.⁵ Let $X \subseteq Z^n$ and (X, d, ρ) be a digital metric space. Then there does not exist a sequence $\{x_n\}$ of distinct elements in X , such that

$$d(x_{m+1}, x_m) < d(x_m, x_{m-1}) \quad \text{for } m = 1, 2, 3, \dots$$

Definition 2.4.⁴ Suppose that (X, d, ρ) is a digital metric space and $P, Q: X \rightarrow X$, and be two self-maps defined on X . then P and Q are compatible if

$$d(PQx, QPx) \leq d(Px, Qx) \text{ for all } x \in X.$$

Definition 2.5.¹⁴ Let (P, Q) be a pair of self-mappings on a metric space (X, d) and $x \in X$. then P is called (G, O) - continuous at X if $PX_n \rightarrow PX$, for every sequence $\{x_n\} \subset X$ with $Qx_n \uparrow Qx$ (resp. $Qx_n \downarrow Qx, Qx_n \uparrow \downarrow Qx$). Moreover, P is called (G, O) - continuous (resp. $((G, \bar{O}) - \text{continuous}, (G, O)$ - continuous) at every point of $x \in X$.

Remark 2.6.¹⁴ In digital metric space, G -continuity $\Rightarrow (G, O) - \text{continuity} \Rightarrow (G, \bar{O}) - \text{continuity}$ (as well as $(G, \underline{O}) - \text{Continuity}$).

Proposition 2.7.⁶ Let (X, d, ρ) is a digital metric space. A sequence $\{x_n\}$ of points of a digital metric space (X, d, ρ) is

- (i) A Cauchy sequence if and only if there is $\alpha \in \mathbb{N}$ such that for all, $n, m \geq \alpha$, then

$$d(x_n, x_m) \leq 1$$

i.e., $x_n = x_m$.

- (ii) Convergent to a point $l \in X$ if for all $\epsilon \geq 0$, there is $\alpha \in \mathbb{N}$ such that for all $n \geq \alpha$ then $d(x_n, l) \leq \epsilon$, i.e., $x_n = l$.

Lemma 2.8.⁶ Let $\{x_n\}$ be a sequence of complete digital metric space (X, d, ρ) . If there exists $\alpha \in (0, 1)$ such that $d(x_{n+1}, x_n) \leq \alpha d(x_n, x_{n-1}) \forall n$, then $\{x_n\}$ converges to a point in X .

Proposition 2.9.⁶ A sequence $\{x_n\}$ of points of a digital metric space (X, d, ρ) converges to a limit $l \in X$ if there is $\alpha \in \mathbb{N}$ such that for all $n \geq \alpha$, then $x_n = l$.

Proposition 2.10.¹³ Let f and g be digitally compatible mappings of type (R) of a digital metric space (X, d, ρ) into itself. Suppose that $\lim_n f x_n = \lim_n g x_n = t$ for some $t \in X$. then

- (a) $\lim_n g f x_n = f t$ if f is digitally continuous at t .
- (b) $\lim_n f g x_n = g t$ if g is digitally continuous at t .
- (c) $f g t = g f t$ and $f t = g t$ if f and g both are digitally continuous at t .

3. MAIN RESULTS.

Theorem 3.1: Let P, Q, G and H be quadruple mappings of a complete digital metric space (X, d, ρ) satisfying the conditions.

- 1) $G(X) \subset Q(X), H(X) \subset P(X)$,

$$2) \quad d(x, y) = \alpha \max \left\{ d(Gx, Hy), d(Gx, Px), d(Hy, Qy), d(Hx, Gy), d(Hx, Qx), \frac{1}{2} (d(Gx, Qy) + d(Hy, Px)) \right\} \text{ for all } x, y \in X, \text{ where } \alpha \in (0, 1).$$

3) In mappings P, Q, G and H one of the mappings is continuous.

Assume the pairs (P, G) and (Q, H) are compatible, then P, Q, G and H have a unique common fixed point in X .

Proof: Let x_0 be an arbitrary point in X . Since $Q(X) \subset G(X)$ and $P(X) \subset H(X)$, we can construct the sequence $\{y_n\}$ in X such that,

$$y_{2n} = G(x_{2n}) = Q(x_{2n+1}) \text{ and } y_{2n+1} = H(x_{2n+1}) = P(x_{2n+2}) \text{ for each } n \geq 0.$$

$$d(y_{2n}, y_{2n+1}) = d(G(x_{2n}), H(x_{2n+1}))$$

$$\leq \alpha \max \left\{ d(P(x_{2n}), Q(x_{2n+1})), d(P(x_{2n}), G(x_{2n})), d(Q(x_{2n+1}), H(x_{2n+1})), d(G(x_{2n}), H(x_{2n+1})), \frac{1}{2} (d(G(x_{2n}), Q(x_{2n+1})) + d(P(x_{2n}), H(x_{2n+1}))) \right\}$$

$$\leq \alpha \max \left\{ d(y_{2n-1}, y_{2n}), d(y_{2n-1}, y_{2n}), d(y_{2n}, y_{2n+1}), d(y_{2n-1}, y_{2n+1}), \frac{1}{2} (d(y_{2n}, y_{2n}) + d(y_{2n-1}, y_{2n+1})) \right\}.$$

$$\leq \alpha \max \left\{ d(y_{2n-1}, y_{2n}), d(y_{2n}, y_{2n+1}), d(y_{2n}, y_{2n-1}), d(y_{2n-1}, y_{2n+1}), d(y_{2n+1}, y_{2n}), \frac{1}{2} (d(y_{2n-1}, y_{2n}) + d(y_{2n}, y_{2n+1})) \right\}$$

Putting $d_n = d(y_n, y_{n+1})$, we have

$$d_{2n} \leq \alpha \max \left\{ d_{2n-1}, d_{2n}, \frac{1}{2} (d_{2n-1} + d_{2n}) \right\}.$$

Now let $d_{2n} > d_{2n-1}$.

Therefore, $d_{2n} \leq 2\alpha d_{2n}$, for all $\alpha \in (0, 1)$, which is a contradiction.

Hence, $d_{2n} \leq d_{2n-1}$.

Let $m, n \in N$ such that $m > n$, we have

$$d(y_m, y_n) \leq \alpha d(y_m, y_{m-1}) + \dots + \alpha d(y_{n+1}, y_n) \\ \leq \alpha^n (d(y_1, y_0)) \rightarrow 0 \text{ as } m, n \rightarrow \infty.$$

It implies $\{y_n\}$ is digitally Cauchy sequence in digital metric space (X, d, ρ) and converges to point z as $n \rightarrow \infty$. Similarly sub sequences $\{Gx_{2n}\}, \{Px_{2n}\}, \{Hx_{2n+1}\}$ and $\{Qx_{2n+1}\}$ also converges to the point z .

Now suppose P is continuous. Since P and G are digitally compatible mapping then from Proposition 2.10 the sequence $\{PPx_n\}$ and $\{GPx_{2n}\}$ converges to Pz as $n \rightarrow \infty$.

Now we claim that $z = Pz$, for this on putting $x = Px_{2n}$ and $y = x_{2n+1}$ in condition 2 we have

$$d(GPx_{2n}, Hx_{2n+1}) \leq \alpha \max \left\{ d(PPx_{2n}, Qx_{2n+1}), d(PPx_{2n}, GPx_{2n}), d(Qx_{2n+1}, Hx_{2n+1}), d(PPx_{2n}, Gx_{2n+1}), \frac{1}{2}(d(GPx_{2n}, Qx_{2n+1}) + d(PPx_{2n}, Hx_{2n+1})) \right\}$$

Letting $n \rightarrow \infty$, we have

$$d(Pz, z) \leq \alpha \max \left\{ d(Pz, z), d(Pz, Pz), d(z, Pz), d(Pz, Pz), d(z, z), \frac{1}{2}(d(Pz, z) + d(Pz, z)) \right\} \leq \alpha d(Pz, z)$$

It implies that $Pz = z$.

Now we claim that $Gz = z$. Putting $x = z, y = x_{2n+1}$ in condition2 we have

$$d(Gz, Hx_{2n+1}) \leq \alpha \max \left\{ d(Pz, Qx_{2n+1}), d(Px, Gx_{2n+1}), d(Qx_{2n+1}, Hx_{2n+1}), d(Px, Hx_{2n+1}), d(Gx_{2n+1}, Qx_{2n+1}), \frac{1}{2}(d(Gz, Qx_{2n+1}) + d(Pz, Hx_{2n+1})) \right\}$$

Letting $n \rightarrow \infty$, we have

$$d(Qz, z) \leq \alpha \max \left\{ d(z, z), d(z, Gz), d(z, Hz), d(z, Qz), d(z, z), \frac{1}{2}(d(Gz, z) + d(Gz, z)) \right\} \leq \alpha d(Gz, z)$$

It implies that $Pz = z$.

Since $G(X) \subset Q(X)$ and hence there exists a point u in X such that $z = Gz = Qu$.

Now we claim that $z = Hu$.

$$\begin{aligned} d(z, Hu) &= d(Gz, Hu) \\ &\leq \alpha \max \left\{ d(Pz, Qu), d(Pz, Gz), d(Qu, Hu), d(Pz, Hu), d(Pu, Qu), \frac{1}{2}(d(Gz, Qu) + d(Pz, Hu)) \right\} \\ &\leq \alpha \max \left\{ d(z, z), d(z, z), d(z, Hu), d(z, Qu), d(z, Gu), \frac{1}{2}(d(z, z) + d(z, Hu)) \right\}. \end{aligned}$$

It implies that $z = Hu$.

Since (Q, H) is compatible and $Qu = Hu = z$, by proposition 2.7 we have $d(QHu, HQu) = 0$ and hence $Qz = QHu = HQu = Hz$. Also from condition 2, we have

$$\begin{aligned} d(z, Qz) &= d(Gz, Hz) \\ &\leq \alpha \max \left\{ d(Pz, Qz), d(Pz, Gz), d(Pz, Hz), d(Gz, Hz), d(Qz, Hz), \frac{1}{2}(d(Gz, Qz) + d(Pz, Hz)) \right\} \\ &\leq \alpha \max \left\{ d(z, Qz), d(z, z), d(z, Hz), d(z, Gz), d(Qz, Qz), \frac{1}{2}(d(z, Qz) + d(z, Qz)) \right\}. \end{aligned}$$

It implies that $z = Qz$.

Hence $z = Qz = Hz = Pz = Gz$.

Therefore P, Q, G and H have common fixed point z .

Similarly, proof can also be completed taking Q as continuous.

Now suppose G is continuous. Since P and G are compatible by Proposition 2.10 we have GGx_{2n} and PGx_{2n} converges to Gz as $\rightarrow \infty$.

We claim that $z = Gz$ and on putting $x = Gx_{2n}, y = x_{2n+1}$ in condition 2 we have,

$$d(GGx_{2n}, Hx_{2n+1}) \leq \alpha \max \left\{ d(PGx_{2n}, Qx_{2n+1}), d(PGx_{2n}, GGx_{2n}), d(PGx_{2n}, Hx_{2n+1}), d(PGx_{2n}, HHx_{2n}), d(Qx_{2n+1}, Hx_{2n+1}), \frac{1}{2}(d(GGx_{2n}, Qx_{2n+1}) + d(PGx_{2n}, Hx_{2n+1})) \right\}$$

Letting $n \rightarrow \infty$, we have

$$d(Gz, z) \leq \alpha \max \left\{ d(Gz, z), d(Gz, Gz), d(Gz, Hz), d(Hz, Gz), d(z, z), \frac{1}{2}(d(Gz, z) + d(Gz, z)) \right\} = \alpha d(Gz, z),$$

It implies $Gz = z$.

Since $GX \subset QX$, hence there exists a point w in X such that $z = Gz = Qw$.

We claim that $z = Hw$ and on putting $x = Gx_{2n}, y = w$ in condition 2 we have

$$d(GGx_{2n}, Hw) \leq \alpha \max \left\{ d(PGx_{2n}, Qw), d(PGx_{2n}, GGx_{2n}), d(PGx_{2n}, HHx_{2n}), d(PGx_{2n}, HGx_{2n}), d(Qw, Hw), \frac{1}{2}(d(GGx_{2n}, Qw) + d(PGx_{2n}, Hw)) \right\}$$

i.e.

$$d(z, Hw) \leq \alpha \max \left\{ d(z, z), d(z, z), d(z, z), d(z, z), d(z, Hw), \frac{1}{2}(d(z, z) + d(z, Hw)) \right\}$$

it implies that $z = Hw$.

Since Q and H are compatible on X and $Qw = Hw = z$, so by Proposition 2.7 $d(QHw, HQw) = 0$.

Hence $Qz = QHw = HQw = Hz$.

Now we claim that $z = Hz$, on putting $x = x_{2n}, y = z$ in condition 2 we have

$$d(Gx_{2n}, Hz) \leq \alpha \max \left\{ d(Px_{2n}, Qz), d(Px_{2n}, Gx_{2n}), d(Px_{2n}, Hx_{2n}), d(Px_{2n}, Hz), d(Qz, Hz), \frac{1}{2}(d(Gx_{2n}, Qz) + d(Px_{2n}, Hz)) \right\}$$

i.e.

$$d(z, Hz) \leq \alpha \max \left\{ d(z, Hz), d(z, z), d(z, z), d(z, z), d(Hz, Hz), \frac{1}{2}(d(z, Hz) + d(z, Hz)) \right\}$$

it implies that $Hz = z$.

Since $HX \subset PX$, there exists a point p in X such that $z = Hz = Pp$.

We claim that $z = Gp$, on putting $x = p, y = z$ in condition 2 we have

$$\begin{aligned} (Gp, z) = d(Gp, Hz) &\leq \alpha \max \left\{ d(Pp, Qz), d(Pp, Gp), d(Pp, Qz), d(Pp, Hz), d(Qz, Hz), \right. \\ &\quad \left. \frac{1}{2}(d(Gp, Qz) + d(Pp, Hz)) \right\} \\ &\leq \alpha \max \left\{ d(z, z), d(z, Gp), d(z, Gz), d(z, z), d(Hz, Hz), \frac{1}{2}(d(Gp, z) + d(z, z)) \right\}, \end{aligned}$$

It implies that $Gp = z$.

Since, P and G are compatible on X , $Gp = Pp = z$. So, by proposition 2.7, we have $d(PGp, GPP) = 0$ and hence $Pz = PGp = GPP = Gz$.

That is, $z = Pz = Gz = Qz = Hz$. Therefore, z is common fixed point of P, G, Q and H .

Similarly, proof can be completed when H is continuous.

Hence, uniqueness follows easily.

Example 3.2. Let $X = [0, \infty)$ be a digital metric space, and $M(x, y, \rho) = \left(\frac{\rho}{\rho+1}\right) d(x, y)$, where $d(x, y) = |x - y|$. Let $P: X \rightarrow X$ be a self-mapping defined by

$$P(x) = \begin{cases} \frac{x}{6} & \text{if } x \leq 1 \\ 0 & \text{if } x \geq 1 \end{cases}$$

We can easily that P is a continuous mapping.

REFERENCES

1. Banach S. *Sur les operations dans les ensembles abstraits et leurs applications aux equations integrales*, Fund. Math. 1922; 3: 133–181.
2. Rosenfeld A. *Digital Topology*. The American Mathematical, Monthly 1979; 86: 621-630.
3. Boxer L. *Properties of Digital Homotopy*. Journal of Mathematical Imaging and vision 2005; 22: 19-26.
4. Boxer L. *Continuous Maps on Digital Simple Closed Curves*. Applied Mathematics 2010; 1: 377-386.
5. Ege O. and Karaca I. *Lefschetz Fixed Point Theorem for Digital Images*. Fixed Point Theory and Applications, 2013;13.
6. Ege O. and Karaca I. *Banach fixed point theorem for digital images*, J. Nonlinear Sci. Appl. 2015; 8: 237-245.
7. Rosenfeld A. *Continuous functions on digital pictures*, Pattern Recognition Letters 1986; 4: 177- 184.
8. Jungck G. *Commuting mappings and fixed point*, Amer. Math. Monthly 1976; 83: 261- 263.
9. Jungck G. *Compatible mappings and common fixed points*, ibid, 11(2): 285-288.

10. Jungck G. Murthy PP. Cho YJ. *Compatible mappings of type (A) and common fixed points*, Math. Japon., 1993; 38: 381 – 390.
11. Dalal S. Masmali I.A. and Alhamzi G.Y. *Common Fixed Point Results for Compatible Map in Digital Metric Space*. Advances in Pure Mathematics 2018; 8: 362-371.
12. Fisher B. and Sessa S. *Common fixed points of two pairs of weakly commuting mappings*, Univ. of Novisad, Math. Ser. 1986; 16: 45-59.
13. Jain D. *Compatible Mappings of Type (R) in Digital Metric Spaces*. International Journal of Advanced Research Trends in Engineering and Technology (IJARTET). 2018; Vol. 5 No. 4: 67-71.
14. Alam A. and Imdad M. *Comparable Linear Contractions in Ordered Metric Spaces* 2015. Available from : <https://arxiv.org/abs/1507.08987>.