

International Journal of Scientific Research and Reviews

Moss Flora of Dachigam National Park (J&K) India

Amarpreet Kour

Department of Botany, Panjab University, Chandigarh-160014

ABSTRACT

Dachigam National Park is 122kms away from the city Srinagar. It is located at an altitude of 5500ft to 14000ft above the sea level and covers an area of 141sqkms. Present study was conducted to record moss flora of the area as there is no previous work done. In present investigation, a total number of 41 species have been recorded from study area. These species fall into 14 families and 21 genera.

KEYWORDS – Dachigam, Cervuselaphus, mosses

Corresponding Author:

Amarpreet Kour

Department of Botany,
Panjab University, Chandigarh-160014

E mail: amarpreetkour@gmail.com

INTRODUCTION

Dachigam National Park is famous for the Kashmir Stag or “Hangul” (*Cervuselaphushanglu*). Dachigam is abode to many animals like Himalayan Brown Bear, Himalayan Black Bear, Musk Deer, Leopard and Hyena. Park constitutes 150 species of birds and 661 species of vascular plants¹. Qureshi *et al.* have reported 35 species of butterflies from the park. It is strictly reserved for the protection of the wildlife and biodiversity. The boundaries are well marked and constrained².

Mosses constitute an important component of the park. The area was previously unexplored for moss flora and it was extensively researched for mosses. Kashyap has documented twenty two hepatic taxa from Kashmir valley and nine from Ladakh region^{3,4}. Kaul and Dhar recorded thirty five species of bryophytes from the Kashmir valley⁵. Kaul and Singh reported 37 mosses from the region⁶. Townsend also made collection of mosses from the valley⁷. Banday *et al.* explored the bryoflora of Kashmir valley⁸. Tanwir and Langer investigated Marchantialean flora of Jammu region. In 2006 same researchers explored liverworts of Ladakh, (J&K)^{9,10}. Tanwir *et al.* explored liverworts and Hornwort of Patnitop and its adjoining areas (J&K)¹¹. Dar and Khuroo, reported a 210 species of bryophytes (95 genera and 41 families) from Kashmir¹². Rashid *et al.* studied bryophytes of district Rajouri (Jammu) and reported of 44 bryophyte species included referable representing 16 families and four orders¹³. Banday *et al.* (2004) studied restricted habitat preference in Kashmir Himalayan mosses. In their study, 16 species of mosses show restricted habitat preference, while others show a wide range in their habitat colonization¹⁴. Kour *et al.* recorded eight species of mosses, namely *Brachythecium kamounense* (Harv.) A. Jaeg., *Brachythecium plumosum* (Hedw.) Schimp., *Brachythecium rutabulum* (Hedw.) Schimp., *Cratoneuron commutatum* (Hedw.) G. Roth., *Drepanocladus aduncus* (Hedw.) Warnst., *Haplocladium microphyllum* (Hedw.) Broth., *Hypnum cupressiforme* Hedw. and *Philonotis falcata* (Hook.) Mitt. for the first time from Doodhpathri (Budgam) J&K¹⁵. In another study Kour *et al.* recorded five species of genus *Mnium* namely; *Mnium confertidens*, *Mnium cuspidatum*, *Mnium integrum*, *Mnium rostratum* and *Mnium succulentum* for the first time from district Budgam¹⁶. Sharma *et al.* documented a total of 21

mosses belonging to 7 orders, 11 families and 16 genera. are reported with the family Pottiaceae being the most speciose¹⁷.

STUDY AREA

Dachigam National Park is located between 34° 04' -34° 14' N latitudes and 74°48' to 75° 85' E longitudes. It is 122kms away from the city Srinagar and covers an area of 141sqkms¹⁸. Park is divided into two parts upper and lower Dachigam. In the north-east, it is bounded by Sindh valley, Tarsar, Lidderwath, Kolhai of Lidder Valley and in the East, Overa-Aru Wildlife Sanctuary. In south- East, it is surrounded by Tral range and in west and south-west, Harwan, Brain and Nishat (Figure A, B & C)¹⁹.

Figure 1: Map of Dachigam.

RESULTS AND DISCUSSION

In present investigation, a total number of 42 bryophyta species have been recorded from the area.

Barbula constricta Mitt. Musci Ind. Or.: 33(1859).

Family: Pottiaceae

Habitat: Moist Soil

Barbulacylindrica (Tayl.) Schimp. var. *vinealis* (Brid) Lindb. Musci. Scand.: 32(1879)

Family: Pottiaceae

Habitat: Moist Soil

Barbula nigrescens Mitt. Musci Ind. Or. 36(1859)

Family: Pottiaceae

Habitat: Moist Soil

Brachythecium bucharanii (Hook.) Jaeg. Ber. S. Gall. Naturw. Gess., 1876-77: 341(1878)

Family: Brachytheciaceae

Habitat: Moist Soil

Brachythecium kamounense (Harv.) Jaeg. Ber. S. Naturw. Ges., 1876-77: 342(1878)

Family: Brachytheciaceae

Habitat: Moist Soil

Brachythecium rivulare B.S. G. Bryol. Eur., 6: 17(1853)

Family: Brachytheciaceae

Habitat: Moist Soil

Brachytheciumrutabulum (Hedw.) B.S.G. Bryol. Eur., 6: 15(1853)

Family:Brachytheciaceae

Habitat: Moist Soil

*Bryumalpinum*Huds. exWith. Syst. Arr. Birt. Pl. ed. 4, 3: 824(1801)

Family:Bryaceae

Habitat: Moist Soil

*Bryumargenteum*Hedw. Musc. : 181 (1801)

Family:Bryaceae

Habitat: Moist Soil

Distichiuminclinatum (Hedw.) B.S.G., Bryol. Eur., 2: 193(1864)

Family:Ditrichaceae

Habitat: Moist Soil

Ditrichumflexicaule (Schwaegr.) Hamp, Flora, 50: 182(1867)

Family:Ditrichaceae

Habitat: Moist Soil

Drepanocladusaduncus (Hedw.) Warnst. Beih.Bot.Centralbl. 13:400 (1903)

Family:Amblystegiaceae

Habitat: Moist Soil

Drepanocladusexannulatus (B. S. G.) Warnst. Beih. Bot. Centralbl., 13:405(1903)

Family:Amblystegiaceae

Habitat: Moist Soil

Entadoncurvatus (Griff) Jaeg. Ber. S. Gall. Naturw. Ges. 1876-77: 293 (1878)

Family:Entodontaceae

Habitat: Wooden Logs

*Entadonnepalensis*Mizushima Hara: Fl. E. Him.: 584, 42 (1966)

Family:Entodontaceae

Habitat: Wooden Logs

Entadonovicarpus Dix J. Bomb. Nat. Hist. Soc., 39: 789 (1937)

Family:Entodontaceae

Habitat: Wooden Logs

Eurhynchiumriparioides (Hedw.) Richs. Ann. Bryol., 9:135(1937)

Family:Brachytheciaceae

Habitat: Moist Soil

Eurhynchiumswartzii(Turn.) Curnow Rabenh.:Bryoth. Eur., 12: 593(1862)

Family:Brachytheciaceae

Habitat: Moist Soil

Funariahygrometrica(Hedw.) Sp. Musc. : 172(1801)

Family:Funariaceae

Habitat: Moist Soil

Grimmaovalis (Hedw.) Lindb. Act. Soc. Sci. Fenn., 10: 75(1871)

Family:Grimmiaceae

Habitat: Rocks

Grimmiaelongata Kaulfuss in J. Sturm *et al.*,Deutschl. Fl. 15: (1816)

Family:Grimmiaceae

Habitat: Rocks

Grimmiafunalis (Schwaegr.) B.S.G Bryol. Eur., 3: 119 (1845)

Family:Grimmiaceae

Habitat: Rocks

Grimmialaevigata (Bridel) Bridel, Bryol. Univ. 1: 183 (1826)

Family:Grimmiaceae

Habitat: Rocks

Hyophilainvoluta (Hook.) Jaeg. Natur. 1871-72: 356 (1873)

Family:Pottiaceae

Habitat: Rocks

Hyophilaspathulata (harv) Jaeg.,Ber. S. Gall. Naturw. Ges., 1817-72: 353(1873)

Family:Pottiaceae

Habitat: Rocks

Hyophilawalkeri Broth. Rec. Bot. Surv. Ind., 1:317 (1899)

Family:Pottiaceae

Habitat: Rocks

*Hypnumcupressiforme*Hedw. Sp. Musc. : 291(1801)

Family:Hypanaceae

Habitat: Sandy soil

Leucodonsciuroide (Hedw.) Schwaegr., Suppl. 1: (1816)

Family:Leucodontaceae

Habitat: Tree

Lindbergiakoelzii Williams, Bryologist, 35 : 3(1932)

Family:Leskeaceae

Habitat: Moist Soil

*Mniumcuspidatum*Hedw. Musc. :192(1801)

Family:Mniaceae

Habitat: Moist Soil

*Mniumrostratum*schrad. Regensburg, 1:79(1802)

Family:Mniaceae

Habitat: Moist Soil

Orthotheciumintriatum(Hartm.) B. S. G., Bryol. Eur.,

Family:Hypnaceae

Habitat: Moist Soil

*Polytrichumalpinum*Hedw. Sp. Musc. : 92 (1801)

Family:Polytrichaceae

Habitat: Moist Soil

*Racomitriumcrispulum*Hook.f. & Wilson,J.D.Hooker, Fl. Nov.-Zel. 2: 76 (1854)

Family:Grimmiaceae

Habitat: Rocks

Racomitriumlanuginosum (Hedwig) Bridel, Muscol. Recent., suppl. 4: (1818)

Family:Grimmiaceae

Habitat: Rocks

RhynchostegiellaScabriseta(Schwaegr.) Broth. Nat. Pfl.1(3): 1161 (1909).

Family:Brachytheciaceae

Habitat: Moist soil

Rhynchostegiumplaniusculum (Mitt.) A. Jaeg. Ber. S. Gall. Naturw. Ges., 1876-77: 364(1878).

Family:Brachytheciaceae

Habitat: Moist soil

Syntrichiaprinceps (De Not.) Mitt. Musc. Ind.Or. :39(1859).

Family:Pottiaceae

Habitat: Rocks

Thuidiumglaucinum (Mitt.) Bosch &Sande Lac.Bryol.Jav., 2: 117 (1876).

Family:Thuidiaceae

Habitat: Moist soil

Thuidium orientale Mitten ex Dixon J. Bot., 51: 329 (1913).

Family: Thuidiaceae

Tortellatortusoa (Hedw.) Limpr. Laubm. Deutschl., 1 : 604 (1888).

Family: Pottiaceae

Habitat: Moist soil

A total number of 41 species of bryophyte have been recorded from Dachigam National Park. These species fall into 21 genera and 14 families. Brachytheciaceae and Pottiaceae families are most speciose, representing highest number of species (8), followed by Grimmiaceae (6) and Entodontaceae (3). Amblystegiaceae, Bryaceae, Ditrichaceae, Hypanaceae, Mniaceae and Thuidiaceae possess two species. Families like Funariaceae, Leskeaceae, Leucodontaceae and Polytrichaceae possess one species. The field notes are also added to depict the habitat preferences of each taxon. These bryophytic species were mostly present on moist soil, trees, rocks and sandy soil.

CONCLUSIONS

Globally 2 to 4% of bryoflora has been threatened in the short term which is a matter of great concern²⁰. In view of the threatening situation the study of the bryophytic flora of this unexplored area assumes greater importance. Global warming is going to be a big threat for bryophytes. According to IPCC, there is 0.6 degree increase to average temperature on earth in last century and will raise to 1.4 to 5.8°C by 2100²¹. Habitat destruction, deforestation, scarcity of water and increase in growing population are other factors that are threat to bryophytes. These plants need attention for their conservation and protection. Some strict policies are required for their protection. In view of the threatening situation the study of the bryophytic flora of this unexplored area assumes greater importance.

ACKNOWLEDGEMENT

I would like to thank Chief Wild life Warden, Mr. Chaturbhuja Behera for granting me permission to collect bryophytes from the highly restricted area. I would also like to thank Nazir Malik (Jungle lover), who helped me during my collection. Last but not least, I would like to thank my loving dad, S. Tejasingh for his infinite love and inspiration.




Figure A: Map of Dachigam

Figure B & C: Field photographs of the study area.

REFERENCES

1. Dar GA, Bhagat RC&Khan MA. Biodiversity of the Kashmir Himalaya. Valley Book House, Srinagar, India ;2002.
2. Qureshi A A, Bhagat RC &Bhat DM. Diversity of butterflies (Lepidoptera: Papilionoidea and Hesperoidea) of Dachigam National Park, Jammu and Kashmir, India. *Journal of Threatened Taxa*, 2014; 6(1): 5389-5392.
3. Kashyap SR. Liverworts of the Western Himalayas and the Panjab Plain I. The University of the Panjab, Lahore, 1929.
4. Kashyap SR. Liverworts of the Western Himalayas and the Panjab Plain II. The University of the Panjab, Lahore, 1932.
5. Kaul RK &Dhar GL. Some bryophytes of Kashmir valley. *Kashmir Science*, 1968; 5: 233 - 237.
6. Kaul RK & Singh G. Some mosses from Kashmir. *The Bryologist*, 1972; 75(4): 531-538.
7. Townsend CC. Mosses from Kashmir. *Journal of Bryology*, 1988; 15: 293-297.
8. Banday FA, Naqshi AR &Dar GH. Liverworts (Hepaticae) of Kashmir Himalaya, a floristic survey. *Oriental Science*, 1998; 3: 1-6.
9. Tanwir M &Langer A. Marchantialean flora of Jammu region (J & K State)-A preliminary survey. A. N. Kamili and A. R. Yousuf (Eds.) *In: Bioresources: Consersns and Conservation*, *CARD*, University of Kashmir, Srinagar, 2004; 23-30.
10. Tanwir M & Langer A. Liverworts of Ladakh, J&K State (North-West Himalaya), India. *Journal of the Indian Botanical Society*, 2006; 85: 71-73.
11. Tanwir M & Langer A, Bhandari M. Liverwort and hornwort of Patnitop and its adjoining areas (J&K), Western Himalaya, India," *Geophytology*, 2008; 37(1-2): 35-41.
12. Dar GH &Khuroo AA. Floristic Diversity in the Kashmir Himalaya: progress. *SainsMalaysiana*, 2013; 42(10): 1377-1386.
13. Rashid A, Mishra R & Sharma A. Bryoflora of district Rajouri - Jammu and Kashmir State, India. *Earth Foundation International. Archive for Bryology*, 2012; 145: 1-11.
14. Banday FA, Reshi Z, Kapila S & Kumar SS. Restricted habitat preference in Kashmir Himalayan mosses. *Punjab University Research Journal*, 2004; 54: 75-77.
15. Kour A, Rao A &Kapila S. Taxonomic study of some mosses of Doodhpathri (Budgam) J&K, India. *International Journal of Advances in Pharmacy, Biology and Chemistry*, 2015; 4(1): 196-208.

16. Kour A, Rao A, Kapila S & Kumar SS. Taxonomical studies of genus *Mnium* from district Budgam (J&K) India. *International Journal of Science and Research*, 2015; 4(3): 1651-1658.
17. Sharma A, Langer A & Uniyal PL. A Preliminary Report on the Mosses of Rajouri and Poonch Districts of Jammu & Kashmir, India. *theamericanbryological and lichenological society inc*, 2016; 33(1): 26-33.
18. Shameem SA & Kangroo IN. Comparative assessment of edaphic features and Phytodiversity in lower Dachigam National Park, Kashmir Himalaya, India. *African Journal of Environmental Science and Technology* Vol. 5(11), pp. 972-984.
19. Kurt, F. Kashmir Deer (*Cervuselaphushanglu*) in Dachigam.. In: Scott, Peter: Threatened deer. *Int. Union Conserv. Nat. Nat. Resour (IUCN)*, Morges, Swetzerland. 1978, 87-109
20. Vanderpoorten A & Hallingbäck T. Conservation biology of bryophytes. Ch 12 in: *Bryophyte Biology*. Second Edition, eds. Goffinet, B. & Shaw, A.J. Cambridge University Press, 2008; 487-532.
21. Kulshrestha UC. Global warming-Present status of research and future strategies. *J. Ind. Geophys. Union*, 2012; 16(4): 143-160.