

## *International Journal of Scientific Research and Reviews*

### **H-Recurrent Finsler Connection**

**Brijendra Krishna Singh**

Department of Mathematics Amity University Chhattisgarh, Raipur, India

Email : [bks0509@gmail.com](mailto:bks0509@gmail.com)

---

#### **ABSTRACT**

The Decomposition of the normal Finsler connection tensor  $N_{jkh}^i$  of a finsler connection in the form of H Recurrent Finsler Connection and assume that decompose vector field  $X^i$  is not independent of directional arguments then thenormal projective curvature tensor are connected by recurrent Finsler connection.

**KEYWORDS:** *Finsler, manifolds, torsion, projective, recurrence*

---

**\*Corresponding author:**

**Brijendra Krishna Singh**

Department of Mathematics

Amity University Chhattisgarh,

Raipur, India

Email : [bks0509@gmail.com](mailto:bks0509@gmail.com)

**INTRODUCTION:**

A Finsler manifold  $F_n$  of dimension  $n$  is a manifold  $F_n$  associated with a fundamental function  $F(x, \dot{x})$ , the metric tensor of  $(F_n, F)$  is given by

$$(1.1) \quad g_{ij} = \frac{1}{2} \dot{\partial}_i \dot{\partial}_j F^2 \text{ where } \dot{\partial}_i = \partial / \partial \dot{x}^i.$$

A Finsler connection of  $(F_n, F)$  is a triad  $(F_{jk}^i, N_k^i, C_{jk}^i)$  of a v-connection  $F_{jk}^i$ , a nonlinear connection  $N_k^i$  and a vertical connection  $C_{jk}^i$  [6]. The h- and v- covariant derivatives of any tensor field  $V_j^i$  corresponding to a given Finsler connection is given by

$$(1.2) \quad V_{j|k}^i = d_k V_j^i + V_j^m F_{mk}^i - V_m^i F_{jk}^m,$$

$$(1.3) \quad V_{j|k}^i = \partial_k V_j^i + V_j^m C_{mk}^i - V_m^i C_{jk}^m$$

where (1.4)  $d_k = \partial_k - N_k^m \partial_m$ ,  $\partial_k = \partial / \partial x^k$ .

From a given Finsler metric we can determine various Finsler connections. In the present studies we shall use the Cartan connection which will be denoted by  $C\Gamma: (\Gamma_{jk}^{-xi}, G_k^i, C_{jk}^i)$ . These connections can be uniquely determined from the metric function  $F$  by the following axioms:

- (A<sub>1</sub>) The connection is h – metrical i.e.  $g_{ij}/k = 0$ ,
- (A<sub>2</sub>) The connection is v – metrical i.e.  $g_{ij}/k = 0$ ,
- (A<sub>3</sub>) The deflection tensor field  $D_k^i$  vanishes,
- (A<sub>4</sub>) The (h) h – torsion tensor field  $T_{jk}^i$  vanishes,
- (A<sub>5</sub>) The (v) v – torsion tensor field  $S_{jk}^i$  vanishes.

All these five axioms have been mentioned in [7]. The individual members of the triad are given as

$$(1.13) \quad \Gamma_{jk}^{xi} = \frac{1}{2} g^{ih} (d_k g_{jh} + d_j g_{kh} - d_h g_{jk}),$$

$$(1.14) \quad a) \quad G_k^i = \partial_k G^i = \gamma_{ok}^i - 2C_{km}^i G^m,$$

$$b) \quad G^i = \frac{1}{2} \gamma_{oo}^i,$$

$$(1.15) \quad C_{j|k}^i = g^{ih} C_{jhk}, \quad C_{jkh} = \frac{1}{2} \partial_h g_{jk},$$

where (1.16)  $\gamma_{jk}^i = \frac{1}{2} g^{ih} (\partial_k g_{jh} + \partial_j g_{kh} - \partial_h g_{jk}),$

**DEFINITION (1.1):**

A Finsler connection will be called h-recurrent Finsler connection  $RF\Gamma$  if it satisfies the following axioms:

(A<sub>1</sub>)' The connection is h-recurrent with recurrence vector  $\alpha_k$  i.e.  $g_{ij|k} = \alpha_k g_{ij}$ .

(A<sub>2</sub>)' The connection is v-metrical i.e.  $g_{ij|k} = 0$ .

(A<sub>3</sub>)' The deflection tensor field is given by  $D_k^i$ .

(A<sub>4</sub>)' The (h) h-torsion tensor field  $T_{jk}^i$  vanishes.

(A<sub>5</sub>)' The (v) v-torsion tensor field  $S_{jk}^i$  vanishes.

In view of equations (1.18), (1.20) and (1.22) we find that the h-recurrent Finsler connection  $RF\Gamma$  are given by

$$(1.23) F_{jk}^i = \overset{c}{F}_{jk}^i - C_{km}^i X_j^m - C_{jm}^i X_k^m + C_{jkm} X^{mi},$$

$$(1.24) N_k^i = \overset{c}{N}_k^i + X_k^i,$$

$$(1.25) C_{jk}^i = \overset{c}{C}_{jk}^i = \frac{1}{4} g^{ih} \dot{\partial}_h \dot{\partial}_j \dot{\partial}_k F^2$$

Where (1.26)  $X_k^i = C_{km}^i B_o^m - B_k^i$ ,

$$(1.27) B_k^i = D_k^i + \frac{1}{2} (\alpha_o \delta_k^o + \alpha_k \dot{x}^i - \alpha^i y_k)$$

$$(1.28) X^{mi} = g^{ji} X_j^m$$

and  $\left( \overset{c}{F}_{jk}^i, \overset{c}{N}_k^i, \overset{c}{C}_{jk}^i \right)$  are the coefficients of Cartan connection  $C\Gamma$ . With the help of the equations (1.8),

(1.23) and (1.24) the (v) hv-torsion tensor  $RF\Gamma$  can be written as

$$(1.29) P_{jk}^i = \overset{c}{P}_{jk}^i + X_j^i |k + C_{jm}^i X_k^m + C_{jkm} (X^{im} - X^{mi})$$

where  $P_{jk}^i$  is the (v) hv-torsion tensor of Cartan connection  $C\Gamma$  and  $|$  means v-covariant differentiation with respect to  $C\Gamma$  or  $RF\Gamma$ . Again using the equations (1.7) and (1.24), we get the following alternative form of (v) hv-torsion tensor of  $RF\Gamma$ .

$$(1.30) R_{jk}^v = \overset{c}{R}_{jk}^v - \overset{c}{P}_{jm}^i X_k^m + \overset{c}{P}_{km}^i X_j^m + X_j^i + C_{|k}^i \\ - X_k^i C_{|j}^i - X_k^m X_j^i |m + X_j^m X_k^i |m - C_{jm}^i X_r^i X_k^m + C_{km}^r X_r^i X_j^m$$

**THE (v) hv-TORSION TENSOR OF THE FORM  $P_{jk}^i = -\dot{\delta}_k^i B_j^i$**

In this section we shall pay our attention to that h-recurrent Finsler connection  $RF\Gamma$  whose (v) hv-torsion tensor  $P_{jk}^i$  is being expressed by the following equation

$$(4.1) P_{jk}^i = -\dot{\delta}_k B_j^i,$$

where  $B_j^i$  is the tensor field of the Finsler connection (1.27). Using (4.11) in (1.29), we get

$$(4.2) \overset{c}{P}_{jk}^i = \dot{\delta}_k (C_{jr}^i B_0^r) + C_{mk}^i X_j^m + C_{jm}^i X_k^m - C_{jkm} X^{mi} = 0.$$

Using  $\dot{\delta}_k g_{ij} = 2C_{ijk}$  in (4.2), we get

$$(4.3) \overset{c}{P}_{ijk} + \dot{\delta}_k (C_{ijr} B_0^r) - 2C_{irk} C_{jm}^r B_0^m + C_{imk} X_k^m + C_{ijm} X_k^m - C_{jkm} X_i^m = 0.$$

Since  $C_{ijk}$  and  $\overset{c}{P}_{ijk}$  are symmetric in  $i$  and  $j$ , hence from (4.3), we get

$$(4.4) S_{ijmk} B_0^m C_{imk} X_j^m - C_{jmk} X_i^m = 0.$$

Multiplying (4.4) by  $x^i$ , we get

$$(4.5) C_{jmk} X_0^m = 0.$$

An obvious of (4.5) is the equation

$$(4.6) X_j^i = -B_j^i \text{ and } C_{ikm} B_j^m = C_{jkm} B_i^m.$$

In the light of these observations from (4.3), we get

$$(4.7) \overset{c}{P}_{ijk} = C_{jkm} B_i^m.$$

Substituting these results into the equations (1.30), (1.31) and (1.32), we get

$$(4.8) R_{jk}^i = \overset{c}{R}_{jk} B_j^i C_{|k} + B_k^i C_{|j} - B_k^m B_j^i \Big|_m + B_j^m B_k^i \Big|_m,$$

$$(4.9) P_{hjk}^i = \overset{c}{P}_{hjk} S_{hjk}^i B_j^r,$$

and  $(4.10) R_{hjk}^i = \overset{c}{R}_{hjk} + \overset{c}{P}_{hjm} B_k^m - \overset{c}{P}_{hkm} B_j^m + S_{hrs}^i B_j^r B_k^s.$

If we now assume that

$$(4.11) P_{ijk}^c = C_{jkm} B_i^m \text{ holds,}$$

then this assumption gives

$$(4.12) C_{ijr} B_k^r = C_{ikr} B_j^r, C_{ijr} B_0^r = 0 \text{ and } X_j^i = -B_j^i.$$

Using (4.12) in (1.27), we get

$$(4.13) P_{jk}^i = -\dot{\delta}_k B_j^i.$$

Therefore, we can state.

### **THEOREM (4.1):**

If  $F_n$  be supposed to be an n-dimensional Finsler space equipped with h-recurrent Finsler connection  $RF\Gamma$  and with the deflection tensor  $D_j^i$  and recurrence vector  $\alpha_k$ , if we further suppose that  $B_j^i = D_j^i + \frac{1}{2}(\alpha_0 \delta_j^i + \alpha_j \dot{x}^i - \alpha^i y_j)$  then the (v) hv-curvature tensor  $P_{jk}^0$  of  $RF\Gamma$  is given by

$P_{jk}^i = -\dot{\delta}_k^i D_j^i$  if and only if the (v) hv-torsion tensor  $\overset{c}{P}_{jk}^i$  of the connection  $C\Gamma$  is represented by  $\overset{c}{P}_{jk}^i = C_{jm}^i B_k^m$  and in such a case the (v) h-torsion tensor  $R_{hjk}^i$  of the hv-curvature tensor  $P_{hjk}^i$  and the h-curvature tensor  $R_{hjk}^i$  of recurrent Finsler connection  $RF\Gamma$  are respectively given by (4.8), (4.9) and (4.10).

### **REFERENCES**

1. Brickell, F.: A theorem on homogeneous functions, J. Lond. Math.Soc., 1967;42: 352-359.
2. Hashiguchi, M. : On determination of Finsler connections by deflection tensor fields, Rep. Fac Sci. Kagoshima Univ. (Math, Phy, Chem) 1969; 2: 29-39.
3. Hashiguchi, M.: On Wagner's generalized Berwald Space, J. Korean Math. Soc. 1975 ;12: 51-61,.
4. Ikeda, F.: Some remarks on deflection tensor, Tensor (N.S.) 1990; 49:1-6.
5. Matsumoto, M.: A Finsler connection with many torsions, Tensor (N.S.) 1966; 17: 217-226.
6. Matsumoto, M.: The theory of Finsler connections, Publ. of the study group of Geometry, 5, Deptt Maths., Okayama Univ. 1970; 120.
7. Matsumoto, M.: Foundations of Finsler geometry and special Finsler spaces, Kaiseisha Press, Saikawa, Otsa, Japan 1986; 520.
8. Prasad, B.N.: On recurrent Finsler connections with deflection and torsion, Publications Shukla, H.S. and Mathe maticae, Hungaria,. Singh, D.D. 1990; 37, 77-84.