

International Journal of Scientific Research and Reviews

Enhancing Acclimatization of Tissue Cultured Plants By Biotization- A Review

G.Indravathi^{1,2*} and P.SureshBabu³

¹Dept. of Biotechnology, KVR Govt. College for Women (A), Kurnool, Andhra Pradesh, India.

²Dept. of Biotechnology, Jawaharlal Nehru Technological University, Anantapur,
Andhra Pradesh, India.

³Department of Biology, Indian Institute of Science Education and Research, Tirupati, Andhra
Pradesh, India.

ABSTRACT

The plants raised from tissue culture experience high mortality following laboratory to land transfer. Apart from the abiotic causes, one major cause of high mortality of such aseptically raised plants is their sudden exposure particularly the root system to microbial communities, including soil pathogens. Efficient antagonistic plant growth-promoting microbes have been used for hardening to increase the survival and to augment overall plant growth. Biotization is the metabolic response of *in vitro* grown plant material to microbial inoculum. Mycorrhization of tissue cultured plants is believed to provide an advantage to the transplanted propagules in terms of nutrient availability, soil pH, aeration and protection from water stress. The tissue culture raised plants when treated with bacterial inoculants produce plant growth promoting substances and some secondary metabolites which enhance nutrient uptake and provide resistance against pathogens. Multimicrobial biotization is inoculation of more than one microbial species to micro plants. Plantlet survival rate was maximum in dual inoculation, this must be due to the positive interaction between two or more species and their ability to enhance stress tolerance by protecting them from subsequent 'transplantation shock'. Thus biological hardening envisages physical, chemical and environmental conditioning of the micro propagated plantlets. Different parameters were reported for control and biotized plantlets to know the effect of microbes on plant performance. This review is focused on the effect of biotization on enhancing acclimatization of tissue culture raised plants.

KEY WORDS: Biotization, acclimatization, micropropagation, biotic factors.

***Corresponding author:**

G. Indravathi

Dept. of Biotechnology,

KVR Govt. College for Women(A),

Kurnool - 518002

Andhra Pradesh, India.

E-mail: gindravathi@gmail.com, Mobile : 9989067747

INTRODUCTION:

Tissue culture raised plants experience high mortality following laboratory to land transfer. One major cause of high mortality of aseptically raised plants apart from the abiotic factors is their sudden exposure, in particular, the root system to microbial communities, including minor and major pathogens, present in the soil. The primary target of several research groups attempting utilization of microbial inoculants in micropropagation is to induce stress resistance¹.

Biotization is the metabolic response of *in vitro* grown plant material to microbial inoculum, leading to the morphological and physiological development enhancing biotic and abiotic stress resistance of the derived propagules. Biotization is an emerging dimension of micropropagation technique where the growth of the host plant is promoted by the formation of secondary metabolites related to plant defense. Such systems allow for mutual adaptation between the host plant and the introduced bacteria. Bacterized plantlets not only grow faster than nonbacterized plantlets but they are sturdier and have a better-developed root system².

The inoculation of seeds with beneficial microorganisms has been practiced for many years, but the inoculation of tissue culture raised plantlets is an innovative aspect. Plant tissue culture is based on aseptic conditions, hence microorganisms including beneficial endophytes are treated as the problem causing contaminants. But nowadays microbial inoculants, primarily bacteria, and mycorrhizae are being evaluated as biopriming agents for successful transplanting. Biotization could be achieved during *in vitro* rooting or under *ex vitro* conditions. Thus efficient antagonistic plant growth-promoting bacteria used for biological hardening envisages physical, chemical and environmental conditioning of the micropropagated plantlets.

All beneficial microorganisms helped in the enhanced uptake of nutrients, nitrogen fixation, resistance to soil-borne diseases and improved plant water relations. Although the soils are not deficient in phosphorous, its unavailability in phosphorous fixation is found to be a major constraint in acidic soils. Effective microbes aids in the uptake of mineral nutrients in the available form where the normal plant roots fail. The microbial association helps in the improved hydraulic conductivity of the roots which contributes to the better uptake of water thereby resulting in the reduction of desiccation and wilting in immature plantlets. These favorable conditions help to increase the number of lateral roots and root length. Also, there was an increase in the activity of all the defense-related enzymes like phenylalanine ammonia lyase, peroxidase, and β -1,3-glucanase in the leaves of the plants under treatment³. Lower incidence of rotting and wilting diseases was

noticed in bioinoculant-treated plantlets. This suggests that bioinoculants were capable of eliciting systemic resistance.

Apart from bacteria and fungi, other endophytes like algae, amoebae, virus, archae, oomycetes also show a symbiotic association with plants⁴. For e.g., Green algae *Coccomyxa species* in *Ginkgo biloba*⁵ and amoebal cysts in *Eleutherococcus sieboldianus*⁶.




Fig.1: Effect of Abiotic and Biotic Factors on Acclimatization of Tissue Culture Raised Plants.

MYCORRHIZATION:

Acclimatization phase is a key step in the micropropagation cycle as it affects the survival and growth of the in vitro produced plantlets. There are many examples showing that the inoculation with Arbuscular Mycorrhizal Fungi (AMF), at the time of transplant from axenic to in vivo conditions, significantly improves plant survival and growth. Mycorrhization of tissue cultured plants is believed to provide an advantage to the transplanted propagules in terms of nutrient availability, soil pH, aeration and protection from water stress. An excellent review of factors affecting the result of mycorrhizal inoculation (timing, medium, fertilization, inoculation, fungus-host specificity, growth substrate, etc.) was published⁷.

Mycorrhization has been proved beneficial for many other microplants of raspberry, wild cherry, apple, pear, grapevine, rose, oil palm, citrus, banana, asparagus, pineapple, common ash,

artichoke^{8,9}. By enhancing access to the growth-limiting nutrients both ectomycorrhizae and endomycorrhizae can significantly increase carbon fixation of the plantlets¹⁰. This gain occurs primarily via increased photosynthetic rates. Mycorrhizal plants can also take up more carbon in drought periods than non-mycorrhizal ones, since they can maintain stomata open at lower soil water potentials.

Plants inoculated with ectomycorrhiza showed increased phosphorus and nitrogen uptake¹¹. In ectomycorrhizas, photosynthates move from the autotroph to the fungal mantle, where they are rapidly converted to metabolic intermediates, of which trehalose and mannitol are dominant. These carbon compounds are used for sustaining the fungal biomass existing in the mycorrhizal root tips of the soil mycelia network and also for producing new fungal biomass. As the photosynthetic rate is primarily limited by the accumulation of its end products in the leaf cells, the drain of carbohydrates by the mycobiont may offset end product limitation facilitating an increased photosynthetic rate. Due to this RuBisCO activity, chlorophyll and protein content of leaves of mycorrhizal plants, expressed in terms of fresh weight or of dry weight was reported significantly higher than in control plants. The increased photosynthetic rates were also due to increased N, P and K absorption⁵⁷. Relationships between foliar phosphorus and net photosynthetic rate in non-mycorrhizal and ectomycorrhizal were compared in Pine seedlings¹².

Among endomycorrhiza, Vesicular Arbuscular Mycorrhiza (VAM) plays a significant role in phosphorus uptake and translocation in addition to uptake of zinc, sulfur, and copper. VAMs are exceptional to establish a symbiotic relationship in the roots of higher plants. A careful selection of functionally compatible host fungus substrate combination was essential for the early establishment of VAM in the nursery or in an open field of major horticultural crops.

The mycorrhizal dependency of banana has been extensively studied by various researchers, emphasizing its importance during hardening stages¹³. A marked increase in the uptake of P, Ca, Mg, Zn, Cu and reduced disease severity was observed in the mycorrhizal plants of banana¹⁴. It was reported that VAM fungi – *Glomus fasciculatum* inoculated at the hardening stage helped the banana plantlets to accumulate maximum plant height, root length, biomass, root colonization and nutritional quality¹⁵. Similar results on growth characters and root colonization of papaya plants with *G. message* and *G. fasciculatum* were reported. Inoculation of micropropagated sugarcane seedlings with *G. diazotrophicus* made the plants not only grow faster but also ensured efficient N₂-fixing plants in fields¹⁶. It was also studied that the effect of inoculation of the fungus, *Glomus intraradices* increased the survival, growth, biomass production and nutritive status of cassava, grape and olive plants during hardening.

AMF can contribute to plant growth and survival by reducing the stress associated with

nutrition, water, aeration, soil structure, pH, salt, toxic metals, and pathogens. Biotization of AMF symbiotic endophytes also protect the juvenile axenic plants from an infestation of the harmful saprophytes. It was reported that AM fungi established a symbiotic association with hazelnut (*Corylus avellana* L.)¹⁷. Micropropagated plants of *Ranunculus asiaticus* inoculated by AMF showed an earlier flowering, an improved flower production and better rhizome yield¹⁸. The colonization of *Glomus*, *P.indica* and *Trichoderma* species is also known to reduce the osmotic potential of plants and also induced in vivo rooting in transplanted micro shoots of *C. borivilianum* giving rise to more than 50% establishment¹⁹. Mycorrhization enhanced the growth of micropropagated Chestnut plants, increased their protein content and photosynthetic rates, decreased respiratory rates and CO₂ compensation point.

Piriformospora indica, a root endosymbiotic fungus, mimics AMF in many morphological, functional aspects and growth promotion. It was reported that *P. indica* treated plantlets of *B. serrata* showed an increase in total biomass production and more than 75 percent ex vitro survival in comparison to control plantlets²¹. In particular, *P.indica* helps in phosphorus acquisition and works as a biocontrol agent. A similar high degree of ex vitro survival of micropropagated plantlets of *Artemisia annua*, *Nicotiana tobaccum*, *B. monnieri*, *T. bellerica*, and *F. limonia* colonized with *P. indica* was reported²². A positive influence of root colonization was reported with *P. indica* on overall growth and development in micropropagated plantlets of *T. bellerica*²³. *P.indica* were more resistant to pathogens and more tolerant to salt stress and showed higher yield²⁴.

The profound effect of *T. viride* on root initiation from in vitro micropropagated Neem shoots had been reported²⁵. It was reported that *T. viride* when applied alone to Broccoli showed maximum N and P content in roots and shoots²⁶. The enhanced vegetative growth of broccoli in *Trichoderma* treated plants could be due to the root-colonizing ability of the fungus that resulted in better nutrient absorption through increased root biomass.

BACTERIZATION:

The tissue culture raised plants when treated with bacterial inoculants produce plant growth promoting substances and few secondary metabolites that enhance nutrient supply and provide resistance against pathogens. These useful soil bacteria are named as plant growth promoting rhizobacteria (PGPR), which are preferentially related to the roots. PGPR include many bacterial genera like *Pseudomonas*, *Bacillus*, *Azospirillum*, *Azotobacter*, etc. Bacterial infection occurs through roots and they are translocated to all parts of the plant via the xylem, the aerenchyma and through interconnecting intercellular spaces²⁷. These diazotrophs would have affected plant growth by the synthesis of phytohormones and vitamins, inhibition of plant ethylene synthesis and improved

nutrient uptake²⁸. Bacterized plantlets were greener, had elevated levels of cytokinins, phenylalanine ammonia lyase, and free phenolics and contained more lignin²⁹. Both in vitro and ex vitro benefits of bacterization relied on plant species, cultivar, and growth conditions.

Among the PGPR, *Pseudomonas* deserves a special mention because it improves the plant growth directly or indirectly by the production of plant growth substances, increasing the uptake of certain nutrients from the soil and additionally shows antagonistic effects against some important plant pathogenic microorganisms. In addition to this *Pseudomonas* species have been used to enhance tolerance to transplanting stress in potato⁶². In vitro co-cultivation of soybean cotyledon explants with two strains of *Pseudomonas maltophilia*, stimulated the development of nodular callus with high regeneration potential³⁰. It was reported that *Pseudomonas* strain PsJN enhanced the tolerance to transplanting stress in potato and was found the most effective plant growth promoting bacterium under in vitro conditions³¹. The Oregano plantlets co-cultured with *Pseudomonas spp.* prevented vitrification and contained a lot of phenolics and chlorophyll than non bacterized controls³². Greenhouse experiments also demonstrated that plants derived from dual cultures of potato and the pseudomonad bacterium had a larger root system, and gave better tuber yield than control³³. The tea plants inoculated with *Bacillus subtilis* and *Pseudomonas corrugate* acted as biocontrol agents and were able to defend pathogenic attack probably due to their antagonistic properties³⁴.

Three plant growth-promoting rhizobacteria viz. *Bacillus megaterium*, *B. subtilis* and *Pseudomonas corrugata* were used for biological hardening of micro-propagated plantlets of *Picrorhiza kurroa*³⁵. These bacterial isolates antagonized the pathogenic fungal species and positively influenced survival and growth parameters. The in vitro grown Oil palm plantlets inoculated with *Azospirillum brasilense* produced higher root and shoot biomass and more secondary roots³⁶. Endophytic colonization in rice callus induced by using *Azorhizobium caulinodans* improved yield, grain weight and nutritional quality³⁷.

Thus the reports suggest the use of efficient antagonistic plant growth-promoting bacteria for biological hardening increase plant survival and augment overall plant growth of micro propagated plants.

MULTIMICROBIAL INTERACTION:

Multimicrobial biotization is inoculation of more than one microbial species to microplants. Root colonization by mycorrhizal fungi will have an affect the chemical composition of root exudates. The development of mycelium around roots (mycorrhizosphere) modifies their physical atmosphere and provides a new source of carbon to the microbial community. The development of a

mycorrhizosphere has both qualitative and quantitative repercussions on microbial populations in either the rhizosphere or rhizoplane³⁸. Mycorrhization increases the dichotomy of roots and creates new compartments (mycorrhizosphere and microsphere), for the growth of symbiotic microbes.

Mycorrhizae-Helping Bacteria (MHB) enhance the growth of the plants. It was reported that rhizosphere strains of *Bacillus mycoides* and *Pseudomonas fluorescens* promoted AMF formation in various crop plants by improving susceptibility of roots to AMF³⁹. Other rhizosphere microorganisms which are known to act as a phytostimulators or which possess antagonistic activities toward plant pathogens may be used in conjunction with AMF for biotizing microplants. They include bacteria like *Pseudomonas spp.* and *Bacillus spp.*, and fungi such as *Gliocladium spp.* and *Trichoderma spp.* Plantlet survival rate was maximum in *Trichoderma spp.* inoculation, this must be due to the positive interaction among the species and their ability to enhance stress tolerance by protecting them from subsequent 'transplantation shock'.

It was found that the combined use of *Glomus mosseae* and *Pseudomonas fluorescens* caused a greater increase in plant growth of tomato as compared to the individual application⁴⁰. Studies showed that plant-mediated interactions among *Pseudomonas fluorescens*, *Rhizobium leguminosarum*, and AMF on Pea enhanced nodulation by four fold⁴¹. *P. fluorescens* and ectomycorrhizal fungus, *Suillus granulatus*, used as dual inoculants for *Pinus halepensis* showed significantly high number of lateral rootlets as compared to single inoculations⁴². Better root system helped in more nutrient uptake, which resulted in healthy plants with more shoot biomass.

Agrobacterium rhizogenes causing the hairy-root syndrome in dicots is being applied successfully for the promotion of growth of roots and their effective mycorrhization. 100% plant survival was observed after biotization of micropropagated raspberry with *Agrobacterium radiobacter* and *Glomus mosseae*⁶⁴. It was reported that dual inoculation of potato microplants with *P. fluorescens* and mycorrhizal fungi enhanced plant growth and protected against soil-borne potato pathogen *Rhizoctonia solani*⁴³. *Trichoderma sp.* DB11 and *Gliocladium catenulatum* were inoculated in the potting substrates of micro propagated strawberry in order to promote growth and protect against root and collar rot induced by *Phytophthora spp*⁴⁴.

Biotization of micropropagated *Chlorophytum sp.* with the fungus, *Piriformospora indica* and the bacterium, *Pseudomonas fluorescens*, improved plantlet survival rate, growth parameters, field performance, micronutrient acquisition, alkaloid and saponin content⁴⁵. Significantly higher root-shoot biomass was reported in different crops like Maize, *Bacopa*, Poplar with dual inoculation of *P. fluorescens* and *P. indica* inoculation⁶⁵. Maximum colonization of *P. indica* and bacteria in dual inoculated plants attributed the fact that mycorrhizal root tips tend to support slightly higher populations of *Pseudomonas* than non-mycorrhizal root tips, possibly due to the provision of

additional colonization sites or altered root exudation in mycorrhizosphere.

It was reported that acclimatization of in vitro rooted tea plantlets in soil amended with bioinoculants like *Pseudomonas fluorescens*, *Azospirillum brasilense*, and *Trichoderma harzianum*, either individually or in various combinations, promoted plantlet survival⁴⁶.

Table: 1 Effect of Various Bioinoculants on Plant Performance

S.No.	Bioinoculant	Plant	Effect
I	Monomicrobial Interaction - Mycorrhization		
1	Ectomycorrhizae	Pine	Increase in phosphorus uptake and net Photosynthetic rate ^{11,12} .
2	<i>Glomus fasciculatum</i>	Banana	Increase in plant biomass ¹⁴
3	<i>Glomus message</i>	Papaya	Increase in nutritional quality ⁵⁰
4	<i>Glomus diazotrophicus</i>	Sugarcane	Efficient Nitrogen fixation ¹⁶
5	<i>Glomus intradices</i>	Cassava,Grape,Olive	Increase in plant survival and nutrient status ^{51,58}
6	AM fungi	Hazlenut	Resistance towards soil pathogens ¹⁷
7	AM fungi	<i>Ranunulus asiaticus</i>	Earlier flowering ,improved flower production and better rhizome yield ¹⁸
8	<i>Piriformospora indica</i>	<i>C.borivilianum</i>	In vivo rooting ¹⁹
9	<i>Mycorrhizae</i>	Chestnut	Increase protein content and photosynthetic rate ²⁰
10	<i>Piriformospora indica</i>	<i>B.serrata</i>	Increase in plant biomass and phosphorus uptake ^{21,23}
11	<i>Piriformospora indica</i>	<i>T.bellerica</i>	Increase in plant survival and photosynthetic rate ⁵²
12	<i>Piriformospora indica</i>	Barley	Resistance to pathogens and tolerance to salt stress ²⁴
13	<i>Trichoderma viride</i>	Broccoli	Maximum N and P content in roots and shoots ²⁶
14	<i>Trichoderma viride</i>	Neem	<i>In vitro</i> rooting ²⁵
II	Monomicrobial Interaction - Bacterization		
15	<i>Pseudomonas</i>	Potato	Enhance tolerance to transplantation stress ³¹
16	<i>Pseudomonas</i>	Potato	Large root system and better tuber yield ⁵³
16	<i>Pseudomonas</i>	Soyabean	Increase in Plant survival ⁵⁴
17	<i>Pseudomonas</i>	Oregano	Prevents vitrification,Increase in phenolic and chlorophyll content ³²
18	<i>Bacillus subtilis</i>	Tea	Resistance towards pathogens ³⁴
19	<i>Bacillus megaterium</i>	<i>Picrorhiza kurrooa</i>	Induced systemic resistance ³⁵
20	<i>Azospirillum brasilense</i>	Oil Palm	Increase root and shoot biomass ³⁶
21	<i>Azorhizobium caulinodans</i>	Rice	Improved grain weight and nutritional quality ³⁷
III	Multimicrobial Interaction		
22	<i>Glomus mossae</i> + <i>P.fluorescens</i>	Tomato	Increase in yield ⁴⁰
23	<i>P.fluorescens</i> + AMF + <i>Rhizobium leguminosarum</i>	Pea	Increased nodulation by four fold ⁴¹
24	<i>P.fluorescens</i> + Ectomycorrhizae	Pinus	Increase in root and shoot biomass ⁴²
25	<i>A.radiobacter</i> + <i>Glomus mossae</i>	Raspberry	Improved plant survival ⁵⁵
26	<i>P.fluorescens</i> + AMF	Potato	Protection against soil borne pathogens ⁴³
27	<i>Trichoderma</i> + <i>Gliocladium catenulatum</i>	Strawberry	Resistance against root rot and collar rot ⁴⁴
28	<i>P.fluorescens</i> + <i>P.indica</i>	Chlorophytum	Efficient nutrient uptake and increased secondary metabolite content ^{19,45}

29	<i>P.fluorescens</i> + <i>P.indica</i>	Maize,Bacopa,Poplar	High root -shoot biomass ⁵⁶
30	<i>P.fluorescens</i> + <i>A. brasilense</i> + <i>T. harzianum</i>	Tea	Prevented wilting and root rot ⁴⁶
31	<i>P.fluorescens</i> + <i>T.viride</i>	Tea	Increase in phosphorus uptake ⁴⁷
32	<i>P.fluorescens</i> + <i>T.viride</i>	Cotton	Increase in yield ⁴⁹
33	<i>P.fluorescens</i> + <i>T.viride</i>	Soyabean	Resistance against root rot and stem rot ⁴⁸

Root rot or wilting of tissue culture derived plants was not observed in bioinoculant-treated plants, as they possessed relatively higher activities of defense enzymes, including peroxidase and phenylalanine ammonia lyase. The synergistic effect of *T. viride* with *P. fluorescens* was that it can solubilize more P in the soil by producing organic acids⁴⁷. Similarly, their combined effect on growth improvement was also reported by other workers^{48,49}.

The application of multimicrobial biotization requires knowledge and understanding of the compatibility between different beneficial microorganisms in their interaction within the mycorrhizosphere, rhizosphere, and rhizoplane.

CONCLUSION:

Microplants represent an ideal material for developing basic research on microbial biotization and for producing results which can be easily applied in technology transfer. Microplants produced in vitro are usually transplanted into an inert substrate, a relatively simple environment, where the development of the introduced microbe and their impact on plant growth can be easily monitored. A drastic increase in our knowledge about microbial and microbial/plant interactions in the rhizosphere, and particularly of those that are considered beneficial to plant development, is necessary to produce the best results in micropropagation.

In the hardening technique, biotization economizes the production process and simplify the micropropagation technique, which could be adopted at village centres to extend the scientific technology from lab to land. Development of new culture methods allowing the establishment of stable associations between plants and beneficial organisms in vitro and ex vitro and understanding of mechanisms of signal recognition and transduction in plant-microbial associations under different environments are probably the most critical elements of this challenge.

REFERENCES:

1. Lazarovits G & Nowak J. Rhizobacteria for improvement of plant growth and establishment. Hort Science, 1997; 32: 188.
2. Nowak J. Review benefits of in vitro "biotization" of plant tissue cultures with microbial inoculants. In vitro Cell Dev Biol Plant, 1998; 34:122.

3. Rammoorthy V, Viswanathan R, Raguchandar J, Prakasham T & Samiyappan R. Induction of systemic resistance by plant growth promoting rhizobacteria in crop plants against pest and diseases. *Crop protection*, 2001; 20: 1.
4. Hardoim PR, Van Overbeek LS, Berg G, Pirttilä AM, Compant S, Campisano A, Döring M & Sessitsch A, The hidden world within plants: ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiol. Mol. Biol. Rev.* 2015; 79 (3): 293.
5. Trémouillaux-Guiller J, Rohr T, Rohr R & Huss VA, Discovery of an endophytic alga in *Ginkgo biloba*. *Am. J. Bot.* 2002; 89 (5): 727.
6. Müller P, and Döring M, Isothermal DNA amplification facilitates the identification of a broad spectrum of bacteria, fungi and protozoa in *Eleutherococcus* sp. plant tissue cultures. *Plant Cell Tissue Organ Cult*, 2009; 98 (1): 35.
7. Vestberg, M.; Estaun, V. Micropropagated plants, an opportunity to positively manage mycorrhizal activities. In: Gianinazzi, S.; Schuepp, H., eds. *Impact of arbuscular mycorrhizas on sustainable agriculture and natural ecosystems*. Basel: Birkhauser Verlag, 1994:217±226.
8. Morandi D, Gianinazzi S & Gianinazzi-Pearson V, Intérêt de l'endo- mycorrhization dans la reprise et la croissance du Framboisier issu de multiplication végétative in vitro. *Ann. Amélior. Plantes*, 1979; 29: 623.
9. Lovato PE, Gianinazzi-Pearson V, Trouvelot A, Gianinazzi S, The state of mycorrhizas and micropropagation. *Adv. Hort. Sci*, 1996; 10: 46.
10. Allen MF, Smith WK, Moore TS and Christensen M, Comparative water relations and photosynthesis of mycorrhizal and nonmycorrhizal *Bouteloua gracilis* (J. B. K.) Lag ex Steud. *New Phytol*, 1981; 88: 683.
11. Bougher NL, Grove TS, Malajczuk N, Growth and phosphorus acquisition of karri (*Eucalyptus diversicolor* F. Muell.) seedlings inoculated with ectomycorrhizal fungi in relation to phosphorus supply. *New Phytol*, 1990; 114: 77.
12. Rousseau JVD & Reid CPP, Effects of phosphorus and ectomycorrhizas on the carbon balance of Loblolly pine seedlings. *Forest Science*, 1990; 36: 101.
13. DeClerck C, Plenchette & Strullu DG, Mycorrhizal dependency of banana (*Musa acuminata*, AAA group) cultivar. *Plant & Soil*, 1995; 176: 173.
14. Thaker MN & Jasrai YT, Increased growth of micro-propagated banana (*Musa paradisiaca*) with VAM symbiont. *Plant tissue Culture*, 2002; 12: 147.

15. Mandhare, VK, Suryawanshi AV & Jamadagani BM, Occurrence of powdery mildew (*Leveillula taurica*) on chickpea in Maharashtra. *Journal of Maharashtra Agricultural Universities* , 2005; 30: 340.
16. Muthukumarasamy R, Revathi G & Lakshminarasimhan C. Influence of N fertilisation on the isolation of *Acetobacter diazotrophicus* and *Herbaspirillum spp.* from Indian sugarcane varieties. *Biol Fertil Soils* , 1999; 29: 157.
17. Mirabelli C, Tullio M , Pierandrei F& Rea E, Effect of Arbuscular Mycorrhizal Fungi n on micropropagated Hazelnut (*Corylus avellana* L.) plants. *Acta Hort*, 2009; 812: 467.
18. Borriello R, Maccario D, Viglione S, Bianciotto V & Beruto M, Arbuscular mycorrhiza fungi and micropropagation of *Ranunculus asiaticus* L.: a useful alliance. *Acta Hort*, 2017; 1155: 81.
19. Archana Mathur, Ajay Kumar Mathur, Priyanka Verma, Shrawan Yadav, Moti Lal Gupta & Mahendra P. Darokar Biological hardening and genetic fidelity testing of micro-cloned progeny of *Chlorophytum borivilianum*. *African Journal of Biotechnology* , 2008; 7 (8): 1046.
20. Martins A, Casimiro MS & Pais ,Influence of mycorrhization on physiological parameters of micropropagated *Castanea sativa* Mill. *Plants.Mycorrhiza* , 1997; 7: 161.
21. Suthar RK & Purohith SD, Biopriming of micropropagated *Boswellia serrate* Roxb plantlets role of endophytic root fungus *P. indica*. *Ind J Biotech* , 2012; 11: 304.
22. Suthar RK and Purohith SD, Root Colonization and growth enhancement of micropropagated *Terminalia bellarica* Roxb plantlets inoculated with *P. indica* during ex vitro acclimatization. *Int J Plant Dev Biol* , 2008; 2: 133.
23. Chittora M, Suthar RK & Purohit SD, Root colonization and improved growth performance of micropropagated *Terminalia bellarica* roxb. plantlets inoculated with *Piriformospora indica* during ex vitro acclimatization - *Acta Hort*, 2010 : 865.
24. Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, Heier T, Huckelhoven R, Neumann C, von Wettstein D, Franken P & Kogel KH , *Proc Natl Acad Sci USA*, 2005; 102: 13386.
25. Lavanya M, Venkateshwarlu B, Devi BP Acclimatization of neem microshoots adaptable to semi-sterile conditions. *Ind J Biotechnol*, 2009; 8: 218.
26. John RP, Tyagi RD, Prevost D, Brar SK, Pouleur S & Surampalli RY, Mycoparasitic *Trichoderma viride* as a biocontrol agent against *Fusarium oxysporum*f.sp. adzuki and *Pythium arrhenomanes* and as a growth promoter of soybean. *Crop Protection*, 2010; 29 :1452.

27. Gynaeshwar P, Reddy PM, Ladha JK, Nutrient amendments influence endophytic colonization of rice by *Serratia marcescens* IRBG500 and *Herbaspirillum seropedicae* Z67. *J Microbiol Biotechnol*, 2000; 10: 694.
28. Dobbelaere S, Vanderlyden J & Okon Y, Plant growth promoting effects of diazotrophs in the rhizosphere. *Crit Rev Plant Sci* , 2003; 22: 107.
29. Richards J, Induced resistance responses in potato inoculated in vitro with a plant growth promoting pseudomonad bacterium, M.Sc. thesis, Dalhousie University, Halifax, NS, Canada;1997.
30. Yang YS, Wada K, Goto M.;et al., Invitro formation of nodular calli in soya bean induced by co cultivated *Pseudomonas maltophilia*. *Japan. J. Breed*, 1991; 41: 595.
31. Frommel MI, Nowak J & Lazarovits G, Treatment of potato tubers with a growth promoting *Pseudomonas spp*- Plant growth responses and bacterium distributing in the rhizosphere. *Plant Soil*, 1993; 150: 51.
32. Shetty K, Curtis OF, Levin RE.; et al. Prevention of vitrification associated with invitro shoot culture of oregano by *Pseudomonas* spp. *J. Plant Physiol*, 1995; 147: 447.
33. Dunbar C, Utilization of see weed extract and plant growth promoting rhizobacterium in green house production of potato minitubers. M.Sc. thesis, Dalhousie University, Halifax, NS, Canada,1997.
34. Palni LMS & Beg N. Biological hardening of tissue culture raised tea plants through rhizosphere bacteria. *Biotechnol. Lett.*, 2000; 22: 1087.
35. Trivedi P & Pandey A, Biological hardening of micro propagated *Picrorhiza kurroa* Royel ex Benth., an endangered species of medical importance. *World J Microbiol Biotechnol* , 2007;23: 877.
36. Azlin CO, Amir HG, Chai Lai Keng & Zamzuri I, Effect of plant growth promoting rhizobacteria on root formation and growth o f tissue cultured Oil palm .*Biotechnology*, 2007; 6(4): 549.
37. Senthilkumar M, Madhaiyan M, Sundaram SP, Sangeetha H & Kannaiyan S, Induction of endophytic colonization in rice (*Oryza sativa* L.) tissue culture plants by *Azorhizobium caulinodans*. *Biotechnol Lett*, 2008; 30: 1477.
38. Barea JM, Rhizosphere and Mycorrhiza of Field Crops. In: *Biological Resource Management Connecting Science and Policy*. (Ed. Balázs E. et al. Springer, Berlin, Heidelberg) 2000, 81.
39. Von A. State of commercial use of AMF-inoculum in Germany. In: *Arbuscular mycorrhizas in sustainable soil plant systems- Report of 1997 activities, Cost Action* (Ed. Gianinazzi S, & Schuepp H , 821, Iceland) 1998;153.

40. Siddiqui ZA & Mahmood I, Effect of a plant growth promoting bacterium, an AM fungus and soil types on the morph metrics and reproduction of *Meloidogyne javanica* on tomato. *Appl. Soil Ecol*, 1998; 8: 77.
41. Andrade G, De Leij FAAM, Lynch JM, Plant mediated interactions between *Pseudomonas fluorescens*, *Rhizobium leguminosarum* and arbuscular mycorrhizae on pea. *Lett. Appl. Microbiol.*, 1998; 26: 311.
42. Rincon A, Diez BR, Fraile GS, Garcia L, Pascual MF et al, Colonization of *Pinus halepensis* roots by *Pseudomonas fluorescens* and interaction with the ectomycorrhizal fungus *Suillus granulatus*. *FEMS Microbiol Ecol*, 2005; 51: 303.
43. Duffy EM, Hurley EM & Cassells AC, Weaning performance of potato microplants following bacterization and mycorrhization. *Potato Research* , 1999;42: 521.
44. Vestberg M, Kukkonena S, Saaria K, Parikkab P, Huttunenc J, Tainioc L, Devosd N, Weekersd F, Keversd C, Thonartd P, Lemoinee MC, Cordierf C, Alabouvetf C, Gianinazzif S, Microbial inoculation for improving the growth and health of micro propagated strawberry. *Applied Soil Ecology*, 2004; 27: 243.
45. Gosal SK, Karlupia A, Gosal SS, Khhibba IM & Varma A, Biotization with *P. indica* and *P. fluorescens* improves survival rate , nutrient acquisition, field performance and saponin content of micropropagated Chlorophytum species. *Ind J Biotechnol* , 2010; 9: 289.
46. Jibu Thomas, Ajay D & Raj Kumar R, Influence of beneficial microorganisms during in vivo acclimatization of in vitro- derived tea (*Camellia sinensis*) plants . *Plant Cell Tiss Organ Cult* , 2010; 101: 365.
47. Avis TJ, Gravel V, Antoun H & Tweddell RS, Multifaceted beneficial effects of rhizosphere microorganisms on plant health and productivity. *Soil Biology and Biochemistry*, 2008; 40: 1733.
48. Mishra DS, Gupta AK, Prajapati CR and Singh US, Combination of fungal and bacterial antagonists for management of root and stem rot disease of soybean. *Pakistan Journal Botany*, 2011;43: 2569.
49. Shanmugaiah V, Balasubramanian N, Gomathinayagam S, Manoharan PT & Rajendran A, Effect of single application of *Trichoderma viride* and *Pseudomonas fluorescens* on growth promotion in cotton plants. *African Journal of Agricultural Research*, 2009; 4: 1220.
50. Kennedy PM, Lowry JB & Conlan LL, Isolation of grass cell walls as neutral detergent fibre increases their fermentability for rumen micro-organisms. *J. Sci. Food Agric*, 1999; 79 (4): 544.

51. Carlos Luis Carretero , Manuel Cantos , José Luis García , Rosario Azcón & Antonio Troncoso , Growth Responses of Micropropagated Cassava Clones as affected by Glomus Intraradices colonization. Journal of Plant Nutrition, 2009; 32: 261.
52. Vyas S, Nagori R & Purohith SD , Root Colonization and growth enhancement of micropropagated Feronia limonia (L) swingle by *P. indica*. Int J Plant Dev Biol 2008; 2: 128.
53. Bashan Y, Inoculants of plant growth promoting bacteria for use in agriculture, Biotechnol Adv, 1998; 16: 729.
54. Lifshitz R , Kloepper JW et al. Growth promotion of canola seedlings by a strain of *Pseudomonas putida* under gnotobiotic conditions. Can .J. Microbiol, 1987; 33: 390.
55. Gianinazzi S, Oubaha L, Chahbandar M, Blal B, Lemoine MC, Biotization of microplants for improved performance. Acta Horticulturae. , 2003; 625: 165.
56. Verma S, Varma A, Karl-Heinz R, Hasse IA, Kost G et al, *Piriformospora indica* gen. nov. a new root-colonizing fungus, Mycologia USA, 1998; 90: 895.
57. Swada S, Igarashi T & Miyachi S, Effects of nutritional levels of phosphate on photosynthesis and growth studied with single, rooted leaf of dwarf bean. Plant Cell Physiol , 1982; 23: 27.
58. Bhargava BS, Raghupathi HB, Analysis of plant materials for macro and micronutrients. In: Methods of analysis of soils, plants, waters and fertilizer. (Ed. Tandon HLS , Fertilizer Association and Consultation Organisation, New Delhi) 2001;49.