

International Journal of Scientific Research and Reviews

On $(1,2)^*$ - G''' -Regular Space In Bitopological Spaces

Dr. A. ArivuChelvam*

Assistant Professor, Department of Mathematics, MannarThirumalaiNaickercollege,
Madurai, India.

Email: arivuchelvam2008@gmail.com

ABSTRACT

In this paper, we introduce $(1,2)^*$ - g''' -regular space in bitopological spaces. We obtain several characterizations of $(1,2)^*$ - g''' -regular space in some preservation theorems for $(1,2)^*$ - g''' -regular.

KEYWORDS: $(1,2)^*$ - g''' -regular spaces, $(1,2)^*$ - g''' -regular, $(1,2)^*$ - g''' -neighborhood, $(1,2)^*$ - g''' -open set, $(1,2)^*$ - g''' -space.

***Corresponding Author**

Dr. A. ArivuChelvam*

Assistant Professor, Department of Mathematics,

MannarThirumalaiNaickercollege,

Madurai, India.

Email: arivuchelvam2008@gmail.com

1. INTRODUCTION

Using g -closed sets, Munshi¹introduced g -regular in topological spaces. In a similar way, Sheik John²introduced ω -regular using ω -closed sets in topological spaces.

2. SOME DEFINITIONS AND THEOREMS

Definition 2.1

A function $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is called a weakly $(1,2)^*$ - continuous if for each point $x \in X$ and each $\sigma_{1,2}$ -open set V in (Y, σ_1, σ_2) containing $f(x)$, there exists a $\tau_{1,2}$ -open set U containing x such that $f(U) \subseteq \sigma_{1,2}$ - $\text{cl}(V)$.

Definition 2.2

A space (X, τ_1, τ_2) is called a $(1,2)^*$ - gT g''' -space if every $(1,2)^*$ - g - closed set in it is $(1,2)^*$ - g''' -closed.

Definition 2.3

A bitopological space (X, τ_1, τ_2) will be termed symmetric if and only if for x and y in (X, τ_1, τ_2) , $x \in \tau_{1,2}$ - $\text{cl}(y)$ implies that $y \in \tau_{1,2}$ - $\text{cl}(x)$.

Definition 2.4

For a subset A of a bitopological space (X, τ_1, τ_2) , $\tau_{1,2}$ - $\text{cl}_0(A) = \{x \in X : \tau_{1,2}$ - $\text{cl}(U) \cap A \neq \emptyset, U$ is $\tau_{1,2}$ -open set containing $x\}$.

Theorem 2.5

A set A is $(1,2)^*$ - g''' -open if and only if $F \subseteq \tau_{1,2}$ - $\text{int}(A)$ whenever F is $(1,2)^*$ - g -closed and $F \subseteq A$.

Theorem 2.6

The space (X, τ_1, τ_2) is symmetric if and only if $\{x\}$ is $(1,2)^*$ - g -closed in (X, τ_1, τ_2) for each point x of (X, τ_1, τ_2) .

Definition 2.7

Let $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ be a function. Then f is called

- (i) $(1,2)^*$ - g''' -continuous if the inverse image of every $\sigma_{1,2}$ -closed set of Y is $(1,2)^*$ - g''' -closed in X .
- (ii) $(1,2)^*$ - g''' -irresolute if the inverse image of every $(1,2)^*$ - g''' -closed set of Y is $(1,2)^*$ - g''' -closed in X .
- (iii) pre- $(1,2)^*$ - g -open if the image of every $(1,2)^*$ - g -open set of X is $(1,2)^*$ - g -open set in Y .
- (iv) $(1,2)^*$ -open if the image of every $\tau_{1,2}$ -open set of X is $\sigma_{1,2}$ -open in Y .
- (v) $(1,2)^*$ - g -irresolute if the inverse image of every $(1,2)^*$ - g -closed set of Y is $(1,2)^*$ - g -closed in X .
- (vi) $(1,2)^*$ - g''' -closed if the image of every $\tau_{1,2}$ -closed set of X is $(1,2)^*$ - g''' - closed in Y .
- (vii) $(1, 2)^*$ -continuous³ if for each $\sigma_{1,2}$ -open set V of Y , $f^{-1}(V)$ is $\tau_{1,2}$ - open in X .

Theorem 2.8

If $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is bijective, pre-(1,2)*-gs-open and (1,2)*-g'''-continuous, then f is (1,2)*- g'''-irresolute.

Definition 2.9

Let (X, τ_1, τ_2) be a bitopological space. Let x be a point of X and G be a subset of X . Then G is called an (1,2)*- g'''-neighborhood of x (briefly, (1,2)*- g'''-neighborhood of x) in X if there exists an (1,2)*- g'''-open set U of X such that $x \in U \subseteq G$.

3. (1,2)*- α g'''-REGULAR SPACE

We introduce the following definition.

Definition 3.1

A space (X, τ_1, τ_2) is said to be (1,2)*- g'''-regular if for every (1,2)*- g'''-closed set F and each point $x \notin F$, there exist disjoint $\tau_{1,2}$ -open sets U and V such that $F \subseteq U$ and $x \in V$.

Theorem 3.2

Let (X, τ_1, τ_2) be a bitopological space.

Then the following statements are equivalent:

- (i) (X, τ_1, τ_2) is a (1,2)*- g'''-regular space.
- (ii) For each $x \in X$ and (1,2)*- g'''-neighborhood W of x there exists an $\tau_{1,2}$ - open neighborhood V of x such that $\tau_{1,2}\text{-cl}(V) \subseteq W$.

Proof

(i) \Rightarrow (ii). Let W be any (1,2)*- g'''-neighborhood of x . Then there exist a (1,2)*- g'''-open set G such that $x \in G \subseteq W$. Since G^c is (1,2)*- g'''-closed and $x \notin G^c$, by hypothesis there exist $\tau_{1,2}$ -open sets U and V such that $G^c \subseteq U$, $x \in V$ and $U \cap V = \emptyset$ and so $V \subseteq U^c$. Now, $\tau_{1,2}\text{-cl}(V) \subseteq \tau_{1,2}\text{-cl}(U^c) = U^c$ and $G^c \subseteq U$ implies $U^c \subseteq G \subseteq W$. Therefore $\tau_{1,2}\text{-cl}(V) \subseteq W$.

(ii) \Rightarrow (i). Let F be any (1,2)*- g'''-closed set and $x \notin F$. Then $x \in F^c$ and F^c is (1,2)*- g'''-open and so F^c is a (1,2)*- g'''-neighborhood of x . By hypothesis, there exists an $\tau_{1,2}$ -open neighborhood V of x such that $x \in V$ and $\tau_{1,2}\text{-cl}(V) \subseteq F^c$, which implies $F \subseteq (\tau_{1,2}\text{-cl}(V))^c$. Then $(\tau_{1,2}\text{-cl}(V))^c$ is an $\tau_{1,2}$ -open set containing F and $V \cap (\tau_{1,2}\text{-cl}(V))^c = \emptyset$. Therefore, X is (1,2)*- g'''-regular^{4,5,6,7}.

Theorem 3.3

For a space (X, τ_1, τ_2) the following are equivalent:

- (i) (X, τ_1, τ_2) is (1,2)*-normal.

- (ii) For every pair of disjoint $\tau_{1,2}$ -closed sets A and B, there exist $(1,2)^*$ - g''' - open sets U and V such that $A \subseteq U, B \subseteq V$ and $U \cap V = \varnothing$.

Proof

(i) \Rightarrow (ii). Let A and B be disjoint $\tau_{1,2}$ -closed subsets of (X, τ_1, τ_2) . By hypothesis, there exist disjoint $\tau_{1,2}$ -open sets (and hence $(1,2)^*$ - g''' -open sets) U and V such that $A \subseteq U$ and $B \subseteq V$.

(ii) \Rightarrow (i). Let A and B be $\tau_{1,2}$ -closed subsets of (X, τ_1, τ_2) . Then by assumption, $A \subseteq G, B \subseteq H$ and $G \cap H = \varnothing$, where G and H are disjoint $(1,2)^*$ - g''' -open sets. Since A and B are $(1,2)^*$ -gs-closed, by Theorem 2.5, $A \subseteq \tau_{1,2}\text{-int}(G)$ and $B \subseteq \tau_{1,2}\text{-int}(H)$. Further, $\tau_{1,2}\text{-int}(G) \cap \tau_{1,2}\text{-int}(H) = \tau_{1,2}\text{-int}(G \cap H) = \varnothing$.

Theorem 3.4

A $(1,2)^*$ -gT g''' -space (X, τ_1, τ_2) is symmetric if and only if $\{x\}$ is $(1,2)^*$ - g''' -closed in (X, τ_1, τ_2) for each point x of (X, τ_1, τ_2) .

Proof

Follows from Definitions 2.2, 2.3 and Theorem 2.6.

Theorem 3.5

A bitopological space (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular if and only if for each $(1,2)^*$ - g''' -closed set F of (X, τ_1, τ_2) and each $x \in F^c$ there exist $\tau_{1,2}$ -open sets U and V of (X, τ_1, τ_2) such that $x \in U, F \subseteq V$ and $\tau_{1,2}\text{-cl}(U) \cap \tau_{1,2}\text{-cl}(V) = \varnothing$.

Proof

Let F be a $(1,2)^*$ - g''' -closed set of (X, τ_1, τ_2) and $x \notin F$. Then there exist $\tau_{1,2}$ -open sets U_0 and V of (X, τ_1, τ_2) such that $x \in U_0, F \subseteq V$ and $U_0 \cap V = \varnothing$, which implies $U_0 \cap \tau_{1,2}\text{-cl}(V) = \varnothing$. Since $\tau_{1,2}\text{-cl}(V)$ is $\tau_{1,2}$ -closed, it is $(1,2)^*$ - g''' -closed and $x \notin \tau_{1,2}\text{-cl}(V)$. Since (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular, there exist $\tau_{1,2}$ -open sets G and H of (X, τ_1, τ_2) such that $x \in G, \tau_{1,2}\text{-cl}(V) \subseteq H$ and $G \cap H = \varnothing$, which implies $\tau_{1,2}\text{-cl}(G) \cap H = \varnothing$. Let $U = U_0 \cap G$, then U and V are $\tau_{1,2}$ -open sets of (X, τ_1, τ_2) such that $x \in U, F \subseteq V$ and $\tau_{1,2}\text{-cl}(U) \cap \tau_{1,2}\text{-cl}(V) = \varnothing$.

Converse part is trivial.

Corollary 3.6

If a space (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular, symmetric and $(1,2)^*$ -gT g''' - space, then it is $(1,2)^*$ -Urysohn.

Proof

Let x and y be any two distinct points of (X, τ_1, τ_2) . Since (X, τ_1, τ_2) is symmetric and $(1,2)^*$ -gT g''' - space, $\{x\}$ is $(1,2)^*$ - g''' -closed by Theorem 3.4. Therefore, by Theorem 3.5, there exist $\tau_{1,2}$ -open sets U and V such that $x \in U, y \in V$ and $\tau_{1,2}\text{-cl}(U) \cap \tau_{1,2}\text{-cl}(V) = \varnothing$.

Theorem 3.7

Let (X, τ_1, τ_2) be a bitopological space. Then the following statements are equivalent:

- (i) (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular.
- (ii) For each point $x \in X$ and for each $(1,2)^*$ - g''' -neighborhood W of x , there exists an $\tau_{1,2}$ -open neighborhood V of x such that $\tau_{1,2}\text{-cl}(V) \subseteq W$.
- (iii) For each point $x \in X$ and for each $(1,2)^*$ - g''' -closed set F not containing x , there exists an $\tau_{1,2}$ -open neighborhood V of x such that $\tau_{1,2}\text{-cl}(V) \cap F = \emptyset$.

Proof

(i) \Leftrightarrow (ii). It is obvious from Theorem 3.2.

(ii) \Rightarrow (iii). Let $x \in X$ and F be a $(1,2)^*$ - g''' -closed set such that $x \notin F$. Then F^c is a $(1,2)^*$ - g''' -neighborhood of x and by hypothesis, there exists an $\tau_{1,2}$ -open neighborhood V of x such that $\tau_{1,2}\text{-cl}(V) \subseteq F^c$ and hence $\tau_{1,2}\text{-cl}(V) \cap F = \emptyset$.

(iii) \Rightarrow (ii). Let $x \in X$ and W be a $(1,2)^*$ - g''' -neighborhood of x . Then there exists a $(1,2)^*$ - g''' -open set G such that $x \in G \subseteq W$. Since G^c is $(1,2)^*$ - g''' -closed and $x \notin G^c$, by hypothesis there exists an $\tau_{1,2}$ -open neighborhood V of x such that $\tau_{1,2}\text{-cl}(V) \cap G^c = \emptyset$. Therefore, $\tau_{1,2}\text{-cl}(V) \subseteq G \subseteq W$.

Theorem 3.8

The following are equivalent for a space (X, τ_1, τ_2) .

- (i) (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular.
- (i) $\tau_{1,2}\text{-cl}_0(A) = (1,2)^*$ - g''' - $\text{cl}(A)$ for each subset A of (X, τ_1, τ_2) .
- (ii) $\tau_{1,2}\text{-cl}_0(A) = A$ for each $(1,2)^*$ - g''' -closed set A .

Proof

(i) \Rightarrow (ii). For any subset A of (X, τ_1, τ_2) , we have always $A \subseteq (1,2)^*\text{-}g'''\text{-cl}(A) \subseteq \tau_{1,2}\text{-cl}_0(A)$. Let $x \in ((1,2)^*\text{-}g'''\text{-cl}(A))^c$. Then there exists a $(1,2)^*$ - g''' -closed set F such that $x \in F^c$ and $A \subseteq F$. By assumption, there exist disjoint $\tau_{1,2}$ -open sets U and V such that $x \in U$ and $F \subseteq V$. Now, $x \in U \subseteq \tau_{1,2}\text{-cl}(U) \subseteq V^c \subseteq F^c \subseteq A^c$ and therefore $\tau_{1,2}\text{-cl}(U) \cap A = \emptyset$. Thus, $x \in (\tau_{1,2}\text{-cl}_0(A))^c$ and hence $\tau_{1,2}\text{-cl}_0(A) = (1,2)^*\text{-}g'''\text{-cl}(A)$.

(ii) \Rightarrow (iii). It is trivial.

(iii) \Rightarrow (i). Let F be any $(1,2)^*$ - g''' -closed set and $x \in F^c$. Since F is $(1,2)^*$ - g''' -closed, by assumption $x \in (\tau_{1,2}\text{-cl}_0(F))^c$ and so there exists an $\tau_{1,2}$ -open set U such that $x \in U$ and $\tau_{1,2}\text{-cl}(U) \cap F = \emptyset$. Then $F \subseteq (\tau_{1,2}\text{-cl}(U))^c$. Let

$V = (\tau_{1,2}\text{-cl}(U))^c$. Then V is an $\tau_{1,2}$ -open such that $F \subseteq V$. Also, the sets U and V are disjoint and hence (X, τ_1, τ_2) are $(1,2)^*$ - g''' -regular^{12,13}.

Theorem 3.9

If (X, τ_1, τ_2) is a $(1,2)^*$ - g''' -regular space and $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is bijective, pre- $(1,2)^*$ -gs-open, $(1,2)^*$ - g''' -continuous and $(1,2)^*$ -open, then (Y, σ_1, σ_2) is $(1,2)^*$ - g''' -regular.

Proof

Let F be any $(1,2)^*$ - g''' -closed subset of (Y, σ_1, σ_2) and $y \notin F$. Since the function f is $(1,2)^*$ - g''' -irresolute by Theorem 2.8, we have $f^{-1}(F)$ is $(1,2)^*$ - g''' -closed in (X, τ_1, τ_2) . Since f is bijective, let $f(x) = y$, then $x \notin f^{-1}(F)$. By hypothesis, there exist disjoint $\tau_{1,2}$ -open sets U and V such that $x \in U$ and $f^{-1}(F) \subseteq V$. Since f is $(1,2)^*$ -open and bijective, we have $y \in f(U)$, $F \subseteq f(V)$ and $f(U) \cap f(V) = \emptyset$. This shows that the space (Y, σ_1, σ_2) is also $(1,2)^*$ - g''' -regular¹⁴.

Proposition 3.10

If $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is $(1,2)^*$ -gs-irresolute $(1,2)^*$ - g''' -closed and A is an $(1,2)^*$ - g''' -closed subset of X then $f(A)$ is $(1,2)^*$ - g''' -closed in Y .

Proof

Let U be a $(1,2)^*$ -gs-open set in Y such that $f(A) \subseteq U$. Since f is $(1,2)^*$ -gs-irresolute, $f^{-1}(U)$ is a $(1,2)^*$ -gs-open set containing A . Hence $\tau_{1,2}\text{-cl}(A) \subseteq f^{-1}(U)$ as A is $(1,2)^*$ - g''' -closed in X . Since f is $(1,2)^*$ - g''' -closed, $f(\tau_{1,2}\text{-cl}(A))$ is an $(1,2)^*$ - g''' -closed set contained in the $(1,2)^*$ -gs-open set U , which implies that $\sigma_{1,2}\text{-cl}(f(\tau_{1,2}\text{-cl}(A))) \subseteq U$ and $\sigma_{1,2}\text{-cl}(f(A)) \subseteq U$. Therefore $f(A)$ is an $(1,2)^*$ - g''' -closed set in Y ¹⁵.

Theorem 3.11

If $f : (X, \tau_1, \tau^2) \rightarrow (Y, \sigma_1, \sigma_2)$ is $(1,2)^*$ -gs-irresolute $(1,2)^*$ - g''' -closed $(1,2)^*$ -continuous injection and (Y, σ_1, σ_2) is $(1,2)^*$ - g''' -regular, then (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular.

Proof

Let F be any $(1,2)^*$ - g''' -closed set of (X, τ_1, τ_2) and $x \notin F$. Since f is $(1,2)^*$ -gs-irresolute $(1,2)^*$ - g''' -closed, by Proposition 3.3.10, $f(F)$ is $(1,2)^*$ - g''' -closed in (Y, σ_1, σ_2) and $f(x) \notin f(F)$. Since (Y, σ_1, σ_2) is $(1,2)^*$ - g''' -regular and so there exist disjoint $\sigma_{1,2}$ -open sets U and V in (Y, σ_1, σ_2) such that $f(x) \in U$ and $f(F) \subseteq V$. i.e., $x \in f^{-1}(U)$, $F \subseteq f^{-1}(V)$ and $f^{-1}(U) \cap f^{-1}(V) = \emptyset$. Therefore, (X, τ_1, τ_2) is $(1,2)^*$ - g''' -regular¹⁶.

Theorem 3.12

If $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is weakly $(1,2)^*$ -continuous $(1,2)^*$ - g''' - closed injection and (Y, σ_1, σ_2) is $(1,2)^*$ - g''' -regular, then (X, τ_1, τ_2) is $(1,2)^*$ -regular.

Proof

Let F be any $\tau_{1,2}$ -closed set of (X, τ_1, τ_2) and $x \notin F$. Since f is $(1,2)^*$ - g''' - closed, $f(F)$ is $(1,2)^*$ - g''' -closed in (Y, σ_1, σ_2) and $f(x) \notin f(F)$. Since (Y, σ_1, σ_2) is $(1,2)^*$ - g''' -regular

by Theorem 3.5 there exist $\sigma_{1,2}$ -open sets U and V such that $f(x) \in U$, $f(F) \subseteq V$ and $\sigma_{1,2}\text{-cl}(U) \cap \sigma_{1,2}\text{-cl}(V) = \varnothing$. Since f is weakly $(1,2)^*$ -continuous it follows that $x \in f^{-1}(U) \subseteq \tau_{1,2}\text{-int}(f^{-1}(\sigma_{1,2}\text{-cl}(U)))$, $F \subseteq f^{-1}(V) \subseteq \tau_{1,2}\text{-int}(f^{-1}(\sigma_{1,2}\text{-cl}(V)))$ and $\tau_{1,2}\text{-int}(f^{-1}(\sigma_{1,2}\text{-cl}(U))) \cap \tau_{1,2}\text{-int}(f^{-1}(\sigma_{1,2}\text{-cl}(V))) = \varnothing$. Therefore, (X, τ_1, τ_2) is $(1,2)^*$ -regular¹⁷.

REFERENCES:

1. Munshi BM et al. Acta Ciencia Indica, 1986;12: 140-144
2. Tong J et al. On decomposition of continuity in topological spaces, Acta Math. Hungar., 1989; 54:51-55.
3. Ravi O, Lellis Thivagar Met al. On stronger forms of $(1,2)^*$ -quotient mappings in bitopological spaces, Internat. J. Math. Game Theory and Algebra. 2004; 14(6) : 481-492.
4. Reilly IL, Ganster, M et al. Locally closed sets and LC-continuous functions, Internat. J. Math. Sci. 1989;12(3) :417-424.
5. Hatir E, Noiri T, Yuksel S et al. A decomposition of continuity, Acta Math. Hungar., 70, No. 1996;2:145-150.
6. Lellis Thivagar M, Nirmala Mariappan, Jafari S et al. On $(1,2)^*$ - $\alpha\hat{g}$ - closed sets, J. Adv. Math. Studies, 2009;2(2):25-34.
7. Ravi O, Thivagar ML, Joseph Israel, M. Kayathri K et al. Decompositions of $(1,2)^*$ -rg-continuous maps in bitopological spaces, Antarctica J. Math., 2009;6(1): 13-23.
8. Kelly J C et al, Bitopological spaces, Proc. London Math. Soc., 1963; 13:71-89.
9. Kopperman R, Meyer P, Kong T et al. A topological approach to digital topology, Amer. Math. Monthly., 1991; 98:901-917.
10. Ravi O, Abd El-Monsef ME, Lellis Thivagar M et al. Remarks on bitopological $(1,2)^*$ -quotient mappings, J. Egypt Math. Soc. 2008; 16(1) :17-25.
11. Ravi O, Thivagar ML et al. Remarks on λ -irresolute functions via $(1,2)^*$ -sets, Advances in App. Math. Analysis. 2010; 5(1) :1-15.
12. Ravi O, Ekici E, Lellis Thivagar M et al. On $(1,2)^*$ -sets and decompositions of bitopological $(1,2)^*$ -continuous mappings, Kochi J. Math. 2008;3:181-189.

13. Ravi O, KayathriK, Thivagar ML, Joseph Israel M et al. Mildly $(1,2)^*$ -normal spaces and some bitopological functions, *Mathematica Bohemica*. 2010; 135(1):1-15.
14. Ravi O, Pious MissierS, SalaiParkunan T et al. On bitopological $(1,2)^*$ -generalized homeomorphisms, *Int J. Contemp. Math. Sciences*. 2010;5(11):543-557.
15. Ravi O, Thivagar ML, Nagarajan A et al. $(1,2)^*$ - α g-closed sets and $(1,2)^*$ -g α -closed sets (submitted).
16. Ravi O, Thivagar ML, Ekici E et al. Decomposition of $(1,2)^*$ - continuity and complete $(1,2)^*$ - continuity in bitopological spaces, *Analele Universitatii Din Oradea. Fasc. Matematica Tom XV*. 2008; 2:29-37.
17. Ravi O, Pandi A, Pious MissierS, SalaiParkunan T et al. Remarks on bitopological $(1,2)^*$ - $r\omega$ - Homeomorphisms, *Int. J. of Math. Arc*. 2011;2(4) :465-475.