

International Journal of Scientific Research and Reviews

Solution of the fractional partial differential equation by using homotopy analysis method

Gohil V.P^{1*}, Dr. Meher R.K.²

Sardar Vallabhbhai National Institute of Technology, Surat, India

vpgohil0909@gmail.com

ABSTRACT:

The solution of fractional partial differential equations is obtained by using the homotopy analysis method. We also discussed the convergence analysis of the homotopy analysis method about the considered fractional partial differential equation.

KEY WORDS: Fractional partial differential equation, Convergence analysis, Homotopy analysis method.

***Corresponding author**

V.P.Gohil

Sardar Vallabhbhai National Institute of Technology,

Surat, India

Email: vpgohil0909@gmail.com

INTRODUCTION OF HOMOTOPY ANALYSIS METHOD:

This method is proposed by Liao in 1992³⁻⁶. The following differential equations are considered by us for this method,

$$N_i[u_i(x,t)] = 0, i = 1, 2, \dots, n \quad (1)$$

Where N_i considered as a nonlinear operators, (x, t) and $u_i(x, t)$ are pair of independent variables and unknown functions respectively.

The so-called zero-order deformation equations defined by

$$(1-q)L[\phi_i(x,t;q) - u_{i,0}(x,t)] = qc_i N_i[\phi_i(x,t;q)] \quad (2)$$

Where q is an embedding parameter which lies between $[0,1]$, c_i and L are nonzero auxiliary functions and auxiliary linear operator respectively, initial guesses of $u_i(x,t)$ are $u_{i,0}(x,t)$ and $\phi_i(x,t;q)$ are unknown functions.

We have freedom to choose auxiliary objects such as c_i and L in HAM, which is main importance of this method.

When $q = 0$ and $q = 1$ we get by (2),

$$\phi_i(x,t;0) = u_{i,0}(x,t) \quad \text{and} \quad \phi_i(x,t;1) = u_i(x,t)$$

By Taylor's series expansion

$$\phi_i(x,t;q) = u_{i,0}(x,t) + \sum_{m=1}^{\infty} u_{i,m}(x,t) \cdot q^m \quad (3)$$

Where

$$u_{i,m}(x,t) = \left[\frac{1}{m!} \cdot \frac{\partial^m \phi_i(x,t;q)}{\partial q^m} \right]_{q=0} \quad (4)$$

If we choose the auxiliary parameter c_i , the auxiliary functions, the auxiliary linear operator and initial guesses properly than the series equation (3) converges at $q = 1$

$$\phi_i(x, t, ; 1) = u_{i,0}(x, t) + \sum_{m=1}^{\infty} u_{i,m}(x, t) \tag{5}$$

This must be one of the solutions of the original nonlinear equations.

If we differentiate equation (2) m times with respect to the embedding parameter q and the setting $q = 0$ and finally dividing them by $m!$ than we get the so-called m^{th} Order deformation equations like this

$$L[u_{i,m}(x, t) - \chi_m u_{i,m-1}(x, t)] = h_i R_{i,m}(\overline{u_{i,m-1}}) \tag{6}$$

Where

$$R_{i,m}(u_{i,m-1}) = \left[\frac{1}{(m-1)!} \cdot \frac{\partial^{m-1} N_i[\phi_i(x, t, ; q)]}{\partial q^{m-1}} \right]_{q=0} \tag{7}$$

χ_m is characteristic function.

FRACTIONAL DERIVATIVE ACCORDING TO RIEMANN-LIOUVILLE:

$$D^\alpha (t^n) = \frac{d^\alpha}{dt^\alpha} (t^n) = \frac{\Gamma(n+1)}{\Gamma(n+1-\alpha)} t^{n-\alpha} \tag{8}$$

Where, Gamma possesses a standard definite. Inverse fractional differential operator notation is J^α

SOLUTION OF FRACTIONAL KDV EQUATION BY USING THE HOMOTOPY ANALYSIS METHOD: $[0 < \alpha \leq 1]$

Let following be the KDV equation [9]

$$\frac{\partial^\alpha u}{\partial t^\alpha} - 3 \frac{\partial u^2}{\partial x} = 0 \tag{9}$$

With initial condition $u(x, 0) = 3x$

First of all we want to define linear and nonlinear terms like as

$$N(\phi(x, t; q)) = \frac{\partial^\alpha \phi(x, t; q)}{\partial t^\alpha} - 3 \frac{\partial \phi(x, t; q)^2}{\partial x} \tag{10}$$

$$L(\phi(x, t; q)) = \frac{\partial^\alpha \phi(x, t; q)}{\partial t^\alpha} \tag{11}$$

Assume initial approximation

$$u_0(x, t) = 6x + tx^3$$

By using the procedure of Homotopy Analysis Method, the zeroth-order deformation equations for (1) can be written as

$$(1 - q)[\phi(x, t; q) - u_0(x, t)] = qc_0 N[\phi(x, t; q)] \tag{12}$$

For $q = 0$ and $q = 1$, it can be written as

$$\phi(x, t; 0) = u_0(x, t) \quad , \quad \phi(x, t; 1) = u(x, t)$$

The m th order deformation equations can be written as

$$L(u_m(x, t) - \chi_m u_{m-1}(x, t)) = c_0 R_m(u_{m-1}) \tag{13}$$

$$\text{Where } R_m(u_{m-1}) = \frac{\partial^\alpha u_{m-1}}{\partial t^\alpha} - 3 \frac{\partial u_{m-1}^2}{\partial x} \tag{14}$$

The approximate solution of equation (9) can be written as

$$u(x, t) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t) \tag{15}$$

$$\text{Where } u_m(x, t) = \chi_m u_{m-1}(x, t) + c_0 L^{-1}[R_m(u_{m-1})] \tag{16}$$

If we take $m=1$ in (14),

$$\begin{aligned} R_1(u_0) &= \frac{\partial^\alpha u_0}{\partial t^\alpha} - 3 \frac{\partial u_0^2}{\partial x} = x^2 \frac{\Gamma(2)}{\Gamma(2-\alpha)} t^{2-\alpha} - 54x - 54tx^2 - 12t^2 x^3 \\ u_1(x, t) &= c_0 D^{-\alpha} [R_1(u_0)] \\ &= c_0 D^{-\alpha} \left[x^2 \frac{\Gamma(2)}{\Gamma(2-\alpha)} t^{2-\alpha} - 54x - 54tx^2 - 12t^2 x^3 \right] \\ &= c_0 \left[\frac{x^2 t^2}{2} x^3 - 54x \frac{1}{\Gamma(1+\alpha)} t^\alpha - 54x^2 \frac{1}{\Gamma(2+\alpha)} t^{1+\alpha} - 12x^3 \frac{1}{\Gamma(3+\alpha)} t^{2+\alpha} \right] \end{aligned} \tag{17}$$

If we take two terms approximation than we get

$$\begin{aligned} u(x, t) &= u_0(x, t) + u_1(x, t) + \dots \\ &= 3x + tx^2 + c_0 \left[\frac{x^2 t^2}{2} x^3 - 54x \frac{1}{\Gamma(1+\alpha)} t^\alpha - 54x^2 \frac{1}{\Gamma(2+\alpha)} t^{1+\alpha} - 12x^3 \frac{1}{\Gamma(3+\alpha)} t^{2+\alpha} \right] \end{aligned} \tag{18}$$

If we take some special case,

$$\alpha = 1, \alpha = 1, \quad u(x,t) = 3x + tx^2 + c_0 \left[\frac{x^2 t^2}{2} x^3 - 54xt - 27x^2 t^2 - 2x^3 t^3 \right]$$

$$\alpha = \frac{1}{2}, \alpha = \frac{1}{2}, \quad u(x,t) = 3x + tx^2 + c_0 \left[\frac{x^2 t^2}{2} x^3 - 108x \frac{1}{\sqrt{\pi}} t^{\frac{1}{2}} - 36x^2 \frac{1}{\sqrt{\pi}} t^{\frac{3}{2}} - \frac{16}{5} x^3 \frac{1}{\sqrt{\pi}} t^{\frac{5}{2}} \right]$$

CONVERGENCE OF HOMOTOPY ANALYSIS METHOD (HAM):

Theorem:-As long as the series equation (15) is convergent where $u_m(x,t)$ is governed by the m^{th} order deformation equation (13) under (14) must be the solution of (9) ⁵.

Proof: Let the series $u(x,t) = u_0(x,t) + \sum_{m=1}^{\infty} u_m(x,t)$ be convergent.

Then
$$\lim_{m \rightarrow \infty} u_m(x,t) = 0 \tag{19}$$

Now we have

$$\begin{aligned} & \sum_{m=1}^n [u_m(x,t) - \chi_m u_{m-1}(x,t)] \\ &= u_1 + (u_2 - u_3) + (u_3 - u_2) + \dots + (u_{n-1} - u_{n-2}) + (u_n - u_{n-1}) \\ &= u_n \end{aligned}$$

So
$$\sum_{m=1}^{\infty} [u_m(x,t) - \chi_m u_{m-1}(x,t)] = \lim_{n \rightarrow \infty} u_n(x,t) = 0 \tag{20}$$

According to the definition of linear operators, we can write

$$\sum_{m=1}^{\infty} L[u_m(x,t) - \chi_m u_{m-1}(x,t)] = L \left(\sum_{m=1}^{\infty} [u_m(x,t) - \chi_m u_{m-1}(x,t)] \right) = L(0) = 0$$

From the above equation and equation (13)

$$R_m(\overline{u_{m-1}}) = 0, \quad (\because c_0 \neq 0) \tag{21}$$

From (14),

$$\begin{aligned}
& \sum_{m=1}^{\infty} R_m(\overline{u_{m-1}}) \\
&= \sum_{m=1}^{\infty} \frac{\partial^\alpha u_{m-1}}{\partial t^\alpha} - \sum_{m=1}^{\infty} \frac{\partial (u_{m-1})^2}{\partial x} \\
&= \frac{\partial^\alpha \sum_{m=1}^{\infty} u_{m-1}}{\partial t^\alpha} - 3 \frac{\partial \left(\sum_{m=1}^{\infty} u_{m-1} \right)^2}{\partial x} + 3 \frac{\partial \left(\sum_{i,j=1}^{\infty} u_i u_j \right)}{\partial x}
\end{aligned} \tag{22}$$

From (21) and (22), proof of the theorem is completed.

REFERENCES

1. BatihaB. ,Noorani M.S.M. ,Hashim I. Numerical simulations of systems of PDEs by variational iteration method. Phys. Lett. A.2008; 372: 822-829.
2. GohilV.P., MeherR.Homotopy Analysis Method for solving Counter current imbibition phenomena of the time positive fractional type arising in heterogeneous porous media, International Journal of Mathematics & Computation.2017; 28:77-85
3. Liao S.J. On the homotopy analysis method for nonlinear problems. Applied Mathematics and Computation. 2004; 147(2): 499-513.
4. Liao S.J. and Tan Y. A general approach to obtain series solutions of nonlinear differential equations. Studies in Applied Mathematics. 2007; 119: 297-354.
5. Liao S.J. Beyond Perturbation: Introduction to the HomotopyAnalysis Method, Chapman.2004.
6. Liao S.J. Comparison between the homotopy analysis method andhomotopy perturbation method. Appl. Math. Com- put.2005; 169:1186-1194.
7. MeherR, GohilV.P..Application of homotopy analysis method for the solution of cubic boussinesq equation and boussinesq-burger equation.2016; 20:379-396.
8. Wang Z. ,Zou I. ,Zhang H. Applying homotopy analysis method for solving differential difference equation .Phys Lett A. 2007;371: 72-82.
9. Abdulaziz Q. ,Hashim I., Ismail E. S. Approximate analytical solution to fractional modified KdV equations. Mathematical and Computer Modelling. 2009; 49: 136-145.